t. p. opp. belongs with v. 2 part 1;
v2(2) had no sep. t. p.

Individual parts of Willdenow's Species Plantarum were published as follows:

vol. 1 - 1798
vol. 2 - 1800
vol. 3[1] - 1801 - 3[2] - 1803 - 3[3] - 1804
vol. 4[2] - 1806

and not in the years on title-page

Internat. Rules - p.12. 1935

Linneana
QK91
.B512
1797-1830
v. 2, pt. 2
c. 2

CAROLI A LINNÉ
SPECIES PLANTARUM

EXHIBENTES

PLANTAS RITE COGNITAS

AD

GENERA RELATAS

CUM

DIFFERENTIIS SPECIFICIS, NOMINIBUS TRIVIALIBUS
SYNONYMIS SELECTIS, LOCIS NATALIBUS

SECUNDUM

SYSTEMA SEXUALE

DIGESTAS.

Mo. Bot. Garden,
1893

EDITIO QUARTA, POST REICHARDIANAM QUINTA
ADJECTIS VEGETABILIBUS HUCUSQUE COGNITIS

CURANTE

CAROLO LUDOVICO WILLDENOW.

Tomus II. PART II

BEROLINI.
IMPENSIS G. C. NAUK.
1799.

Weiſe haſt du, Natur! der Pflanzen Erzeugung geordnet,
Gütig und weiſe die Kräfte der erdeverſchönernden Pflanzen;
Nicht der Schüler allein der rettenden Göttin Hygea
Kennt ſie die heilenden Kräfte der aromatiſchen Staude,
Fern am Ganges geholt und vom Haupte der Cordilleras,
Oeſter verkannt am Ufer der vaterländiſchen Bäche:
Sicherer weiſs der Wilde die ſchmerzenlindernde Wurzel
Und den geheimeren Stand der ſieberheilenden Rinde:
Aber er kennt ſie auch die tödtenden Gifte der Pflanzen,
Kennt der Euphorbien Kraft und der giftigen Mancinella,
Die den geflügelten Pfeil mit ſchnellerem Tode bewaffnet,
Als das gorgoniſche Scheuſal der ſchlangenbehaarten Echidna. —

Baron von der Lühe.

SPECIES PLANTARUM.

Tomus II. Pars II.

Claſſis XI.

DODECANDRIA.

MONOGYNIA.

927. BOCCONIA *Cor.* o. *Cal.* 2-phyllus, inferus. *Capſ.* 2-valvis, 1-ſperma.

925. ASARUM. *Cor.* o. *Cal.* 3-fidus, ſuperus. *Capſ.* 6-locularis.

955. STERCULIA. *Cor.* o. *Cal.* 5-partit. *Nectar.* ſtaminiferum. *Germen* pedicellatum. *Capſ.* 5.

931. RHIZOPHORA. *Cor.* 4-partita. *Cal.* 4-partitus, inferus. *Sem.* 1, clavatum receptaculo carnoſo.

938. GARCINIA. *Cor.* 4-petala. *Cal.* 4-phyllus, inferus. *Bacca* 8-ſperma, coronata.

943. CRATAEVA. *Cor.* 4-petala. *Cal.* 4-fidus, inferus. *Bacca* 2-locularis, pedicellata.

928. DODECAS. *Cor.* 4-petala. *Cal.* 4-fidus. *Capſ.* 1-locul. 4-valvis.

932. CRENAEA. *Cor.* 4-petala. *Cal.* 4-fidus. *Capſ.* 5-locularis polyſperma.

939. HALESIA. *Cor.* 4-fida. *Cal.* 4-dentatus, ſuperus. *Pericarp.* 4-ſpermum, 4-angulatum.

933. APACTIS. *Cor.* 4-petala. *Cal.* o. *Stam.* 16.

926. TOMEX. *Cor.* 5-petala. *Cal.* o. *Involucrum.* 4-5-phyllum. 5-12-florum. *Bacca* 1-ſperma.

945. EURYA. *Cor.* 5-petala. *Cal.* duplex. *Capſula* 5-locularis.

944. TRIUMFETTA. *Cor.* 5-petala. *Cal.* 5-phyllus, inferus. *Capſ.* 4-locul. 2-ſperma, muricata.

946. PEGANUM. *Cor.* 5-petala. *Cal.* 5-phyllus, inferus. *Capſ.* 3-locul. *Stam.* 15.

954. KLEINHOFIA. *Cor.* 5-petala. *Cal.* 5-phyll. *Nectar* ſtaminiferum. *Germen* pedicellatum. *Capſ.* 5-angularis inflata.

948. NITRARIA. *Cor.* 5-petala. *Cal.* 5-fidus, inferus. *Drupa* 1-ſperma. *Stam.* 15.

941. ARISTOTELIA. *Cor.* 5-petala. *Cal.* 5-partitus. *Bacca* 3-locularis.

937. GRANGERIA. *Cor.* 5-petala. *Cal.* 5-fidus. *Drupa* monoſperma.

936. VATICA. *Cor.* 5-petala. *Anth.* 15-quadriloculares: loculis interioribus brevioribus.

947. HUDSONIA. *Cor.* 5-petala. *Cal.* 3-phyllus, inferus. *Capſ.* 1-locularis. 3-valvis. 3-ſperma.

942. CANELLA. *Cor.* 5-petala. *Cal.* 3-lobus, inferus. *Bacca* 1-locul. 2-ſ. 4-ſperma. *Nectarium* antheriferum.

949. PORTULACA. *Cor.* 5-petala. *Cal.* 2-fidus, inferus. *Capſ.* 1-locul. circumſciſſa.

950. TALINUM. *Cor.* 5-petala. *Cal.* 2-phyllus. *Capſ.* 1-locularis, trivalvis. *Sem* arillata.

951. LYTHRUM. *Cor.* 6-petala. *Cal.* 12-fidus, inferus. *Capſ.* 2-locularis.

952. CUPHEA. *Cor.* 6-petala, inaequalis. *Cal.* 6-dentatus, inaequalis. *Capſ.* 1-locul. ante maturitatem dehiſcens.

953. GINORIA. *Cor.* 6-petala. *Cal.* 6-fidus, inferus. *Capſ.* 1-locul. 4-valvis.

934. BLAKEA. *Cor.* 6-petala. *Cal.* fruct. 6-phyllus: *Floris* ſuperus indiviſus. *Capſ.* 6-locularis. *Antherae* connexae.

929. AGATHOPHYLLUM. *Cor.* 6-petala. *Cal.* truncatus. *Drupa.* monoſperma.

935. BEFARIA. *Cor.* 7-petala. *Stam.* 14. *Bacca.* exſucca, 7-locularis.

930. BASSIA. *Cor.* 8-fida. *Stam.* 16. *Drupa.* 5-ſperma.

940. DECUMARIA. *Cor.* 10-petala. *Cal.* 10-phyllus, ſuperus.

† *Cleome viſcoſa, dodecandra. Chlora dodecandra. Samyda pubeſcens, ſerrulata. Rivina octandra. Paſſerina capitata.*

DIGYNIA.

956. HELIOCARPUS. *Cor.* 4-petala. *Cal.* 4-phyllus. *Capſ.* 2-locul. 1-ſperma, compreſſo-radiata.

957. AGRIMONIA. *Cor.* 5-petala. *Cal.* 5-fidus. *Sem.* 1. ſ. 2.

TRIGYNIA.

958. RESEDA. *Cor.* petalis multifidis. *Cal.* partitus. *Capſ.* 3-locularis, hians.

960. VISNEA. *Cor.* petalis ellipticis. *Cal.* 5-phyll. *Nux.*
959. EUPHORBIA. *Cor.* petalis peltatis. *Cal.* ventricoſus. *Capſ.* 3-cocca.

TETRAGYNIA.

962. APONOGETON. *Cor.* o. *Cal.* o. *Capſ.* 4.
961. CALLIGONUM. *Cor.* o. *Cal.* 5-partitus. *Nux.* monoſperma.

† *Tormentilla erecta. Reſedae aliquot.*

PENTAGYNIA.

963. GLINUS. *Cor.* o, niſi ſetulae. *Cal.* 5-phyllus *Capſ.* 5-locul.
964. BLACKWELLIA. *Cor.* 15-petala. *Cal.* 5-fid. *Capſ.* 1-locul. polyſperma.

† *Reſeda pupuraſcens.*

DODECAGYNIA.

965. SEMPERVIVUM. *Cor.* 12-petala. *Cal.* 12-partitus. *Capſ.* 12.

† *Aliſma cordifolia.*

MONOGYNIA.

925. ASARUM.

Gen. plant. edit. Schreb. n. 801.

Cal. 3. ſ. 4-fidus, germini inſidens. *Cor.* 0. *Capſ.* coriacea, coronata.

1. ASARUM *europaeum.*

A. foliis reniformibus obtuſis binis. *Mat. med.* 119. *Roy. lugdb.* 207. *Gmel. ſib.* 3. *p.* 16. *Mill. dict. n.* 1. *et ic. t.* 53. *Willich. obſ n.* 23. *Scop. carn. n.* 563. *Pollich. pal. n.* 448. *Mönch. haſſ. n.* 388 *Flor dan. t.* 633. *Ludw. ect. t.* 182. *Blackw. t.* 383. *Hoffm. germ.* 161. *Roth. germ. I.* 202. *II.* 518.

A. foliis reniformibus ſubhirſutis. *Hall. helv. n.* 1547.

A. foliis ſubcordatis petiolatis. *Hort. cliff.* 178. *Fl. ſuec.* 392; 421. *Guett. ſtamp.* 198.

Aſarum. *Bauh. pin.* 197. *Cam. epit.* 19.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 2.

Europäiſche Haſelwurz. W.

Habitat in Europae *nemoribus.* ♃ (v. v.)

2. ASARUM *canadenſe.*

A. foliis reniformibus mucronatis. *Mill. dict. n.* 2. *et illuſtr. ſyſt. Thunb. jap.* 190.

Aſarum latifolium.. *Saliſb. prod chap. allert.* 344.

A. foliis ſubcordatis petiolatis. *Gron virg.* 52.

A. canadenſe, mucronato folio. *Moriſ. hiſt.* 3. *p.* 511. *ſ.* 13 *t.* 7. *f* 4

A. canadenſe *Corn. canad.* 24. *t.* 25.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 3.

Canadiſche Haſelwurz. W.

Habitat in Canada. ♃

An varietas praecedentis? ſic ſuadent Folia bina, disſuadent alia.

Folia *utrinque minutiſſime pubeſcentia.* Germen *obſolete trigonum.* Calyx *foliolis inferne incurvis cavis, ſuperne plus minus patentibus, planiuſculis, marginibus revolutis.* Filamenta *vix altitudine ſtigmatum.* Saliſb. l. c.

2. ASARUM *virginicum.*

A. foliis cordatis obtuſis glabris petiolatis. *Gron. virg.* 162. *Mill. dict. n.* 3. *Thunb. jap.* 190.

A. virginianum, piſtolochiae foliis ſubrotundis, cyclaminis more maculatis. *Pluk. alm.* 53. *t.* 78. *f.* 2. *Moriſ. hiſt.* 3. *p.* 511. *ſ.* 17. *t.* 7. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 3.
Virginiſche Haſelwurz. W.
Habitat in Virginia, terra Mariana, Carolina. ♃

*926. TOMEX.

Gen. plant. edit. Schreb. n. 802.

Involucrum 4-5-phyllum. *Cal.* 0. *Cor.* 5-petala. *Nectar* ſquamae 5, inter ſtamina inferiora. *Bacca* 1-ſperma.

*1. TOMEX *japonica.* †
T. floſculis corollatis, foliis ſubtus tomentoſis. W.
Tomex japonica. *Thunb. jap.* 190.
Japaniſcher Tomex. W.
Habitat in Japonia. ♄

Caulis *arboreus excelſus ramoſus, plusquam orgyalis.* Rami *et* ramuli *tomentoſi ultimi angulati.* Folia *alterna petiolata oblonga obtuſa integerrima erecta, parallelo-nervoſa (venoſa? W.); ſupra glabra viridia, ſubtus incano-tomentoſa palmaria, junioribus minoribus.* Petioli *angulati ſtriati tomentoſi pollicares.* Flores *axillares in capitulum collecti pedicellati dioici.* Pedunculi *angulati ſtriati ſolitarii tomentoſi bracteati ſemiunguiculares.* Bracteae *oppoſitae prope baſin minutae tomentoſae.* Pedicelli *intra capitulum breviſſimi quini raro ſex uniflori tomentoſi ſemilineam longi.* Fructus *et* foemineos *flores nunquam vidi.* Thunb. l. c.

*2. TOMEX *Tetranthera.* W.
T. floſculis corollatis, foliis glabris. W.
Tetranthera laurifolia. *Jacq. hort. Schoenb.* 1. *p.* 59. *t.* 113.
Lorbeerblättriger Tomex. W.
Habitat in China. ♄

Rami *teretes flaveſcentes, juniores hirti.* Folia *alterna petiolata oblongo-obovata obtuſa integerrima venoſa glabra, nervo medio ſub lente villoſiuſculo.* Petioli *ſemiunciales hirti.* Pedunculi *laterales in nuda rami junioris parte bi-usque quinqueflori.* Involucrum *tetraphyllum, foliolis ſubrotundis concavis pubeſcentibus multiflorum, flores circiter* 12 *pedi-*

cellatos continens. Pedicelli *florum hirti.* Calyx *nullus.* Corolla *pentapetala alba, petalis apice ſubdenticulatis.* Nectaria *clavata viridia ſtaminibus breviora.* Filamenta 12 - 17 *ſubulata piloſa.* Antherae *oblongae planae, loculis quatuor diſtinctis ſeggregatis.* Germen *oblongum.* Stylus *breviſſimus.* Stigma *ſimplex.* Bacca *globoſa rubra, magnitudine Ribetis rubri, monoſperma.* W.

*3. TOMEX *Sebifera.* W.

T. floſculis apetalis, foliis glabris. W.
Berrya chinenſis. *Klein in Litt.*
Sebifera glutinoſa. *Loureiro cochin.* 783. *
Laurus (*involucrata*) foliis obovatis, umbellis involucratis. *Retz obſ.* 6. *p.* 27. *
Tallow Tree. *Anglor.*
Talgbringender Tomex. W.
Habitat in ſylvis Cochinchinae, Chinae. ♄ (v. ſ.)

Arbor *grandis.* Rami *teretes noduloſi, cortice flaveſcenti nitido tecti.* Ramuli *tenui pube induti.* Folia *ſparſa petiolata obovata obtuſa, vel etiam oblongolanceolata obtuſa, integerrima venoſa glabra.* Petioli *unciales pubeſcentes.* Pedunculi *tetragoni tri-vel quinqueflori pubeſcentes axillares et laterales.* Pedicelli *florum tetragoni valde hirti.* W.

Sequentem deſcriptionem partium fructificantium in recenti planta factam mecum communicavit amicus cariſſimus D. Klein: Involucrum *tetraphyllum, foliolis ovatis concavis extus tomentoſis, includens floſculos apetalos* 8 - 10 *breviter pedicellatos.* Calyx *nullus niſi margo pedicelli integerrimus piloſus floſculorum.* Corolla *nulla.* Nectarium *corpuſcula ad baſin filamentorum ſita.* Filamenta 12 *ad* 15 *filiformia piloſa.* Antherae *oblongae ovatae ſulcatae.* Germen *ovatum ſtaminibus arcte cinctum.* Stylus *filiformis ſtaminibus duplo longior.* Stigma *incraſſatum excavatum margine reflexo.* Bacca *monoſperma globoſa.* W.

927. BOCCONIA.

Gen. plant. edit. Schreb. n. 803.

Cal. diphyllus. *Cor.* 0. *Stylus* bifidus. *Capſ.* 2-valvis, 1-ſperma.

1. BOCCONIA *frute ſcens.*

B. foliis oblongis ſinuatis. W.

Bocconia. *Hort. cliff.* 202. *Iacq. amer.* 146. *Mill. dict. Willich. obſ. n.* 35. *Swartz obſ.* 187. *
B. ramoſa, ſphondyli folio tomentoſo. *Plum. gen.* 55. *Trew ehr. t.* 4.
B. ramoſa, foliis majoribus ſinuatis, racemis terminalibus. *Brown. jam.* 244.
Chelidonium majus arboreum, foliis quercinis. *Sloan. jam.* 82. *hiſt.* 1. *p.* 195. *t.* 125.
Cocoxihvitl. *Hernand. mex.* 158.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 601.
Srauchartige Bocconie. W.
Habitat in Mexico, Iamaica, Cuba, Domingo. ♄ (v. ſ.)
Petala *nulla.* Stamina *ſex.* Willich. l. c. R.

*2. BOCCONIA *cordata.*

B. foliis cordatis ſublobatis. W.
Herzblättrige Bocconie. W.
Habitat in China? ♄

Folia *ſubtus glauca ut in praecedente, ſed ſubrotunda cordata, obſolete lobata, obtuſe dentata nervoſo-venoſa.* Panicula *elongata pedalis, ramis ſimplicibus nec diviſis.* Calyx *albus, ut in praecedente, ſed major.* Stamina *numeravi* 24. Stylus *nullus.* Stigma *bilamellatum ſeſſile.* W.

928. DODECAS.

Gen. plant. ed. Schreb. n. 804.

Cal. 5-fidus corollifer ſuperus. *Petala* 5. *Capſ.* 1-locul. 4-valv. polyſperma calyce coronata.

1. DODECAS *ſurinamenſis.*

D. ſurinamenſis. *Lin. ſuppl.* 245.
Surinamiſche Dodecas. W.
Habitat in Surinamo. C. G. *Dalberg.* ♄

Frutex *Lycii barbari facie laevis ſubtetragonus, ramis oppoſitis brevibus.* Folia *oppoſita obovato-oblonga ſ. cuneata ſubpetiolata laevia obtuſa integerrima.* Stipulae *nullae.* Pedunculi *axillares uniflori breves ſolitarii longitudine calycis.* Bracteae *binae minutae ſub calyce.* Calyx *ante explicationem tetragonus quadrivalvis inferne turbinatus teres.* Petala *calyci inſerta, ad diviſuras.* Stamina *receptaculo inſerta juxta germen, nec calyci.*

*929. AGATHOPHYLLUM.

Gen. pl. ed. Schreb. n. 1754.

Cal. minimus truncatus. *Petala* 6 calyci inſerta. *Drupa* ſubgloboſa. *Nux* ſemiquinquelocul. 1-ſperma. *Nucleus* 5-lobus.

*1. AGATHOPHYLLUM *aromaticum.* W.
Evodia Ravenſura. *Gaert. ſem.* 2. *p.* 101. *t.* 103. *f.* 2.
Ravenſura aromatica. *Sonnerat. itin. ind. ed. germ* 2. *p.* 177. *t.* 127. *
Gewürzhaftes Gutblatt. W.
Habitat in inſula Madagaſcar. ♄
Arbor *craſſa, cortice aromatico rufo, ligno inodoro gravi.* Folia *petiolata alterna obovata obtuſa coriacea integerrima glabra, inferne glauceſcentia venoſa.* Flores *parvi in corymbis terminalibus.* W.

930. BASSIA.

Gen. pl. ed. Schreb. n. 805.

Cal. 4-phyllus. *Cor.* 8-fida, tubo inflato. *Stamina* 16. *Drupa* 5-ſperma.

1. BASSIA *longifolia.*
B. foliis lanceolatis, pedunculis unifloris longiſſimis horizontalibus axillaribus congeſtis. W.
B. longifolia. *Mant.* 563.
Ilippe Malabaris. *Koenig. mſſ.*
Houttuyn Lin. Pfl. Syſt. 3. *p.* 606.
Langblättrige Baſſie. W.
Habitat in Malabaria. ♄ (v. ſ.)
Arbor *procera:* Ramis *ultimis recurvis, craſſiuſculis, griſeo-pubeſcentibus.* Folia, *in extremis ramulis, alterna, approximata, petiolata, ovato-lanceolata, integerrima, acuta, venoſa, nuda, ſemipedalia, decidua.* Petioli *teretiuſculi, breves.* Pedunculi *axillares* 1-5, *ſimplices, filiformes, uniflori erecti: poſt floreſcentiam elongati, penduli.*

*2. BASSIA *latifolia.*
B. foliis ellipticis acutis, pedunculis unifloris nutantibus terminalibus congeſtis. W.
Baſſia latifolia. *Roxb. corom* 1. *p.* 20. *t.* 19. *
Mahwah Tree. *Act. ſoc. Bengal.* 1. *p.* 300.
Breitblättrige Baſſie. W.

Habitat in montibus Indiae *orientalis.* ♄

*3. BASSIA *obovata.*

B. foliis obovatis, pedunculis congeſtis terminalibus unifloris. *Forſt. prod. n.* 200.

Stumpfblättrige Baſſie. W.

Habitat in inſula Tanna. ♄

931. RHIZOPHORA.

Gen. pl. ed. Schreb. n. 806.

Cal. 4-partitus. *Cor.* 4-partita. *Sem.* 1, longiſſimum, baſi carnoſum.

1. RHIZOPHORA *conjugata.*

R. foliis ovato-oblongis obtuſiuſculis integerrimis, calycibus ſeſſilibus, fructibus cylindraceo-ſubulatis. *Fl. zeyl.* 181.*

Anonyma. *Herm. pict.* 279. *Burm. ind.* 108. R.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 672.

Gepaarter Wurzelbaum. W.

Habitat in India. ♄

2. RHIZOPHORA *gymnorhiza.*

R. foliis ovato-lanceolatis integerrimis, radice terrae ſuperimpoſita. *Sp. pl.* 634.

R. calycum laciniis perſiſtentibus patentibus verſus fructum incurvatis. *Wachend. ult.* 89.

Mangium celſum. *Rumph. amb.* 3. *p.* 102. *t.* 68.

Candel. *Rheed mal.* 6. *p.* 27. *t.* 31. 32. *Raj. hiſt.* 1769. *Burm. ind. p.* 108.

Houttuyn Lin. Pfl. Syſt. 5. *p.* 673.

Hoher Wurzelbaum. W.

Habitat in Indiae *ripis ſalſis.* ♄

3. RHIZOPHORA *Candel.*

R. foliis obtuſis, pedunculis bigeminatis folio longioribus, fructibus ſubulatis. *Sp. pl.* 634.

Tſierou-Candel. *Rheed. mal.* 6 *p.* 63. *t.* 25. *Raj. hiſt.* 1770. *Burm. ind.* 108.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 675.

Pfriemenfrüchtiger Wurzelbaum. W.

Habitat in Indiae *ſalſis.* ♄

4. RHIZOPHORA *Mangle.*

R. foliis acutis, fructibus ſubulato-clavatis. *Sp. pl.* 634.

R. pedunculis bifidis trifidisque, fructibus ſubulato-clavatis. *Iacq. amer.* 141. *t.* 89.

R. ſegmentis calycum perſiſtentibus reflexis, fructu acuminato. *Wack. ultr.* 90.
R. utrinque brachiata, foliis elliptico-ovatis: ſummis ramis diſpoſitis. *Brown jam.* 211.
Mangle arbor pyrifolia, fructu oblongo tereti, ſummis ramis radicoſa. *Pluk. alm.* 241. *t.* 204. *f.* 3.
Candela americana foliis laurinis. *Catesb. car.* 2. *p.* 63. *t.* 63.
Mangle aquatica, foliis ſubrotundis et punctatis. *Plum. gen.* 13.
Mangle pyri foliis, cum ſiliquis longis ficui indicae affinis. *Bauh hiſt.* 1. *p.* 415. *Sloan. jam.* 155. *hiſt.* 2. *p.* 63. *Rai. hiſt.* 1772.
Mangium calendarium *Rumph. amb.* 3. *p.* 108. *t.* 71. 72.
Peekandel. *Rheed.* 6 *p.* 91. *t.* 34. *Raj. hiſt.* 1770.
Margue Guapariba. *Piſ. braſ. l.* 4. *c.* 87. *R.*
Houttuyn Lin. Pfl. Syſt. 1. *p.* 675.
Gemeiner Wurzelbaum. W.
Habitat in paludibus Caribaearum, Malabariae. ♄ (v. ſ.)
Folia *oppoſita. petiolata, coſtata.* Cal. 4-*fidus. Petala* 4. Stamina 8.

5. RHIZOPHORA *cylindrica.*
R. fructibus cylindricis obtuſis. *Sp pl.* 635:
Karil-Candel. *Rheed. mal.* 6. *p.* 59. *t.* 33. *Raj. hiſt* 1770.
Mangium minus. *Rumph. amb.* 3. *p.* 106. *t.* 69. *Burm. ind.* 108.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 682.
Walzenfrüchtiger Wurzelbaum. W.
Habitat in uliginoſis Malabariae. ♄

*932. CRENAEA.

Gen. plant. edit. Schreb. n. 807.

Cal. 4-fidus corollifer. *Petala* 4. *Capſ.* 5-locularis polyſperma.

*1. CRENAEA *maritima.*
C. maritima *Aubl. guj* 1. 525. *t.* 209. *
Meerſtrands Kerbling. W.
Habitat in maritimis Gujanae. ♃
Caules *e radice plures tetragoni nodoſi.* Folia *ſubpetiolata oppoſita obovato-lanceolata integerrima venoſa glabra.* Flores *albi parvi in corymbis*

pauciſloris axillaribus. Bracteae *binae lanceolatae parvae oppoſitae in quolibet pedunculo.*

*933. APACTIS.

Gen. plant. ed. Schreb. n. 808.

Cal. 0. *Petala* 4 crenata inaequalia. *Germen* ſuperum. *Fructus.* - - - -

*1. APACTIS *japonica.* †
A. japonica. *Thunb. jap.* 191. *
Japaniſche Apactis. W.
Habitat in Japonia. ♄
Arbor. Folia *alterna petiolata ovata ſerrata baſi integerrima glabra.* Flores *in ultimis ramulis racemoſi.* Racemi *ſaepe ovati, excreſcentes magis oblongi.* Pedunculus *et* pedicelli *villoſo-ſcabri.* Thunb. l. c.

934. BLAKEA.

Gen. plant. ed. Schreb. n. 810.

Cal. inferus 6-phyllus: ſuperus integer. *Pet.* 6. *Capſ.* 6-locularis, polyſperma.

1. BLAKEA *trinervia.*
B. bicalyculata, foliis trinerviis, transverſim ſubtiliſſime ſtriatis. *Lin. ſuppl.* 246. *Swartz obſ.* 188. *
B. foliis ellipticis trinerviis, nitidis, floribus lateralibus. *Brown jam.* 323. *t* 35.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 607.
Dreynervige Blakie. W.
Habitat in Iamaica. ♄
Folia *oblongo-ovata, petiolata, integerrima, coriacea, oppoſita, trinervia: nervis ſubtus protuberantibus nigricantibus.* Flores *oppoſiti, ſolitarii.*

2. BLAKEA *triplinervia.*
B. ecalyculata, foliis triplinerviis, pedunculis axillaribus ſubtrifloris. *Vahl. ſymb.* 3. *p.* 61.
B. ecalyculata, follis tripilnerviis. *Lin. ſuppl.* 246.
B. foliis triplinerviis, coſtatis. *Amoen. acad p.* 257.
B. (*quinquenervia*) foliis ovato-oblongis acuminatis quinquenerviis. *Aubl guj.* 1. *p.* 525. *t.* 210. *
Dreyfachrippige Blakie. W.
Habitat in ſylvis Gujanae. ♄

Arbor 16-*pedalis.* Folia *oppoſita petiolata, triplinervia ſpithamaea, ſubtus itidem coſtata et nervo marginali.* Pedunculi *triflori.* Flores *magnitudine Punicae diſtincti, absque perianthio infero.* Perianthium *campanulatum* 3-5-*fidum coriaceum receptaculo a peripheria ad centrum ſtriato perſiſtens.* Petala *circiter* 7 *ovato-oblonga calyci inſerta ſeſſilia lutea, intus alba carnoſa.* Filamenta *circites* 12 *ſ.* 15 *ſubcylindrica calyci inſerta, deſinentia in antheras.* Antherae *ovatae compreſſae parallelae incurvae, dorſo planiuſcutae corolla breviores, quaſi truncatae, non concatenatae, e filamentis continuatae et non filamentis inſidentes.* Germen *inferum.* Stylus *clavatus teres longitudine petalorum deflexus ad latus inferius corollae, extra circulum ſtaminum.* Stigma *capitatum ſtriatum.* Bacca *ſubrotunda multilocularis coronata calyce, diſco concavo, annulo crenato, centro umbilicato.* Semina *minuta.* Fructus *luteus ſapidus. Raro flos in Herbario conſervari poteſt, quia frangatur.*

*3. BLAKEA *pulverulenta.*

B. ecalyculata, foliis triplinerviis, racemo terminali. *Vahl ſymb.* 3. *p.* 61. *

Beſtaubte Blakie. W.

Habitat in America *meridionali.* ♄

Folia *elliptica acuminata integerrima, utrinque glabra, exceptis nervis ſubtus.* Vahl. l. c.

935. BEFARIA.

Gen. plant. edit. Schreb. n. 811.

Cal. 7-fidus. *Petala* 7. *Stam.* 14. *Bacca* 7-locularis, polyſperma.

1. BEFARIA *reſinoſa.*

B. foliis ovatis, floribus congeſtis. *Lin. ſuppl.* 246.
Icon. *Mutis amer. v.* 1. *t.* 8.

Harzige Befarie. W.

Habitat in Nova Granada. *Mutis.* ♄

Arbor *ramis proliferis, cortice rudi, tenello ſubpubeſcente.* Folia *ſparſa ſubpetiolata ovata laevia perennia integerrima, longitudine pollicaria.* Flores *e ſummitate ramorum congeſti: pedunculis pubeſcentibus unifloris, in racemis minime elongatis.* Corolla

purpurea valde reſinoſa ſ. viſcoſa. Stamina *prope baſin pubeſcentia.*

2. BEFARIA *aeſtuans.*

B. foliis lanceolatis, floribus racemoſis. *Lin. ſuppl.* 247.
B. aeſtuans. *Mant.* 242.
Icon. *Mutis amer. v.* 1. *t.* 7.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 608.
Heiſse Befarie. W.
Habitat in Nova Granada. *Mutis.* ♄

Frutex *biorgyalis: Ramis teretiuſculis, patentibus.* Folia *ſaepius alterna, conferta obovata ſ. oblonga, obtuſa, integerrima, avenia, nitida, ſubtus tomentoſa.* Petioli *breviſſimi, planiuſculi.* Flores *axillares, pauci; terminales plurimi.* Pedunculi *racemoſi: Pedicellis teretibus, hirſutis.* Corollae *incarnatae.* Plantam *non ipſe vidi.* Mant.

Similis B. reſinoſae, altior. Folia *ſimilia B. reſinoſae, ſed lanceolata.* Pedunculi *racemoſi, pedicellis longitudine floris nec ut in B. reſinoſa brevioribus.* Suppl.

Befariae ſaporem peculiarem amarum habent. Affinitas cum Rhododendris. Suppl.

936. VATICA.

Gen. plant. ed. Schreb. n. 812.

Cal. 5-fidus. *Petala* 5. *Antherae* 15, ſeſſiles, 4-loculares.

1. VATICA *chinenſis.*

V. chinenſis. *Mant.* 242. *Smith. ic. ined.* 1. *p.* 36. *t.* 36. *
Houttuyn Lin. Pfl. Syſt. 3. *p.* 609.
Chineſiſche Vatica. W.
Habitat in China. ♄

Statura *accedit ad* Toberam *Kaempf. amoen.* 797. *aut Citrum.* Rami *ſtriati, ſ. angulati, ſubtomentoſi.* Folia *alterna, petiolata, cordato-ovata, integerrima, utrinque glabra, venoſa, alterne nervoſa, ſpithamaea, latitudine palmaria.* Flores *paniculati, praecipue terminales, magnitudine Citri.*

*937. GRANGERIA.

Juſſ. gen. ed. Uſt. 376.

Cal. 5-fidus. *Petala* 5. *Stamina* 15. *Drupa* ſubtriquetra. *Nux* triquetra oſſea monoſperma.

*1. GRANGERIA *borbonica.*

G. borbonica. *Lamarck. encyclop.* 3. *d.* 20. * *illuſt. gen. t.* 427.

Borboniſche Grangerie. W.

Habitat in inſula Borbonia. ♄

Arbor *magnitudine Quercus cortice albicante.* Folia *alterna petiolata ovata integerrima glabra venoſa.* Stipulae *binae ſubulatae.* Racemi *ſesquiunciales terminales et axillares.* Flores *parvi albi.* Lamarck. l. c.

938. GARCINIA.

Gen. plant. edit. Schreb. n. 813.

Cal. 4-phyllus, inferus. *Petala* 4. *Bacca* 8-ſperma, coronata *Stigmate* peltato.

1. GARCINIA *Mangoſtana.*

G. foliis ovatis, pedunculis unifloris. *Hort. cliff.* 182. *Mill. dict.*

Mangoſtans. *Garc. act. angl.* 431. *t.* 1. *Bont. jav.* 115.

Mangoſtana. *Rumph. amb.* 1. *p.* 132. *t.* 43.

Laurifolia javanenſis. *Bauh. pin.* 461. *Raj. hiſt.* 1662.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 687.

Wohlſchmeckende Mangoſtane. W.

Habitat in Java. ♄

2. GARCINIA *celebica.*

G. foliis lanceolatis, pedunculis trifloris. *Sp. pl.* 635.

Mangoſtana celebica. *Rumph. amb.* 1. *p.* 134. *t.* 44.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 689.

Celebiſche Mangoſtane. W.

Habitat in India. ♄

3. GARCINIA *Cambogia.*

G. foliis ellipticis acutis, floribus ſolitariis terminalibus ſubſeſſilibus. W.

Mangoſtana Cambogia. *Gaert. ſem.* 2. *p.* 106. *t.* 105.

Cambogia Gutta. *Sp. pl.* 728. *Amoen. acad.* 1. *p.* 402. *Fl. zeyl.* 195. *Mat. med.* 135. *Blackw. t.* 392.

Coddam-pulli. *Rheed. mal.* 1. *p.* 41. *t.* 24.

Carcapuli. *Bauh. hiſt.* 1. *p.* 105. *Raj. hiſt.* 1661.
Carcapuli Acoſtae, fructu malo aureo ſimili. *Pluk. alm.* 81.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 5.
Guttabringende Mangoſtane. W.
Habitat in India. ♄ (v. ſ.)

4, GARCINA *cornea.*
G. foliis lanceolatis aveniis, pedunculis unifloris cernuis. *Syſt. veg.* 368.
Lignum corneum. *Rumph. amb.* 2. *p.* 55. *t.* 30.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 690.
Hornartige Mangoſtane. W.
Habitat in India. ♄
Obſ. Mangium ſylveſtre *Rumph. amb.* 3. *p,* 77. *t.* 30. *diſtincta ſpecies videtur.*

939. HALESIA.

Gen. plant. edit. Schreb. n. 814.

Cal. 4-denatus, ſuperus. *Cor.* 4-ſida. *Nux* 4-angularis, 2-ſperma.

1. HALESIA *tetraptera.*
H. foliis ovatis acuminatis, venis ſubtus piloſis, alis fructus aequalibus. *Willd. arb.* 137. *
H. foliis lanceolato-ovatis: petiolis glanduloſis. *Sp. pl.* 636. *Mill. dict. n.* 1.
H. foliis ovatis acuminatis argute ſerratis, fructu tetraptero. *Cavan. diſſ.* 6. *p.* 338. *t.* 186. *
H. fructibus membranaceo-quadrangulatis. *Ellis act. angl. v.* 51. *p.* 931. *t.* 22. *f. A.*
Frutex padi foliis ſerratis, floribus monopetalis albis campaniformibus, fructu craſſo tetragono. *Catesb. car.* 1. *p.* 64. *t.* 64.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 610.
Vierflüglige Haleſie. W.
Habitat in Carolina. ♄ (v. v.)

2. HALESIA *diptera.*
H. foliis oblongo-ovatis obtuſe acuminatis, utrinque viridibus, ſubtus molliſſimis, alis fructus alternatim majoribus. *Willd. arb.* 138. *
H. foliis ovatis: petiolis laevibus. *Sp. pl.* 636. *Mill. dict. n.* 2.
H. foliis ovatis acutis ſerratis, fructibus bialatis, pedunculis elongatis. *Cavan. diſſ.* 6. *p.* 338. *t.* 187. *
H. fructibus alatis. *Ellis act. angl. v.* 51. *t.* 931. *f. B.*

Houttuyn Lin. Pfl. Syſt. 3. *p.* 611.
Zweyflügelige Haleſie.
Habitat in Carolina. *D. Garden.* ♄ (v. v.)

Folia *hujus ſextuplo majora praecedentis, nec ſubtus ſubtomentoſa.* Fructus *mucronatus alis duabus oppoſitis magnis; duabus minutis. Ordine naturali proxima Styraci.*

Nullas glandulas *in Haleſiae ſpecierum petiolis obſervavi. Differt haec ſecundum obſervationes meas a praecedente*: foliis *duplo majoribus, apice vix acuminatis, ſubtus tactu molliſſimis a pilis parvis nudo oculo inviſibilibus*, fructus *alis inaequalibus binis majoribus, mucrone fructus, qui eſt ſtylus perſiſtens, breviore obtuſo.* W.

940. DECUMARIA.

Gen. pl. ed. Schreb. n. 815.

Cal. 10-phyllus, ſuperus. *Petala* 10. *Capſ.* 8-ſ. 9-locularis polyſperma.

1. DECUMARIA *barbara.*
D. foliis omnibus ovatis, caule ſcandente. W.
Decumaria barbara. *Sp. pl.* 1663.
Tinus flore pleno. *Willich. obſ.* 1762. *p.* 40. *Schreb. Nov. Act. Upſal. vol.* 1. *p.* 87.
Cluſia foliis venoſis. *Fabric. helmſt.* 2. *p.* 293.
Cluſia minor. *Kniph. cent.* 2. *n.* 15.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 612.
Gemeine Decumarie. W.
Habitat in Carolina. ♄ (v. v.)

Rami *inconditi, geniculis nodoſis radicantes.* Folia *oppoſita, petiolata, obovata, coriacea, venoſa, verſus baſin remote ſerrata.* Stipulae *nullae.* Gemmae *pubeſcentes.* Panicula *corymboſa, terminalis.* Flores *Tiliae ſimiles, albidi, odoratiſſimi.* Calyx *dentibus* 8 *ſ.* 16, *ſuperus.* Fructus *in Europaeorum hortis ignotus. An dioica?*

Caulis *tri. ſ. quadripedalis ſcandens.* W.

*2. DECUMARIA *ſarmentoſa.*
D. foliis inferioribus rotundatis, ſuperioribus ovato-lanceolatis, caule ſarmentoſo. *Boſc in Act. hiſt. nat. Pariſ.* 1. *p.* 76. *t.* 4. *
Forſythia (*ſcandens*) caule geniculato altiſſime ſcandente radicante, foliis ſimplicibus oppoſitis petiolatis acumi-

nato-ovatis apice dentatis, ſupra laevibus, pagina inferiore marginibus petiolisque ſubtomentoſis, floribus albis odoratis, bracteis ſubulatis in corymbis trichotomis terminalibus. *Walt. carol.* 154.
Kletternde Decumarie. W.
Habitat in Carolinae *uliginoſis.* ♄

*941. ARISTOTELIA.

Gen. pl. ed. Schreb. n. 816.

Cal. 5-phyllus. *Petala* 5. *Styl.* trifidus. *Bacca* trilocularis. *Sem.* bina.

*1. ARISTOTELIA *Macqui.*
A. Macqui *l'Herit ſtirp.* 1. *p.* 31. *t.* 16.* *Ait. kew.* 2. *p.* 146. *Willd. arb.* 27.*
Chiliſche Ariſtotelie. W.
Habitat in Chili. ♄ (v. v.)
Frutex *octopedalis ramoſus.* Folia *oppoſita ſempervirentia ovata nitida ſerrata trinervia venoſa.* Flores *albi in racemis axillaribus* 2-*ſ.* 4-*floris.* W.

942. CANELLA.

Gen. plant. edit. Schreb. n. 817.

Cal. 3-lobus. *Petala* 5. *Antherae* 16, adnatae, *Nectario* urceolato. *Bacca* unilocularis 2-ſ. 4-ſperma.

1. CANELLA *alba.*
C. alba. *Murr. ſyſt. veg.* 443. *Swartz. obſ.* 190.* *Swartz. in Act. ſoc. Lin. Lond.* 1. *p.* 96. *t.* 8.*
Winterania Canella. *Sp. pl.* 2. 636. *Lin. ſuppl.* 247.
Winterania. *Hort. cliff.* 488. *Mat. med.* 119.
Winteranus cortex. *Blackw. t.* 206.
Laurus foliis enerviis obovatis obtuſis. *Sp. pl.* 1. *p.* 371.
Canella foliis oblongis obtuſis nitidis, racemis terminalibus. *Brown. jam.* 215. *t.* 37. *f.* 3.
Pſeudo-Caſſia cinnamomea aromatica. *Bauh. pin.* 409.
Caſſia lignea jamaicenſis, laureolae foliis ſubcinereis cortice piperis modo acri. *Pluk. alm.* 89. *t.* 81. *f.* 1.
Caſſia cinnamomea ſ. Cinnamomum ſylveſtre barbadenſium. *Pluk. alm.* 89. *t.* 161. *f.* 7.
Arbor baccifera laurifolia aromatica, fructu viridi caly-

culato racemoſo. *Sloan. jam.* 165. *hiſt.* 2. *p.* 87. *t.* 191. *f.* 2. *Catesb. carol.* 2. *p* 50. *t.* 50.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 691.
Weiſser Canellbaum. W.
Habitat in interioribus ſylvis Indiae *occidentalis.* ♄

943. CRATAEVA.

Gen. plant. edit. Schreb. n. 818.

Cor. 4-petala. *Cal.* 4-ſidus. *Bucca* 1-locularis, polyſperma.

1. CRATAEVA *gynandra.*
C. inermis, foliis integerrimis, floribus gynandris. *Sp. pl.* 636. *Swartz. obſ.* 191.*
C. arborea triphylla, foliis ovatis glabris, racemis terminalibus. *Brown. jam.* 246.
Anona trifolia, flore ſtamineo, fructu ſphaerico ferrugineo ſcabro minori, allii odore. *Sloan. cat.* 206. *hiſt* 2. *p.* 170. *Raj. hiſt.* 3. *p.* 78.
Arbor americana triphylla, numeroſis ſtaminibus purpureis apicibus praeditis, floris umbilicum occupantibus. *Pluk. phyt.* 147. *f.* 6.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 694.
Weſtindiſche Crataeve. W.
Habitat in Jamaicae *fruticetis aridis.* ♄
Differt *a. C. Tapia foliis tenuibus nec craſſis. Petalis lanceolatis nec ovatis; imprimis vero quod columna longitudine calycis attollat ſtamina, ut in Cleome gynandra. Piſtillum in meis non aderat.*

2. CRATAEVA *Tapia.*
C. foliolis ovatis acuminatis, petalis ovato-ſubrotundis obtuſis germinibus globoſis. *Vahl ſymb.* 3. *p.* 61.*
C. inermis, foliis integerrimis: foliolis lateralibus baſi antica brevioribus. *Sp. pl.* 637. *Mill. dict. n.* 1.
C. inermis. *Fl. zeyl.* 211. *Sp. pl.* 1. *p.* 444. *Hort. cliff.* 484.
Apioſcorodon ſ. Arbor americana triphyllos, allii odore, poma ferens. *Pluk. alm.* 34. *t.* 157. *f.* 7.
Malus americana trifolia, fructu pomi aurantii inſtar colorato. *Comm hort* 1. *p.* 129. *t.* 67.
Tapia arborea triphylla. *Plum. gen.* 22. *t.* 21. *R.*
Tapia *Marggr. braſ.* 89. *Piſ. braſ.* 68. *t.* 69. *R.*
Houttuyn Lin. Pfl. Syſt. 1. *p.* 694.

Spitzblättrige Crataeve. W.
Habitat in Indiis. ♄ (v. ſ.)

Flos *etiam in hac, ut in* C. *gynandra, eſt germine ſtipitato more Cleomes gynandrae.*

*3. CRATAEVA *obovata.*

C. foliis petaliſque obovatis, germine oblongo. *Vahl. ſymb.* 3. *p.* 61. *

Stumpfblättrige Crataeve. W.
Habitat in Madagaſcar. ♄

An modo varietas C. Tapiae? vix credo, differt: figura *foliolorum et petalorum*, foliolis *baſi ſubaequalibus*, petalis *longius unguiculatis*, calycis *laciniis oblongis*, filamentis *longioribus*, germine *oblongo. Ceterum habitus reliquorum.* Vahl. l. c.

*4. CRATAEVA *religioſa*

C. foliolis petaliſque lanceolato-ellipticis utrinque acutis. *Vahl. ſymb.* 3. *p.* 62. *

C. (*religioſa*) inermis foliolis ſubaequalibus. *Forſt. prodr. n.* 203. *pl. eſcul.* 45 *

Niirvala. *Rheed. mal.* 3. *p.* 49. *t.* 42. *Raj. hiſt.* 1644.

Lanzettenblättrige Crataeve. W.
Habitat in India *orientali inque* Societatis *inſulis.* ♄

5. CRATAEVA *Marmelos.*

C. ſpinoſa, foliis ſerratis. *Fl. zeyl.* 212. *Mill. dict. n.* 2.

Cucurbitifera trifolia ſpinoſa medica, fructu pulpa cydonii aemula. *Pluk. alm.* 125. *t.* 170. *f.* 5.

Cydonia exotica. *Bauh. pin.* 425.

Bilanus. *Rumph. amb.* 1. *p.* 197. *t.* 81.

Covalam. *Rheed. mal.* 3. *p.* 37. *t.* 37. *Burm. ind.* 109.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 696.

Dornige Crataeve. W.
Habitat in India. ♄ (v. ſ.)

Calyx 5-*fidus.* Petala *nulla? Millero* 5. Stamina 60. An *diſtincti Generis?*

Folia *omnium ſpecierum ſunt ternata.* W.

944. TRIUMFETTA.

Gen. pl. ed. Schreb. n. 819.

Cor. 5-petala. *Cal.* 5-phyllus. *Capſ.* hiſpida, in 4 diſſiliens.

1. TRIUMFETTA *Lappula.*

T. foliis baſi emarginatis, floribus ecalyculatis. *Syſt. nat.* 13. *p.* 327.

T. floribus calyce deſtitutis. *Iacq. ame* 146.
T. foliis baſi emarginatis. *Hort. cliff* 10.* *Hort. upſ.* 148 *Roy. lugdb.* 476. *Mill. dict.*
T. fructu echinato racemoſo. *Plum. gen.* 50. *ic.* 255.
Lappula bermudenſis althaedoides ſpicata, fructu orbiculari majore. *Pluk. alm.* 206. *t.* 245. *f.* 7.
Agrimonia lappacea inodora, folio ſubrotundo dentato. *Sloan. jam.* 92. *hiſt.* 1. *p.* 211. *Raj. ſuppl.* 247.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 5.
Kelchloſe Triumfette. W.
Habitat in Jamaica, Braſilia, Bermudis. ♄
Singulare *quod calyx nullus ſit.*

*2. TRIUMFETTA *glanduloſa.*

T. floribus completis, foliis ovato-lanceolatis ſubtus tomentoſo-incanis. *Vahl. ſymb.* 3. *p.* 62.
Triumfetta glanduloſa. *Forſk. cat. arab. n.* 297.
Drüſige Triumfette, W,
Habitat in Arabia *felici,* India. ♄ (v. ſ.)
Rami *lignoſi teretes villoſi.* Folia *petiolata alterna frequentia tripollicaria, ſuperne verſus ſenſim minora, utrinque pilis ſtellatis molliter villoſa, ſupra viridiora, ſecundum nervos lineata, ſubtus incana ſerrata, ſerraturis infimis foliorum ſuperiorum glanduloſis, baſi integra.* Petioli *breves, e pilis ſtellatis villoſi.* Stipulae *ſubulatae.* Flores *ex alis foliorum ſuperiorum.* Germen *ſetis minutis tectum. A T. velutina differt:* foliis *duplo anguſtioribus extrorſum verſus non anguloſis, acutis nec acuminatis,* pilis *ſubtus magis diſtinctis,* ſtipulis *ſetaceo-ſubulatis.* Vahl. l. c.

3. TRIUMFETTA *Bartramia.*

T. foliis baſi integris indiviſis. *Syſt. nat.* 13. *p.* 327.
Bartramia *Fl. zeyl.* 174. *Sp. pl.* 1. *p.* 389.
Lappula benghalenſis tetraſpermos, ribeſii folio, echinis orbiculatis ad foliorum ortum plurimis ſeſſilibus. *Pluk. alm.* 206. *t.* 41. *f.* 5.
Agrimonia maderaſpatana, folio rotundo ſingulari ſubtus incano. *Pet. gaz.* 51. *t.* 42. *f.* 10.
Lappago amboinica. *Rumph. amb.* 6. *p.* 59. *t.* 25. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 5.
Bartramſche Triumfette. W.
Habitat in India. ☉ (v. ſ.)

*4. TRIUMFETTA *velutina.*

T. floribus completis, foliis ovatis ſubangulatis acuminatis, ſubtus tomentoſo-incanis. *Vahl. ſymb.* 3. *p.* 62.*

Weichblättrige Triumfette. W.
Habitat in inſulae Mauritii.

*5. TRIUMFETTA *procumbens.*
T. foliis ſubrotundo-cordatis ſubtrilobis tomentoſis, caule procumbente. *Forſt. prodr. n.* 204.
Liegende Triumfette. W.
Habitat in Societatis *inſulis.*

*6. TRIUMFETTA *hirta.*
T. floribus completis, foliis trilobis, paniculae terminalis ramis dichotomis hirtis. *Vahl. ſymb.* 3. *p.* 63.*
Rauhe Triumfette.. W.
Habitat in St. Martham Americae. ♄
Folia *ſupra glabra ſeptemnervia, oculo armato pilis ſtellatis tecta.* Vahl. l. c.

7. TRIUMFETTA *ſemitriloba.*
T. foliis ſemitrilobis, floribus completis. *Mant.* 73.
T. fructibus calyculatis; foliis ſemitrilobis. *Iacq. amer.* 147.
T. villoſa, foliis inferioribus angulato-ovatis ſerrato-dentatis, floribus ternatis, faſciculis geminatis folio ſuboppoſitis. *Brown. jam.* 233.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 6.
Dreylappige Triumfette. W.
Habitat in America *calidiore.* ♄ (v. ſ.)
Simillima *T. Lappulae, ſed* flores *calyce inſtructi.*

*8. TRIUMFETTA *grandiflora.*
T. foliis ſubcordato-ovatis integris ſerratis piloſiuſculis, floralibus lanceolatis, ramis hirtis, floribus completis. *Vahl. eclog.* 2. *p.* 34.*
Groſsblütige Triumfette. W.
Habitat in Montſerrat. ♄
A reliquis mihi notis ſpeciebus differt, corollis *duplo vel triplo majoribus.* Vahl. l. c.

*9. TRIUMFETTA *macrophylla.*
T. foliis ovato-cordatis integris inaequaliter ſerratis acuminatis tomentoſis baſi glanduloſis, floribus completis. *Vahl. eclog.* 2. *p.* 34.*
Groſsblättrige Triumfette W.
Habitat in America *meridionali.* ♄
Simils T. Lappulae, cum vero flores calyculati, nec ecalyculati ut in illa, diverſum eſſe credo. A T. rhombeaefolia tomentoſitate *et* figura *foliorum dignoſcitur.* Vahl. l. c.

*10. TRIUMFETTA *rhombeaefolia.*

T. foliis rhomboideis, ſummis lanceolato-ovatis, floribus completis. *Swartz. prod.* 76. *Fl. ind. occid.* 2. *p.* 865. *

T. (*rhomboidea*) floribus calyculatis, foliis rhombeo-ovatis. *Jacq. amer.* 147. *t.* 90. *

Rauthenförmige Triumfette. W.

Habitat in India *occidentali.* ♄

11. TRIUMFETTA *annua.*

T. foliis ovatis indiviſis, rarius lobatis. *Mant.* 73.

T. foliis oblongo-ovatis obtuſis ſerratis, petiolis longiſſimis. *Mill. ic.* 199. *t.* 29.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 7.

Jährige Triumfette. W.

Habitat in India *orientali.* ☉

*945. EURYA.

Gen. plant. edit. Schreb. n. 820.

Cal. duplex, exterior 2-phyllus, interior 5-phyllus. *Petala* 5. *Capſ.* 5-locul. polyſperma.

*1. EURYA *japonica.*

E. japonica. *Thunb. jap.* 191. *t.* 25. *

Fiſakaki. *Koempf. am. exot.* 5. *p.* 778.

Japaniſcher Weitaſt. W.

Habitat in Iaponiae *montibus.* ♄

Folia *alterna petiolata oblonga acuminata emarginata ſempervirentia glabra venoſa reticulata dentata, baſi integerrima attenuata.* Flores *parvi bini pedunculati cernui axillares petiolo breviores.* W.

946. PEGANUM.

Gen. plant. edit. Schreb. n. 821.

Cor. 5-petala. *Cal.* 5-phyllus, ſ. o. *Capſ.* 5-locularis, 3-valvis, polyſperma.

1. PEGANUM *Harmala.*

P. foliis multifidis, caule herbaceo. W.

P. foliis multifidis. *Hort. upſ.* 144. *Gron. orient.* 165. *Gmel. ſib.* 4. *p.* 177. *n.* 96. *Fabric. helm. p.* 228. *Willich. obſ. n.* 50. *Mill. dict. n.* 1. *Giſecke ic. faſc.* 1. *n.* 11. *Blackw. t.* 310.

Peganum. *Hort. cliff.* 206. *Hort. upſ.* 144. *Roy. lugdb.* 461.
Ruta ſylveſtris, flore magno albo. *Bauh. pin.* 336.
Harmala. *Dod. pempt.* 121.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 613.
Gemeine Harmala. W.
Habitat in arena Madriti, Alexandriae, Cappadociae, Galatiae, Sibiriae. ♃ (v. v.)
Petala 5. Stam. 15. *Fabric. l. c.* R.
Calycis *foliola baſi utrinque dente ſubulato inſtructa.* W.

*2. PEGANUM *crithmifolium.*

P. foliis multifidis, caule fruteſcente. *Retz. obſ. p.* 34.
Bacillenblättrige Harmala. W.
Habitat ad mare Caſpium ♄
Similis P. Harmalae, ſed: caulis 3-4-*pedalis perennans ſempervirens*, Flores *minores*, Calyx *ſimplex flore brevior. In P. Harmala eſt* caulis *annuus herbaceus vix ſesquipedalis*, flores *majores*, calycis *foliola diviſa corolla longiora.* Retz. l. c.

*3. PEGANUM *retuſum.*

P. foliis cuneiformibus, caule fruticoſo. *Vahl.* 3. *p.* 63*
P. (*retuſum*) foliis ſubcarnoſis ſeſſilibus cuneiformibus retuſis. *Forſk. deſcr.* 211.
Keilblättrige Harmala. W.
Habitat in Aegypto. ♄

4. PEGANUM *dauricum.*

P. foliis oblongis acutis, caule herbaceo. W.
P. foliis indiviſis. *Sp. pl.* 638.
Ruta foliis ſimplicibus alternis. *Gmel. ſib.* 4. *p.* 176. *t.* 68. *f.* 1.
Harmala montana daurica perennis multicaulis, polygalae folio, flore albo. *Amm. ruth.* 91.
β. Harmala montana, polygalae foliis, floribus luteis. *Amm. ruth.* 92.
Houttuyn Lin. Pfl. Syſt. 13. *p.* 614.
Flachsblättrige Harmala. W.
Habitat in Sibiria. ♃

947. HUDSONIA.

Gen. pl. ed. chreb. n. 822.

Cor. 5-petala. *Cal.* 3-phyllus, tubuloſus. *Stam.* 15. *Capſ.* 1-locularis, 3-valvis, 3-ſperma.

1. HUDSONIA *ericoides.*

H. ericoides. *Mant.* 74. *Berg. act. holm.* 1778. *p.* 18. *t.* 2. *

Ericaeformis ſuffrutex virginianus, floribus exiguis, vaſculo ſeminali oblongo trifariam diviſo. *Pluk. mant.* 88.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 615.

Heidekrautartige Hudſonie. W.

Habitat in Virginia. ♄

Frutex *facie et habitu Ericae.* Ramuli *filiformes, ſparſi, imbricati.* Foliis *ſubulato-aceroſis, hirſutis; tenellis hirſutioribus, incanis, unde quaſi adſperſi gemmis foliaceis, quae ultimi ramuli.* Pedunculi (*e gemmis foliorum*) *ſolitarii, filiformes, foliis longiores.* Calyx *erectus, pubeſcens.*

948. NITRARIA.

Gen. pl. ed. Schreb. n. 823.

Cor. 5-petala, petalis apice fornicatis. *Cal.* 5-fidus. *Stam.* 15. *Drupa* 1-ſperma.

1. NITRARIA *Schoberi.*

N. foliis integerrimis obtuſis. W.

Nitraria. *Act. nov. petrop.* 7. *p.* 315. *t.* 10.. *Murray. prodr.* 191. *f.* 2. *Pall. it.* 1. *p.* 384. *Lerche in Nov. Act. A. N. C. tom.* 5. *app, p.* 162. *Pall. roſſ.* 1. *p.* 80. *t.* 50. *f. A.* *

Oſyris foliis obtuſis. *Hort. upſ.* 295. *Gmel. ſibir.* 2. *p,* 237. *t.* 98.

Caſia fructu nigro. *Amm. ruth.* 256.

β. foliis latioribus, baccis rubris, nuce conica et caule inermi. W.

O. Schoberi. *Pall. roſſ.* 1. *p.* 79. *t.* 50. *f. B.* *

Houttuyn Lin. Pfl. Syſt. 2. *p.* 616.

Ganzblättriger Salpeterſtrauch. W.

Habitat in campis ſalſis Sibiriae *inter* Irtim *et* Ob *flu-*

vios, ad lacus ſalſos circa Jeniſeam *fluvium; β. vero ad mare* Caſpium. ♄

Stamina 15. *Murray l. c.*

Varietas β. eſt forte ſpecies diſtincta; hinc ulterius examinanda. W.

*2. NITRARIA *tridentata.*

N. foliis apice tridentatis. W.

N. (*tridentata*) ramis ſpinoſis, foliis carnoſis. *Desfont. in Uſteri Annal.* 16. *p.* 101.

Nitrariae Schoberi varietas. *Pall. roſſ. p.* 80. *t.* 50. *f. C.?*

Dreyzähniger Salpeterſtrauch. W.

Habitat in maritimis regni Tunetani. ♄

949. PORTULACA.

Gen. pl. ed. Schreb. n. 824.

Cor. 5-petala. *Cal.* 2-fidus. *Capſ.* 1-locularis, circumſciſſa.

1. PORTULACA *oleracea.*

P. foliis cuneiformibus, floribus ſeſſilibus. *Roy. lugdb.* 473. *Gron. virg.* 59. *Hort. upſ.* 147. *Mat. med. p.* 119. *Dalib. par.* 168. *Sauv. monſp.* 168. *Brown. jam.* 233. *Scop. carn. n.* 564. *Pollich. pal. n.* 449. *Blackw t.* 287. *Hoffm. germ.* 161. *Roth. germ. I.* 202. *II.* 518.

P. foliis dilatatis laevibus, ſeſſilibus, confertis. *Hall. helv. n.* 968.

P. foliis cuneiformibus verticillatis ſeſſilibus, floribus ſeſſilibus. *Hort. cliff.* 207.

P. anguſtifolia ſylveſtris. *Bauh. pin.* 288.

P. ſylveſtris. *Matth. epit.* 258. R.

P. ſylveſtris minor. ſ. ſpontanea. *Bauh. hiſt.* 3. *p.* 678. R.

β. Portulaca latifolia ſativa. *Bauh. pin.* 288.

P. domeſtica. *Lob. ic.* 388.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 8.

Gemeiner Portulac. W.

Habitat in Europa, India, *Inſ.* Aſcenſionis, America. ☉ (v. v.)

Capſula *circumſciſſa.* Horologi florae vigilias 11-12 habet.

Piſtilla *quinque habere. Haller. Scop.* Stamina 8 *ad*

15. *Haller.* Stamina 6 *ad* 12 *et plura.* *Stigma* 4 *- partitum* *Gmel. it.* 3. *p.* 395, R.

2. PORTULACA *piloſa.*

P. foliis ſubulatis alternis: axillis piloſis, floribus ſeſſilibus terminalibus. *Mill. dict. n.* 2. *Kniph. cent.* 4. *n.* 64.

P. foliis lienearibus acutis, caule procumbente piloſo. *Roy. lugdb.* 473.

P. curaſſavica, anguſto longo lucidoque folio, procumbens. *Comm. hort* 1. *p.* 9. *t.* 5.

P. curaſſavica lanuginoſa procumbens. *Herm. par.* 215. *t.* 215.

P. lanuginoſa procumbens, vermiculatae foliis americana. *Pluk. alm.* 304. *t.* 247. *f.* 7. *Volk. norib.* 341. *t.* 341.

Anacampſeros ſupina minor, foliis linearibus turgidis, floribus ſummis ramulis confertis. *Brown jam.* 234.

β. P. curaſſavica lanuginoſa, pſylii folio, erectior et elatior, flore dilutius rubente. *Herm. par.* 214. *t* 214.

P. curaſſavica lanuginoſa, kali folio, erectior et elatior. *Pluk. alm.* 303. *t.* 105. *f.* 4.

Houttuyn Lin. Pfl. Syſt. 7. *p* 9.

Haariger Portulac. W.

Habitat in America *meridionali.* ⊙ (v.v.)

Folia *alterna, ſed terminalia ſ. floralia octo in orbem digeſta.* Flores *ſeſſiles, terminales, calyce bifido;* Staminibus *quindecim;* Stylo *quinquefido,* Capſula *circumſciſſa.*

3. PORTULACA *quadrifida.*

P. bracteis quaternis, floribus quadrifidis, caule geniculis piloſis. *Mant.* 73. *Iacq. Collect.* 2. *p.* 356. *t.* 17. *f.* 4. *

P. linifolia. *Forſk aegypt. p* 92. R.

P. lanuginoſa procumbens, flore luteo. *Herm. par.* 216.

Sedum minus arabicum, flore luteo, foliis planis mucronatis. *Pluk. mant.* 169.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 10.

Viertheiliger Portulac. W.

Habitat in Aegypto. ⊙

Radix *fibroſa.* Caules *proſtrati, numeroſi, oleracei, teretes, laeves, alterni, ramoſi, ſemipedales, geniculis radicantibus piloſis.* Folia *oppoſita, patentia, diſtantia, ovato - lanceolata, carnoſa, laevia, ſeſſilia, integerrima, ſubtus concava, adſperſa mi[illegible]apha.*

nis. Petioli *florentes terminales, adulti laterales facti, uniflori, ſuperne incraſſati, longitudine foliorum, terminati.* Bracteis *quaternis ſ. cruciatis, patentibus folia ſimilibus.* Flos *ſeſſilis, flavus, cinctus pilis albis, ut genicula.* Perianthium *bivalve, membranaceum, ovatum.* Corolla 4-*partita, ovato-oblonga, marceſcens.* Filamenta 8, *ſetacea, baſi extrorſum curvata, longitudine corollae.* Antherae *ſubrotundae.* Piſtillum: *Germen inferum, rotundatum:* Stylus *cylindricus, ſuperne craſſior, longitudine ſtaminum.* Stigmata 4. *oblonga obtuſa, recurvata, pubeſcentia.* Capſula *ovata, leviter tetragona, acutiuſcula, unilocularis, circumſciſſa* Semina *plura, ſubrotunda, muricata.* H. V.

4. PORTULACA *halimoides.*

P. foliis oblongis carnoſis, caule corymboſo, floribus ſeſſilibus. *Sp. pl* 639.

P. erecta, ſedi minoris facie, capitulo tomentoſo. *Sloan. jam.* 88. *hiſt.* 1. *p.* 205. *t.* 129. *f.* 3.

Halimus minimus, foliis oblongis ſucculentis tumentibus, ſummis ramulis denſiſſime ſitis. *Brown. jam.* 206.

Houttuyn Lin. Pfl Syſt. 7. *p.* 10.

Meldenartiger Portulac. W.

Habitat in Iamaica. ⊙

Calyx *diphyllus.* Capſula *circumſciſſa.*

5. PORTULACA *meridiana.*

P. foliis ellipticis carnoſis planis, geniculis piloſis, floribus ſeſſilibus terminalibus. *Lin. ſuppl.* 248.

Mittagblütiger Portulac. W.

Habitat in India *orientali.* ⊙ (v. v.) H U.

Caules *ſtatura Sedi acris filiformes herbacei repentes articulati teretes laxi rubentes.* Genicula *pilis aliquot adſperſa radicantia.* Folia *elliptico-lanceolata laevia ſubcarnoſa terminalia oppoſita etiam quaterna.* Flores *lutei terminales ſolitarii ſeſſiles, cincti lana foliisque quatuor.* Calyx *bivalvis tenerrimus, foliolis ovatis acutis rubicundis.* Stamina 4-8 *numero inconſtantia. Floret hora duodecima per diem.*

Genus novum ex hac conſtituit b. m. Parens et Meridianam *vocavit; at differentia non tanta mihi viſa, quin cum Portulaca combinari poſſet, conſentiente affinitate. Polymorphum valde eſt Portulacae genus quoad fructificationem et numerum partium ſi itaque una ſepararetur ſpecies, ſeparandae etiam plures.*

*950 TALINUM.

Juſſ. gen. ed. Uſt. p. 346.

Cor. 5-petala. *Cal.* 2-phyllus. *Capſ.* 3-valvis polyſperma. *Semina* arillata.

1. TALINUM *triangulare.*

T. foliis planis canaliculatis cuneiformibus emarginatis mucronatis, racemo ſimplici pedunculo triquetro. W.

Rülingia (*triangularis*) foliis obovatis planiuſculis, racemo ſimplici, pedunculo triquetro. *Ehrh. Beitr.* 3. *p.* 134.

Portulaca (*racemoſa*) foliis obovatis planiuſculis, racemo ſimplici triquetro. *Sp. pl.* 2. *p.* 640.

Portulaca (*triangularis*) pedunculis multifloris, triangularibus, *Iacq. amer.* 147. *obſ.* 1. *p.* 35. *t.* 23.

Chamaeciſtus americana, portulacae folio, flore albo. *Herm. parad.* 21.

Helianthemum fruteſcens, portulacae folio. *Plum. ſpec.* 7. *t.* 150. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 11.

Dreyeckiges Talinum. W.

Habitat in America *ad maris littora.* ♄

Calyx *diphyllus.* Capſ. *trivalvis.*

*2. TALINUM *craſſifolium.*

T. foliis planis obovatis mucronatis, corymbo elongato, pedunculis triquetris. W.

Portulaca (*craſſifolia*) foliis lanceolatis planis, racemis triquetris caule erecto. *Murr. ſyſt. veg.* 14. *p.* 466 *Iacq. hort.* 3. *t.* 52.

Dickblättriges Talinum. W

Habitat - - - - ♄ (v. v.)

Simile praecedenti, differt foliis *longioribus latioribus non emarginatis,* floribus *in corymbo paucifloro (nec in racemo ſimplici) collectis,* corollis *rubris nec flavis,* caule *erecto nec decumbente. Ceterum* calyx *eſt diphyllus et* capſula *trivalvis.* W.

3. TALINUM *Anacampſeros.*

T. foliis ovatis ſubtus convexo-gibbis, racemo ſimplici, pedunculo tereti. W.

Rülingia (*Anacampſeros*) foliis ovatis gibbis, racemo ſimplici, pedunculo tereti. *Ehrh. Beit.* 3. *p.* 133.

Portulaca (*Anacampſeros*) foliis ovatis gibbis pedunculo multifloro, caule fruticoſo. *Mill. dict. n.* 3.

Anacampſeros foliis ovatis convexis acutiuſculis. *Hort. upſ.* 146.

Anacampſeros foliis acuminatis. *Hort. cliff.* 207. *Roy. lugdb.* 412.

Telephiaſtrum folio globoſo. *Dill. elth.* 375. *t.* 281. *f.* 363.

Portulaca africana ſempervirens, flore rubicundo. *Comm. hort.* 2. *p.* 177. *t.* 89. *Raj. ſuppl.* 507.

Oxalis affinis planta aphylla mammillaris. *Burm. afr.* 76. *t.* 30. *f.* 2. *vitiata.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 12.

Nabelkrautartiges Talinum. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Axillae foliorum piloſae ſunt, et Folia ſucculenta conjungunt hanc cum Portulacis, quamvis ſtamina plurima et Capſula trivalvis. Calyx diphyllus eſt.

4. TALINUM *patens.*

T. foliis ovatis planis, panicula terminali, pedunculis dichotomis. W.

Rülingia (*patens*) foliis lanceolato-ovatis planis, panicula ramoſa. *Ehrh. Beitr.* 3. *p.* 135.

Portulaca (*patens*) foliis lanceolato-ovatis planis, panicula ramoſa, calycibus diphyllis. *Iacq. hort. t.* 151.

P. (*paniculata*) floribus paniculatis. *Iacq. amer.* 148.

β. Talinum (*reflexum*) foliis lanceolato-ovatis ſeſſilibus oppoſitis, florum panicula ramoſa. *Cavan ic.* 1. *p.* 1. *t.* 1.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 12.

Riſpenblütiges Talinum. W.

Habitat in America. ♄ (v. v.)

Caulis *pedalis, flexilis:* Ramis *alternis, laevibus, tetragonis; inferioribus longioribus.* Folia *alterna, ſubpetiolata, lanceolato-ovata, ſubſucculenta, ſemipollicaria, integerrima, carinata fragilia.* Panicula *terminalis, elongata; pedicellis filiformibus, dichotomis.* Calyx *diphyllus, ruber.* Petala 5, *ovata, rubra.* Antherae *luteae.* Styli 3. Capſula *globoſa, 1-locularis, 3-valvis: valvulis duplicatis.* Semina *plura, nigra.*

Diſtincta *omnino a T.* fruticoſo, *cui flores albi et calyx 5-phyllus.*

In varietate α. caulis *eſt perennis, et* petala *rubra, in β. vero* caulis *eſt annuus et* petala *flava an varietates?* W.

*5 TALINUM *cuneifolium.*

T. foliis cuneiformibus, panicula terminalis, pedunculis trifloris. W.

Portulaca (*cuneifolia*) foliis cuneiformibus planis, pedunculis inferioribus racemi trifloris, calycibus diphyllis. *Vahl. ſymb.* 1. *p.* 33.

Orygia (*portulacifolia*) foliis alternis cuneatis glabris. *Forſk. deſcr.* 103 *

Keilblättriges Talinum. W.

Habitat in Arabia *felici.* ♄

Affinis T. patenti, dignoſcitur ab illo, pedunculis *craſſioribus, inferioribus apice* floribus *ternis pedicellatis.* Vahl. l. c.

Calyx *diphyllus.* Corolla *violaceo-rubra.* Capſula *trivalvis.* W.

*6. TALINUM *decumbens.*

T. foliis obovatis planis mucronatis, racemis axillaribus, calycibus pentaphyllis, caule decumbente. W.

Portulaca (*decumbens*) foliis obovatis mucronatis, calycibus pentaphyllis, caule fruticoſo decumbente. *Vahl. ſymb.* 1 *p.* 33. *

Orygia (*decumbens*) caule decumbente fruticoſo, foliis ovatis acutis ſubfarinoſis. *Forſk. deſcr.* 103. *

Liegendes Talinum. W.

Habitat in Arabia *felici.* ♄

Calyx *pentaphyllus.* Corolla *polypetala rubro-violacea.* Capſula *quinquevalvis.* W.

7. TALINUM *fruticoſum.*

T. foliis obovatis planis emarginatis, corymbo terminali, calycibus pentaphyllis, caule erecto. W.

Portulaca (*fruticoſa*) foliis obovatis, planiuſculis, pedunculis racemoſis, calycibus pentaphyllis, caule fruticoſo. *Syſt. Veg. p.* 371.

P. (*paniculata*) foliis ovatis planis, panicula racemoſa, *Spec. pl.* 2. *p.* 640.

P. americana latifolia erecta, floribus albis.] *Comm. hort.* 1. *p.* 7. *t.* 4.

Portulacae facie maritima fruticoſa erecta americana. *Sloan. jam.* 87. *hiſt.* 1. *p.* 205.

Anacampſeros foliis radicalibus mollibus ovatis glabris, ſcapo aſſurgenti paniculato. *Brown. jam.* 234.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 13.

Strauchartiges Talinum. W.

Habitat in Americae *maritimis.* ♄

Calyx pentaphyllus. Capſula trivalvis. Flores *albi.*

951. LYTHRUM.

Gen. pl. ed. Schreb. n. 825.

Cal. 12-dentatus. *Petala* 6, calyci inſerta. *Capſ.* 2-locularis, polyſperma.

1. LYTHRUM *Salicaria.*

L. foliis oppoſitis cordato-lanceolatis, floribus ſpicatis dodecandris. *Mill. dict. n.* 1. *Scop. carn. n.* 565. *Pollich. pal. n.* 450. *Blackw. t.* 520. *Flor. dan. t.* 671. *Kniph. cent.* 5. *n.* 55. *Hoffm. germ.* 162. *Roth. germ. I.* 203. *II.* 510.

L. foliis oppoſitis. *Hort. cliff.* 178. *Fl. ſuec.* 393; 422. *Roy. lugdb.* 458. *Guett. ſtamp.* 2. *p.* 122.

L. foliis verticillatis. *Fl. lapp.* 197.

Salicaria foliis lanceolatis, ſubhirſutis, floribus ſpicatis. *Hall. helv. n.* 854.

Lyſimachia ſpicata purpurea. *Bauh. pin.* 246.

Lyſimachia purpurea communis major. *Cluſ. hiſt.* 2. *p.* 51.

β. Blattaria rubra ſpicata major, folio ſubrotundo. *Moriſ. hiſt.* 2. *p.* 490. *ſ.* 5. *t.* 10. *f.* 11.

γ. Lyſimachia trifolia ſpicata purpurea. *Bocc. muſ.* 167.

δ. Lyſimachia quadrifolia purpurea, mollioribus et longioribus foliis. *Bocc. muſ.* 167.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 14.

Gemeiner Weiderich. W.

Habitat in Europa *ad ripas aquarum.* ♃ (v. v.)

Folia *terna, quaterna rarius. Pollich. l. c.*

2. LITHRUM *virgatum*

L. foliis oppoſitis lanceolatis, panicula virgata, floribus dodecandris ternis. *Sp. pl.* 642. *Gmel. ſib.* 4. *p.* 175. *Hoffm. germ.* 162.

L. (*auſtriacum*) foliis oppoſitis lineari-lanceolatis ſubſeſſilibus nitidis. *Iacq. vind.* 243. *auſt. t.* 7.

Salicaria glabra ramoſiſſima, longiſſimis et anguſtiſſimis foliis. *Amm. ruth.* 89.

Salicaria glabra, longioribus et anguſtioribus foliis. *Amm. ruth.* 88.

Lyſimachia rubra non ſiliquoſa. *Bauh. pin.* 246.

Lyſimachia rubra 2. ſ. minor. *Cluſ. hiſt. p.* 51.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 15.
Ruthenförmiger Weiderich. W.
Habitat in Auſtria, Sibiria, Tartaria. ♃ (v. ſ.)

Caulis *paniculatus, ramis alternis, longis.* Flores 2. ſ. 3, *pedicellati, ex ſingula ala alterna racemi.* Variat *rarius foliis alternis.*

*3. LYTHRUM *acuminatum.* W.

L. foliis oppoſitis lanceolato-acuminatis, racemo elongato, pedunculis ſolitariis. W.
Salicaria orientalis ſalicis folio acutiſſimo et glabro. *Tournef. cor.* 18.
Langſpitziger Weiderich. W.
Habitat in Iberia. ♃. (v. ſ.)

Caulis *bipedalis erectus ramo uno alterove inſtructus.* Folia *oppoſita ſeſſilia lanceolata baſi attenuata, apice valde acuminata glabra, margine ſcabra integerrima, parum venoſa; intermedia majora bipollicaria et ultra.* Folia *floralia lineari-lanceolata, ſumma ſubulata calyce breviora.* Flores *pedunculati axillares oppoſiti ſolitarii, in ſummo caule interdum bini.* Calyx *cylindraceo-clavatus ſtriatus* 12 *breviſſime dentatus.* Petala *ſex lanceolata acuta.* Stamina *longitudine corolla.* Caulis *ſumma pars ob folia floralia anguſtiſſima minuta format racemum ſemipedalem elongatum tenuem.* W.

Diverſiſſima a praecedente, licet habitu quemadmodum conveniat: foliis *longioribus magis acuminatis margine ſcabris,* floribus *anguſtiſſimis,* pedunculis *ſolitariis,* calycibus *anguſtioribus,* petaliſque *lanceolatis. W.*

4. LYTHRUM *triflorum.*

L. glaberrimum, foliis oppoſitis ſubſeſſilibus lanceolatis integris, pedunculis axillaribus oppoſitis, capitulo trifloro. *Lin. ſuppl.* 249.
* Dreyblumiger Weiderich. W.
Habitat in America. ♃.

A reliquis facile diſtinguitur: pedunculis *filiformibus terminatis,* bracteis *binis lanceolatis canaliculatis patentibus flore longioribus, inter quas* flores *tres aequales brevius pedicellati coerulei parvi.*

5. LYTHRUM *verticillatum.*

L. foliis oppoſitis ſubtomentoſis ſubpetiolatis, floribus verticillatis lateralibus. *Sp. pl.* 641. *Mill. dict. n.* 6.

L. foliis oppoſitis, floribus verticillatis. *Gron. virg.* 52.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 16.
Wirtelblütiger Weiderich. W.
Habitat in Virginia.
Pedunculi *multiflori*, *breviſſimi.*

6. LYTHRUM *petiolatum.*

L. foliis oppoſitis linearibus petiolatis, floribus dodecandris. *Sp. pl* 641. *Mill. dict. n.* 7.
Lythrum foliis petiolatis. *Gron virg.* 52.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 17.
Geſtielter Weiderich. W.
Habitat in Virginia.

7. LYTHRUM *racemoſum.*

L. diffuſum, foliis oppoſitis petiolatis ovatis, racemis terminalibus, floribus oppoſitis. *Lin. ſuppl.* 250.
Traubenblütiger Weiderich. W.
Habitat in America *meridionali.* ♄. *Mutis.*
Planta *ſuffruticoſa diffuſa.* Folia *ovata integerrima venoſa.* Racemi *ex dichotomia ſ. terminales longi.* Flores *remoti per paria oppoſiti*, *longi*, *pedicellati erecti bracteati.* Bracteae *ovatae.* Apices *ramulorum*, Calyces *et* Pedicelli *piloſo-viſcidi.*

*8. LYTHRUM *ciliatum.*

L. foliis oppoſitis petiolatis ovatis glabris margine ciliatis, racemis terminalibus, floribus ſubſecundis decandris. *Swartz prodr.* 76. *Fl. ind. occid.* 2. *p.* 868. *
Gefranzter Weiderich. W.
Habitat in montibus altis petroſis Jamaicae auſtralis. ♄ (v. ſ.)
Proximum L. racemoſo, *ſed differt:* foliis *ciliatis*, floribus *racemoſis*, *breviter pedunculatis ebracteatis ſubſecundis.* Swartz l. c.

9. LYTHRUM *Pemphis.*

L. fruticoſum hirſutum, foliis oppoſitis oblongis integris, floribus axillaribus pedunculatis ſolitariis, capſula circumſciſſa uniloculari. *Lin. ſuppl.* 249. *Forſt. prodr. n.* 205.
Pemphis acidula. *Forſt. gen. n.* 34. *t.* 34.
Mangium porcellanicum. *Rumph. amb.* 3. *p.* 126. *t.* 84.
Strauchartiger Weiderich. W.
Habitat ad littora maris inſularum Zeylonae *et* Teauteae. ♄

Frutex *canus.* Folia *approximata in ſummitate ramorum.* Calyx 12-*dentatus, denticulis inaequalibus.* Petala 6 *calyci inſerta alba.* Stamina 12. Capſula *turbinata unilocularis circumſciſſa.*

Genus videtur diverſum ob fructum unilocularem circumſciſſum, ſed ſi una valeret differentia ad genus ſeparandum, nimis multiplicarentur genera: Botanicus, quantum permittat differentia generica, genera naturalia ſeparare non debet.

10. LYTHRUM *dipetalum.*

L. hiſpido-viſcoſum, foliis ternis ſ. oppoſitis ſeſſilibus ovatis, floribus axillaribus nutantibus dipetalis. *Lin. ſuppl.* 250.

Zweyblättriger Weiderich. W.

Habitat in America. *Mutis.* ♄

Frutex *hiſpido-viſcoſus fuſcus.* Rami *alterni.* Folia *ſeſſilia ovato-oblonga integerrima, utrinque hiſpiditate ſcaberrima, ſupra ſplendentia; inferiora verticillata, quarterna, terna; ramulorum oppoſita.* Flores *axillares pedicellati nutantes.* Calyx *linearis hiſpidus viſcoſus.* Petala *bina majora, margini ſuperiori calycis inſerta erecta obovata violacea ſ. coerulea. Floribus ſpecioſa et diſtincta planta.*

11. LYTHRUM *lineare.*

L. foliis oppoſitis linearibus; floribus oppoſitis hexandris. *Sp. pl.* 641. *Mill. dict. n.* 8.

L. foliis linearibus, floribus hexandris ſolitariis. *Gron. virg.* 162.

Habitat in Virginia.

Caulis *filiformis erectus angulatus,* Foliis *oppoſitis linearibus, terminatus* Spica *majore et duabus lateralibus minoribus compoſitis ex floribus oppoſitis remotis, ſub ſingulis folium.* Facies *L. Hyſſopifoliae.*

12. LYTHRUM *Parſonſia.*

L. foliis oppoſitis ovalibus, floribus ſeſſilibus alternis hexandris, caule diffuſo. *Sp. pl.* 641. *Swartz obſ.* 193.*

Parſonſia herbacea, foliis ovatis oppoſitis, floribus ſingularibus foliis ad alterum latus interpoſitis. *Brown. jam.* 199. *t.* 21. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 18.

Ausgebreiteter Weiderich. W.

Habitat in Jamaicae *et* Hiſpaniolae *aridis.* ♃

Caules *proſtrati.* Folia *ſubſeſſilia, nuda, integerrima, trinervia.* Racemi *alterni.* Flores *ſolitarii, alterni, ſeſſiles.*

13. LYTHRUM *Melanium.*

L. foliis oppoſitis ovatis, floribus alternis ſubdecandris, caule proſtrato. *Sp. pl.* 641. *Swartz obſ.* 193.*

Melanium herbaceum reclinatum, foliis ovatis oppoſitis, floribus ſingularibus ad alas alternas. *Brown. jam.* 217.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 18.

Liegender Weiderich. W.

Habitat in Jamaica. ♃

*14. LYTHRUM *cordifolium.*

L. foliis oppoſitis ſubſeſſilibus cordatis acutis ſcabris, racemis terminalibus axillaribusque, floribus decandris. *Swartz prodr.* 76. *Fl. ind. occid.* 2. *p.* 866.*

Herzblättriger Weiderich. W.

Habitat in Domingo *in ſylvis interioribus.* ♄

15. LYTHRUM *Hyſſopifolia.*

L. foliis alternis linearibus, floribus hexandris. *Hort. upſ.* 118. *Mill. dict. n.* 3. *Scop. carn. n.* 566. *Iacq. auſtr. t.* 133. *Scholl. barb. n.* 372. *Pollich. pal. n.* 451 *Hoffm. germ.* 162. *Roth. germ. I.* 203. *II.* 520.

L. foliis alternis. *Hort. cliff.* 178. *Sauv. monſp.* 53. *Roy lugdb.* 458. *Gort. gelr.* 271.

Salicaria foliis linearibus, floribus in alis ſeſſilibus. *Hall. helv. n.* 855. R.

Salicaria hyſſopi folio latiore. *Hall. Rupp. jen.* 147. *t.* 6. *f.* 3.

Hyſſopifolia. *Bauh. pin.* 218.

β. Lyſimachia linifolia purpureo-caerulea. *Bauh. pin.* 246.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 19.

Hopblättriger Weiderich. W.

Habitat in Germaniae, Helvetiae, Angliae, Galliae *inundatis.* ⊙ (v. v.)

Caules *proſtrati, rigiduli, ramoſi, virgati.* Folia *alterna, ſeſſilia, lineari-lanceolata, obtuſa, integerrima.* Flores *axillares, ſolitarii.* Calyx *tubuloſus dentibus* 5 *ſ.* 6, *patulis.* Petala 6, *purpurea, baſi alba.* Stamina 6, *calyce breviora.*

16. LYTRHUM *Thymifolia.*

L. foliis alternis linearibus, floribus tetrapetalis. *Sp. pl.* 642. *Lerche in nov. Act. A. N. C. V. app.* 186. *Hoff. germ.* 162.

L. foliis linearibus. *Sauv. monſp.* 49.

Pentagloſſum linifolium. *Forſk. aegypt. p.* 11.

Salicaria minima tenuifolia. *Tournef. inſt.* 254.

Salicaria hyſſopi folio anguſtiore. *Buxb. hal.* 290.*

Hyſſopifolia minor ſ. potius thymifolia maritima. *Bauh. hiſt.* 3. *p.* 792.

Polygonum aquaticum minus. *Barr. ic.* 773. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 19.

Thymianblättriger Weiderich. W.

Habitat in Italiae et G. Narbonenſis *vliginoſis, in* Hircania. *Lerche.* ⊙ (v. v.)

Petala 4, *ſtamina* 2, *calyx* 4-*dentatus.* Folia *lineari-lanceolata, alterna, ima oppoſita.* Flores *ſolitarii, axillares, ſeſſiles.* Calyx *ſtipulis* 2 *lateralibus fulcitus.* Caulis *erectus.*

Forte ſola praecedentis varietas? W.

*952. CUPHEA,

Iacq. hort. 2. *p.* 83.

Cal. 6 dentatus inaequalis. *Petala* 6 inaequalia calyci inſerta. *Capſ.* 1-locularis; conceptaculo triquetro.

1. CUPHEA *viſcoſiſſima.*

C. vicoſciſſima. *Iacq. hort.* 2. *p.* 83. *t.* 177. *Roth. beytr.* 1. *p.* 124. *Ait. Kew.* 2 *p.* 129.

Balſamona Pinto. *Vandell. faſc. p.* 15. *t.* 3.

Lythrum (*Cuphea*) foliis oppoſitis ovato-oblongis ſcabriuſculis, floribus dodecandris. *Lin. ſuppl.* 249.

Klebrige Cuphea. W.

Habitat in Braſiliae *umbroſis humentibus.* ⊙ (v. v.)

Radix *fibroſa annua.* Caulis *herbaceus teres erectus pedalis pubeſcenti-viſcidus purpuraſcens.* Rami *alterni ſimplices.* Folia *oppoſita petiolata ovato-oblonga integerrima laevia.* Flores *purpurei laterales, breviſſime pedunculati ſolitarii decumbentes.* Calyx *monophyllus cylindricus* 12 *ſtriatus,* 6 *dentatus, dente ſuperiore latiore, baſi ſuperiore producta, pubeſcenti-viſcidus.* Coralla 6 *petala, petulis calycis fauci inſer-*

tis, ſuperioris lateris duobus majuſculis. Nectarium *ſquamula reflexa intra gibbum calycis.* Stamina 12, *in fauce calycis.* Capſula *oblongiuſcula unilocularis.* Semina *plura lentiformia tandum nigra.* Capſula *una cum calyce ante maturitatem longitudinaliter rumpitur et exit adſcenditque* receptaculum *ſeminum cum immaturis ſeminibus a latere interiori, ut nuda matureſcant, more Phrymae.*

953. GINORIA

Gen. plant. edit. Schreb. n. 826.

Cal. 6-fidus. *Petala* 6. *Capſ.* 1-locularis, 4-valvis, colorata, polyſperma.

1. GINORIA *americana.*

G. americana. *Sp. pl.* 642. *Iacq. amer.* 148. *t.* 91.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 618.
Americaniſche Ginorie. W.
Habitat in Cuba *ad ripas fluviorum rupeſtres et glareoſas.* ♄
Frutex *pulcher habitu Myrti.*

954. KLEINHOFIA.

Gen. pl. ed. Schreb. n. 827.

Cal. 5-phyllus. *Petala* 5. *Nectar.* campanulatum 5-dentatum ſtaminiferum columnae germinis adnatum. *Germen* pedicellatum. *Capſula* 5-angularis 5-locularis inflata, loculis monoſpermis.

1. KLEINHOFIA *Hoſpita.*

K. Hoſpita. *Sp. pl.* 1365. *Cavan. diſſ.* 5. *p.* 288. *t.* 146.*
Cati-marus. *Rumph. amb* 3. *p.* 177. *t.* 113.
Houttuyn Lin. Pfl. Syſt. p. 4. 251.
Herzblättrige Kleinhofie. W.
Habitat in Java, Amboina *inque inſulis* Philippinis. ♄
Kleinhof. horti Bot. cultor in Java.

Arbor foliis *alternis petiolatis cordatis acuminatis vix ſerratis nudis.* Flores *quodammodo affines Ayeniae.* Fructus *Cardioſpermi: e floris receptaculo* columna *declinata, apice adſcendens ultra apicem ſupremi*

petali, ferens: Nectarium *campanulatum minimum intra quod* Stamina *et* Piſtillum.

955. STERCULIA.

Gen. pl. ed. Schreb. n. 828.

Cal. 5-partitus. *Cor.* 0. *Nectar* campanulatum 5-dentatum ſtaminiferum columnae germinis adnatum. *Germen* pedicellatum. *Capſulae* 5, uniloculares interiori latere dehiſcentes polyſpermae.

*1. STERCULIA *lanceolata.*

S. foliis lanceolatis, capſulis oblongis. W. †
S. (*lanceolata*) foliis lanceolatis, capſulis kermiſinis pauciſpermis. *Cavan. diſſ.* 5. *p.* 287. *t.* 144. *f.* 1. *
Lanzettenblättriger Stinckbaum. W.
Habitat in China. ♄

2. STERCULIA *Balanghas.*

S. foliis ovato-lanceolatis, capſulis obovatis. W.
S. foliis ovatis integerrimis alternis petiolatis, floribus paniculatis. *Sp. pl.* 1430. *Fl. zeyl.* 350. *Cavan. diſſ.* 5. *p.* 286. *t.* 143. *
Nux malabarica ſulcata mucilaginoſa fabacea. *Pluk. alm.* 266.
Clompanus minor. *Rumph. amb.* 3. *p.* 169. *t.* 107.
Cavalam. *Rheed. mal.* 1. *p.* 89. *t.* 49. *Raj. hiſt.* 1754.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 381.
Eyförmiger Stinckbaum. W.
Habitat in India. ♄ (v. ſ.)

Fructus *nutans quinquelobus: lobis reflexis, more Paeoniae dehiſcentibus.*

*3. STERCULIA *crinita.*

S. foliis ovatis trilobisve, capſulis baſi crinitis. W.
S. (*crinita*) foliis ovatis integerrimis alternatim ſparſis, fructus baſi crinita. *Cavan. diſſ.* 5. *p.* 285. *t.* 142. *
S. (*Ivira*) hermaphrodita, foliis ovatis trilobisve, floribus paniculatis, capſulis quinque. *Swartz prodr.* 98.
Ivira pruriens. *Aubl. guj.* 2. *p.* 694. *t.* 279. *
Borſtiger Stinckbaum W.
Habitat in Gujanae *ſylvis, inque* India *occidentali.* ♄

* 4. STERCULIA *cordifolia.*
S. foliis cordatis obſolete trilobis, capſulis acuminatis tomentoſis. W.
S. foliis cordatis latis acuminatis ſeptemnerviis valde venoſis, pericarpio extus tomentoſo, intus ſetoſo. *Cavan. diſſ.* 5. *p.* 286. *t.* 144. *f.* 2. *
Herzblättriger Stinckbaum. W.
Habitat in Senegal. ♄

*5. STERCULIA *colorata.*
S. foliis quinquelobis, lobis acuminatis, calycibus cylindraceo-clavatis, capſulis oblongis glabris coloratis. W.
Sterculia colorata. *Roxb. corom.* 1. *p.* 26. *t.* 25. *
Scharlachrother Stinckbaum. W.
Habitat in Indiae *orientalis montibus.* ♄
Calyces *cylindraceo-clavati apice dentati coccinei.* Capſulae *magnae rubrae.* W.

*6. STERULIA *urens.*
S. foliis quinquelobis, lobis acuminatis, calycibus campanulatis, capſulis ovatis hiſpidis. W.
Sterculia urens. *Roxb. corom.* 1. *p.* 25. *t.* 24. *
Brennender Stinckbaum. W.
Habitat in Indiae *orientalis montibus.* ♄
Folia *praecedentis Aceris ſaccharini facie, ſed multo majora pubeſcentia.* Calyx *viridis campanulatus apice dentatus.* Bracteae *lineares ad baſin pedunculorum, quod non in reliquis ſpeciebus.* Capſulae *virides hiſpidae.* W.

7. STERCULIA *platanifolia.*
S. foliis palmato-quinquelobis calycibus rotatis reflexis. W.
S. hermaphrodita, foliis cordatis lobatis, floribus paniculatis. *Lin. ſuppl.* 423, *Cavan. diſſ.* 5. *p.* 288. *t.* 145. *et diſſ.* 6. *p.* 352. * *Vahl. ſymb.* 1. *p.* 80. *
Firmiana. *Marſili in Act. acad. patav.* 1. *p.* 106. *t.* 1. 2.
Culhamia. *Forſk. deſcr. p.* 96.
Hibiſcus (*ſimplex*) foliis cordatis trilobis repandis integerrimis. *Sp. pl.* 977.
Outom-chu. *Le Comte mem. de la Chine* 1. *p.* 241. *c. fig.*
Ou tong chu. *Du Halde chin.* 2. *p.* 149. *c. fig.*
Houttuyn Lin. Pfl. Syſt. 4. *p.* 152.
Platanblättriger Stinckbaum. W.

Habitat in Japonia, China. ♄ (v. v. ſ. fl.)

Arbor *altiſſima.* Folia *magna alterna in ſummitatibus ramorum conferta longe petiolata cordata integerrima trilobata ſ. quinquelobata: ſinubus rotundatis obtuſiuſculis, glaberrima, quiquenervia.* Panicula *magna lignea ſubpulverulenta.* Flos *magnitudine S. foetidae (eſt multo minor W.)* Calycis *laciniae extus canae, intus glabrae luteſcentes.* Columna *ſtaminum calyce brevior terminata antheris plurimis ovatis, quae tegunt baſin, quamvis columnae inſideant.* Germen *angulatum canum.* Stylus *filiformis canus.* Stigma *emarginatum.*

8. STERCULIA *foetida.*

S. foliis digitatis. *Sp. pl.* 1431. *Fl. zeyl.* 349. *Cavan. diſſ.* 5. *p.* 284. *t.* 141.* *Sonnerat itin. ind. ed. germ.* 2. *p.* 183. *t.* 132. *

Nux zeylanica folio multifido digitato. *Pluk. alm.* 266. *t.* 208. *f.* 3.

Clompanus major. *Rumph. amb.* 3. *p.* 168. *t.* 107.
Karil. *Rheed. mal.* 4. *p.* 75. *t.* 36. *Raj. hiſt.* 1564.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 382.
Gefingerter Stinckbaum. W.
Habitat in India. ♄ (v. ſ. ſ. fl.)

DIGYNIA.

956. HELIOCARPUS.

Gen. pl. ed. Schreb. n. 829.

Cal. 4-phyllus. *Petala* 4. *Styli* ſimplices. *Capſ.* 2-locularis, compreſſa, utrinque longitudinaliter radiata.

1. HELIOCARPUS *americana.*

Heliocarpus. *Hort. cliff.* 211. *t.* 16.* *Trew. ehret. t.* 45. *Mill. dict.*

Houttuyn Lin. Pfl. Syſt. 3. *p.* 620.
Amerikaniſche Sonnenfrucht.
Habitat in Vera Cruce. ♄ (v. ſ. ſ. fl.)

957. AGRIMONIA.

Gen. plant. edit. Schreb. n. 830.

Cal. 5-dentatus altero obvallatus. *Petala* 5. *Sem.* 2, in fundo calycis.

1. AGRIMONIA *Eupatoria.*

A. fructibus hiſpidis, foliis caulinis pinnatis, foliolis oblongo-ovatis, ſpicis elevatis, petalis calyce duplo longioribus. *Ait. Kew.* 2. *p.* 129.

A. foliis caulinis pinnatis: impari petiolato, fructibus hiſpidis. *Mat. med.* 120. *Hort. cliff.* 179. *Fl. ſuec.* 394, 423. *Hort. upſ.* 118. *Roy. lugdb.* 240. *Guett. ſtamp.* i. *p.* 293. *Dalib. pariſ.* 139. *Oed. dan.* 588. *Blackw. t.* 21. *Gmel. ſib.* 3. *p.* 145. *Mill. dict. n.* 1. *Scop. carn. n.* 567. *Pollich. pal. n.* 452. *Ludw. ect. t.* 29. *Kniph. cent.* 5. *n.* 7. *Hoffm. germ.* 163. *Roth. germ. I.* 203. *II.* 321.

A. foliis pinnatis, pinnis alterne minimis. *Hall. helv. n.* 991.

Eupatorium veterum ſ. Agrimonia. *Bauh. pin.* 351.

Eupatorium. *Fuchſ. hiſt.* 244. *Camer. epit.* 756. *Mill. ic.* 15.

Agrimonia (*minor*) foliis caulinis pinnatis, foliis obtuſis dentatis. *Mill. dict. n.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 21.

Gemeiner Odermennig. W.

Habitat in Europae *pratis, viis.* ♃ (v. v.)

*2. AGRIMONIA *odorata.*

A. fructibus hiſpidis, foliis pinnatis, foliolis oblongis, inferioribus diminutis, petalis calyce duplo longioribus. *Ait. Kew.* 2. *p.* 130. *Mill. dict. n.* 3.

Agrimonia odorata. *Camer. hort* 7.

Eupatorium odoratum. *Lob. adverſ.* 308.

Wohlriechender Odermennig. W.

Habitat in Italia. ♃ (v. v.)

3. AGRIMONIA *repens.*

A. fructibus hiſpidis, foliis caulinis pinnatis, foliolis oblongis, ſpicis ſubſeſſilibus, petalis calyce triplo longioribus. *Ait. Kew.* 2. *p.* 130.

A. foliis caulinis pinnatis: impari ſeſſili, fructibus hiſpidis. *Sp. pl.* 643. *Mill. dict. n.* 4.

A. orientalis humilis, radice craſſiſſima repente, fructu in ſpicam brevem et denſam congeſto. *Tournef. cor.* 21.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 22.
Kriechender Odermennig. W.
Habitat in Armenia. ♃ (v. ſ.)
Affinis nimium A. Eupatoriae, ſed Caules *breviores, craſſiores, hirſuti.* Stipulae *longitudine interſtitiorum, reflexae, fere totum obumbrant caulem.* Foliorum *impar non pedicellatum.* Spica *brevis, denſa, ſeſſilis,* Bracteis *longitudine florum.*

*4. AGRIMONIA *parviflora.*
A. fructibus hiſpidis, foliis caulinis pinnatis, foliolis plurimis lanceolatis, petalis calyce ſeſquilongioribus. *Ait. Kew.* 2. *p.* 130.
Kleinblütiger Odermennig. W.
Habitat in America boreali. ☉ (v. v.)

5. AGRIMONIA *Agrimonoides.*
A. foliis caulinis ternatis, fructibus glabris. *Hort. cliff.* 179. *Roy. lugdb.* 179. *Roy. lugdb.* 241. *Mill. dict. n.* 5. *Scop. carn.* 2. *n.* 568. *Hoffm. germ.* 163.
Agrimoniae ſimilis. *Bauh. pin.* 321.
Agrimonoides. *Col. ecphr.* 1. *p.* 145. *t.* 144. *Boerh. lugdb.* 1. *p.* 179. *Barr. ic.* 612.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 23.
Dreyblättriger Odermennig. W.
Habitat in Italiae, Carnioliae *nemoribus umbroſis humentibus.* ♃ (v. ſ.)
Stamina ſaepe vix ultra 8.

TRIGYNIA.

958. RESEDA.

Gen. plant. edit. Schreb. n. 831.

Cal. 1-phyllus, partitus. *Petala* lacinata. *Capſ.* ore dehiſcens, 1-locularis.

1. RESEDA *Luteola.*
R. foliis lanceolatis integris baſi utrinque unidentatis, calycibus quadrifidis. *Mill. dict. n.* 8. *Necker. gal.*

lob. p. 210. *Pollich. pal. n.* 453. *Willich. obſ. n.* 40. *Blackw. t.* 283. *Kniph. cent.* 5. *n.* 74. *Hoffm. germ.* 163. *Roth. germ. I.* 203. *II.* 522.

R. foliis ellipticis, obtuſe lanceolatis, undulatis, calycibus quadrifidis. *Hall. helv. n.* 1058.

R. foliis ſimplicibus lanceolatis integris. *Hort. cliff.* 212. *Vir. cliff.* 49. *Fl. ſuec.* 439, 424. *Roy. lugdb.* 483. *Dalib. pariſ.* 160.

Luteola herba ſalici folio. *Bauh. pin.* 100.

Lutum herba. *Dod. pempt.* 80.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 24.

Wau-Reſede. W.

Habitat in Europa *ad vias et pagos.* ⊙ (v. v.)

Folia *ad baſin utrinque denticulum habent calloſum.*

2. RESEDA *caneſcens.*

R. foliis lanceolatis undulatis piloſis. *Syſt. veg.* 448. *Vahl. ſymb.* 2. *p.* 52. *

R. (*hexagyna*) foliis lanceolatis repandis undulatis, floribus hexagynis. *Forſk. deſcr.* 92. *

R. foliis ſubulatis ſparſis. *Sauv. monſp.* 48. *Mill. dict.* 7.

R. foliis ſimplicibus oblongis integris. *Guett. ſtamp.* 1. *p.* 225.

R. alba minor. *Bauh. pin.* 100.

Seſamoides ſalmanticum parvum. 2. *Cluſ. hiſt.* 1. *p.* 296. *t.* 295.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 26.

Graue Reſede. W.

Habitat in Salmantica. ♃

Radix *perennis.* Caulis *pedalis, erectiuſculus, piloſus, pilis albis.* Rami *ex alis ſuperioribus ſeriores.* Folia *alterna, ſeſſilia, lanceolata, carina piloſa, vix ciliata, integerrima absque dente, undulata.* Racemus *terminalis:* Floribus *albis* Antherae *luteae* 11. Germina 6, *inſidentia ſtipiti communi.*

Synonyma Sauvageſii et Cluſii forte ad aliam incognitam ſpeciem ſpectant. W.

4. RESEDA *glauca.*

R. foliis linearibus baſi dentatis, floribus tetragynis. *Sp. pl.* 644.

R. lineariae foliis. *Bauh. pin.* 100. *prodr.* 41. *Burſ. IV.* 88.

R. linariae folio, glaucum pyrenaicum, flore ſtamineo. *Moriſ. hiſt.* 3. *p.* 601. *ſ.* 15. *t.* 1. *f.* 4.

R. linariae folia pyrenaica, glauco folio. *Pluk. alm.* 317. *t.* 107. *f.* 2. *Raj. ſuppl.* 511.
Luteola pumila pyrenaica, linariae folio. *Tournef. inſt.* 424.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 27.
Meergrüne Reſede. W.
Habitat in Pyrenaeis. ♃ (v. v.)

*4. RESEDA *dipetala.*

R. foliis linearibus integerrimis, floribus tetragynis dipetalis, petalis indiviſis. *Aiton. Kew.* 2. *p.* 132. *Vahl. ſymb.* 2. *p.* 52.
R. (*capenſis*) caule fruticoſo, foliis linearibus ternis apice mucronatis. *Burm. prodr. cap.* 13.
Flachsblättrige Reſede. W.
Habitat ad Cap. b. ſpei. ♂ (v. ſ.)
Facies *R. ſeſamoides.* Caulis *ſuffruticoſus erectus ramoſus.* Rami *teretes ſparſi glabri.* Folia *ſparſa linearia unguicularia craſſiuſcula acuta glabra.* Racemus *terminalis.* Flores *remotiuſculi.* Calycis *foliola ſex minuta albo-marginata.* Petala *cuneata integerrima.* Vahl. l. c.

5. RESEDA *purpuraſcens.*

R. foliis linearibus obtuſis, floribus pentagynis. *Sp. pl.* 644.
R. alba minor. 1. *Bauh. pin.* 100. *Raj. hiſt.* 1054.
Seſamoides ſalmanticum parvum. 1 *Cluſ. hiſt.* 1. *p.* 296.
Seſamoides foliis craſſis, floribus ex herbaceo purpuraſcentibus. *Tournef. inſt.* 424.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 27.
Röthliche Reſede. W.
Habitat in collibus Salmanticis, Monſpelii.

6. RESEDA *Seſamoides.*

R. foliis lanceolatis integris, fructibus ſtellatis. *Sp. pl.* 644.
Seſamoides fructu ſtellato. *Tournef. inſt.* 424.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 27.
Sternſamige Reſede. W.
Habitat Monſpelii *in Horto Dei.*

7. RESEDA *fruticuloſa.*

R. foliis pinnatis apice recurvatis, floribus tetragynis, calycibus quinquepartitis patentibus, caule baſi fruticoſo.

Sp. pl. 645. *Iacq. Collect.* 3. *p.* 195.* *Icon. rar.* 3. *t.* 474.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 28.
Strauchartige Reſede. W.
Habitat in Hiſpania. ♃

Radix *ſ. caulis ſupra terram perennans, craſſitie ſaepe duplo pollicis, breviſſimus, e cuius apice* Caules *plures, adſcendentes, laeves ſtriis elevatis.* Folia *pinnata: pinnis* 5 *ſ.* 7, *lanceolatis, decurrentibus, fere undatis, laevibus, apice recurvatis.* Spicae *plures, alternae, verſus caulis apicem primores.* Petala 6 *ſ.* 5, *alba, trifida, calyce majora.* Stamina 11, *quaſi media inter R. undatam et albam.*

8. RESEDA *alba.*

R. foliis pinnatis, floribus tetragynis, calycibus ſexpartitis. *Hort. upſ.* 149. *Mill. dict.* 5. *Willich. obſ. n.* 41. *Hoff. germ.* 163. *Roth. germ. I.* 204. *II.* 522.
R. foliis pinnatis integerrimis. *Hort. cliff.* 212.
R maxima. *Bauh. pin.* 100. *Lob. hiſt.* 110.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 28.
Weiſse Reſede. W.
Habitat Monſpelii *in* Hiſpania. ⊙ (v. v.)

9. RESEDA *undata.*

R. foliis pinnatis undulatis, floribus trigynis tetragynisve. *Mill. dict. n.* 4. *Kniph. cent.* 9. *n.* 84. *Vahl. ſymb.* 2. *p.* 52.
R. (*decuſſiva*) foliis decuſſive pinnatis, foliolis ſubaequalibus, floribus trigynis. *Forſk. cat. pl. aegypt. p.* 66.
R. minor foliis inciſis. *Barr. rar.* 78. *t.* 588.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 29.
Wellenblättrige Reſede. W.
Habitat in Hiſpania. ♃ (v. v.)

Caulis *pedalis, ſtriato-angulatus, ſtrictus.* Folia *radicalia et caulina pinnata: foliolis lanceolatis, decurrentibus, acutis, undatis: alternis minoribus, erectioribus. Superiora folia excludunt foliola alterna minora.* Racemi *terminales.* Calyx 5-*partitus.* Petala 5, *trifida, alba.* Antherae *lutae,* 10. Piſtilla *in aliis floſculis* 3, *in aliis* 5. Capſulae *kujus generis maximae.*

10. RESEDA *lutea.*

R. foliis omnibus trifidis: inferioribus pinnatis. *Hort. cliff.* 212. *Hort. upſ.* 149. *Roy. lugdb.* 483. *Dalib.*

pariſ. 159. *Gort. gelr.* 310. *Mill. dict. n.* 1. *et.* 2. *Jacq. auſt. t.* 352. *Scop. carn. n.* 569. *Pollich pal. n.* 454. *Gmel. tub. p.* 137. *Moench. haſſ. n.* 393. *Mattuſch. ſil. n.* 338. *Hoffm. germ.* 163. *Roth. germ. I.* 204. *II.* 523.

R. hexapetala, foliis pinnatis, undulatis, calyce ſexfido. *Hall. helv. n.* 1056.

R. vulgaris. *Bauh. pin.* 100. *Raj. hiſt.* 1055.

R. lutea. *Bauh. hiſt.* 3. *p.* 467.

β. Reſeda gallica criſpa. *Bocc. ſic.* 77. *t.* 41. *f.* 3.

R. maſſilienſis, foliis latioribus criſpis. *Pluk. alm.* 317. *t.* 55. *f.* 4.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 30.

Gelbe Reſede. W.

Habitat in Europae *auſtralioris montibus cretaceis.* ⊙ (v. v.)

Folia *ſaepius undulata.*

11. RESEDA *Phyteuma.*

R. foliis integris trilobisque, calycibus ſexpartitis maximis. *Hort. cliff.* 412. *Dalib. pariſ.* 159. *Mill. dict. n.* 3. *Jacq. auſt. t.* 132. *Willich. in nov. Act. A. N. C. tom.* 4. *p.* 107. *Scop. carn.* 2. *n.* 570.

R. foliis radicalibus integris, caulinis integris et ſemitrilobis, calycibus ſexfidis, florem ſuperantibus. *Hall. helv. n.* 1057.

R. floribus trigynis, calycibus maximis hexaphyllis. *Hort. upſ.* 150.

R. minor vulgaris. *Tournef. inſt.* 423.

Reſedae affinis Phyteuma. *Bauh. pin.* 100. *prodr.* 42. *t.* 42. *Bauh. hiſt.* 3. *p.* 306. *quoad iconem.*

Erucago apula trifida quinquefolia. *Col. ecphr.* 267. *t.* 269.

β. Reſeda minor vulgaris: folio minus inciſo. *Tournef. inſt.* 423.

γ. Reſeda minor vulgaris, foliis integris. *Tournef. inſt.* 423.

Phyteuma. *Bauh. pin.* 3. *p.* 386. *fig.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 31.

Kleine Reſede. W.

Habitat in Gallia, Italia, Oriente, Auſtria, Helvetia. ⊙

12. RESEDA *mediterranea.*

R. foliis integris trilobisque, calycibus flore brevioribus.

Mant. 564. *Iacq. collect.* 1. *p.* 147.* *Icon. rar.* 3. *t.* 475. *Vahl. ſymb.* 2. *p.* 53.

R. (*tetragyna*) tetragyna alba, foliis lanceolatis repandis, interdum baſi pinnatim inciſis calycis dente ſupremo aequali. *Forſk. deſcr.* 90.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 31.

Mittländiſche Reſede. W.

Habitat in Palaeſtina. ⊙ (v. v.)

Caulis *pedalis, adſcendens ſ. erectus, ſuperne ramoſus, ſcaber.* Folia *inferiora lanceolata, alterna, indiviſa, ſubtus ſcabra; caulina ſuperiora trifida.* Racemus *terminalis, longus, erectus.* Calyx *hexaphyllus, breviſſimus, linearis, patens.* Corolla *hexapetala, alba: Petala* 2 *ſuperiora tripartita: media minore;* 2 *laterialia bipartita: inferiore minore;* 2 *infima linearia.* Stamina *pallida.* Germen *tricuſpidatum, calyce vix longius.* Styli *nulli.* Stigmata *ſimplicia.*

13. RESEDA *odorata.*

R. foliis integris trilobisque, calycibus florem aequantibus. *Weig. obſ. bot. p.* 30. *obſ.* 15. *Kniph. cent.* 10 *n.* 73.

R. foliis integris trilobisque, floribus tetragynis. *Mill. dict. t.* 217.

R. foliis integris, floribus odoratis. *Hall. goett.* 95. *Zinn. goett.* 123.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 32.

Gemeine Reſede. W.

Habitat in Aegypto ♂ (v. v.)

Simillima R. Phyteumae, ſed Folia *magis undulata.* Calyces *corolla non longiores.* Rami *declinati, ſetis vagis adſperſi.* Flores *triſtes ambroſiaci, ſaepius tetragyni.* Petalorum *laciniae laterales in hac, intermediae vero in R. phyteumate, breviores.*

959. EUPHORBIA.

Gen. pl. ed. Schreb. n. 823.

Cor. 4-ſ. 5-petala, calyci inſidens. *Cal.* 1-phyllus, ventricoſus. *Capſ.* 3-cocca.

* *Fruticoſae, aculeatae.*

1. EUPHORBIA *antiquorum.*

E. aculeata ſubnuda triangularis articulata: ramis paten-

tibus. *Hort. cliff.* 196.* *Hort. upſ.* 138. *Fl. zeyl.* 199. *Roy. lugdb.* 194. *Amoen. acad.* 3. *p.* 106.* *Mill. dict. n.* 1. *Forſkaehl. aegypt. p.* 93. *Blackw. t.* 339.

Euphorbium antiquorum verum. *Comm. hort.* 1 *p.* 23. *t.* 12.

Schadidacalli. *Rheed. mal.* 2. *p.* 81. *t.* 42.

β. Euphorbia aculeata nuda triangularis articulata, ramis erectis. *Mill. dict.*

Euphorbium trigonum et tetragonum ſpinoſum, ramis compreſſis. *Iſn. act.* 120.

Tithymalus aizoides triangularis et quadrangularis articuloſus et ſpinoſus, ramis compreſſis. *Comm. prael.* 55. *t.* 5.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 34.

Wahre Wolfsmilch. W.

Habitat in India ♄

Stamina *tantum* 5 *numerat Forſkaehl.* l. c. R.

2. EUPHORBIA *canarienſis.*

E. aculeata nuda ſubquadrangularis, aculeis geminatis uncinatis, floribus ſubſeſſilibus. W.

E. aculeata nuda ſubquadrangularis, aculeis geminatis, *Hort. cliff.* 196.* *Hort. upſ.* 138. *Roy. lugdb.* 194. *Amoen. acad.* 3. *p.* 107.* *Mill. dict. n.* 2. *Blackw. t.* 340. *f.* 1.

Tithymalus aizoides fruticoſus canarienſis aphyllus quadrangularis et quinquangularis, ſpinis geminis aduncis atro-nitentibus armatus. *Comm. hort.* 2. *p.* 207. *t.* 104.

Tithymalus aizoides lactifluus ſ. Euphorbia canarienſis quadrilatera et quinquelatera, cerei effigie. *Pluk. alm.* 370. *t.* 320. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 35.

Canariſche Wolfsmilch. W.

Habitat in Canariis. ♄

Flos *infra par aculeorum, ſeſſilis, utrinque* Bractea *breviore ovata concava viridi.* Calyx *clauſus laciniis* 5, *conniventibus, purpureis.* Petala 5, *extra calycem poſita, obſcure purpurea, ſeſſilia, carnoſa, integra, obtuſiſſima.*

*3. EUPHORBIA *viroſa.* W. †

E. aculeata nuda multangularis, aculeis geminatis ſubulatis, flore pedunculato. W.

Euphorbia. *Patterſon. itin. ed. germ p. 60. t. 9. 10.*
Giftigſte Wolfsmilch. W.
Habitat in interioribus regionibus Cap. b. ſpei. ♄
Similis praecedenti, ſed diſtincta: margine *angulorum caulis plano*, aculeis *multo longioribus ſubulatis rectis*, *et* flore *pedunculato Convenit vero floris ſtructura, ad baſin floris ſunt* bracteae *binae ellipticae obtuſae, et petala ſunt obtuſiſſima.* W

4. EUPHORBIA *heptagona.*
E. aculeata nuda ſeptemangularis: ſpinis ſolitariis ſubulatis floriferis. *Hort. cliff.* 196.* *Roy. lugdb.* 194. *Amoen. acad.* 3. *p.* 109.* *Mill. dict. n* 6
Euphorbium heptagonum, ſpinis longiſſimis in apice frugiferis. *Boerh. lugdb.* 1. *p.* 258. *t.* 258.
Euphorbium capenſe, ſpinis longis ſimplicibus. *Bradl. ſucc.* 2. *p.* 4. *t.* 13.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 36.
Siebeneckige Wolfsmilch. W.
Habitat in Aethiopia. ♄

5. EUPHORBIA *mammillaris.*
E. aculeata seminuda, angulis tuberculatis ſpinis interſtinctis, tuberculis junioribus folio inſtructis. W.
E. aculeata nuda: angulis tuberoſis ſpinis interſtinctis. *Amoen. acad.* 3. 108 *Mill. dict. n.* 8.
Euphorbium polygonum, aculeis longioribus ex tuberculorum internodiis prodeuntibus. *Iſn. act.* 1720. *p.* 386.
Tithymalus aizoides africanus, validiſſimis ſpinis ex tuberculorum internodiis provenientibus. *Comm. prael.* 59. *t.* 9.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 36.
Zitzenförmige Wolfsmilch.
Habitat in Aethiopia. ♄
Differt ab Heptagona, angulis duplicatis toruloſis; ſpinae ſolitariae inter ſingula tubercula longitudinaliter digeſta. Flores *non vidi; an ex ſpinarum apice, ut heptagonae?*

6. EUPHORBIA *cereiformis.*
E. aculeata nuda multangularis, ſpinis ſolitariis ſubulatis *Roy. lugdb* 195. *Amoen acad.* 2. *p.* 108. *Mill. dict. n.* 9. *Kniph. cent.* 10 *n.* 41.
Euphorbium cerei effigie, caulibus gracilioribus. *Boerh. lugdb.* 1. *p.* 258.

Euphorbium aphyllum anguloſum, florum coma denſiſſima. *Burm. afr.* 19. *t.* 9. *f.* 3.
Tithymalus africanus ſpinoſus, cerei effigie. *Moriſ. hiſt.* 3. *p.* 245. *Pluk. alm.* 370. *t.* 231. *f.* 1.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 37.
Fackeldiſtelartige Wolfsmilch. W.
Habitat in Aethiopia. ♄

7. EUPHORBIA *officinarum.*

E. aculeata nuda multangularis; aculeis geminatis. *Hort. cliff.* 196. * *Hort. upſ.* 138. *Roy. lugdb.* 195. *Amoen. acad* 3. *p.* 107. *. *Mat. med. p.* 120. *Mill. dict. n.* 4. *Blackw. t.* 340. *f.* 2. *Regn. bot.*
Euphorbium polygonum ſpinoſum, cerei effigie. *Iſnard. act.* 1720. *p.* 500. *t.* 10.
Euphorbium cerei effigie, caulibus craſſioribus ſpinis validioribus armatum, *Comm. hort.* 1. *p.* 21. *t.* 11. *Seb. theſ.* 1. *p.* 29. *t.* 19. *f.* 2.
Euphorbium. *Bauh pin.* 387.
β. Kolquall. *Bruce. itin. ed. angl. app. p.* 41. *c. fig. ed. germ.* 5. *p.* 52. *t.* 10. 11.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 37.
Harzbringende Wolfsmilch. W.
Habitat in Aethiopia et Africa *calidiore.* ♄ (v. v.)

Varietas β. *angulis caulis repandis differre videtur, ſed ex brevi confuſaque deſcriptione Clariſſ. Bruce, non liquet, an ſit diſtincta ſpecies, vel varietas.* W.

*8. EUPHORBIA *triaculeata.*

E. aculeata nuda, caule tereti ſulcato, aculeis ternis. *Vahl. ſymb.* 2. *p.* 53.
E. (*triaculeata*) caule tereti ſeptemſulcato, floribus ternis ſeſſilibus ex ala aculeorum. *Forſk. deſcr.* 94. *
Dreyſtachliche Wolfsmilch. W.
Habitat in Arabia felici *ad urbem* Muſa *frequens in planitie ſubhumida.* ♄

Aculeus *intermedius pollicaris, laterales quintuplo minores recurvi.* Vahl. l. c.

9. EUPHORBIA *nereifolia.*

E. aculeata ſeminuda, aculeis geminatis, angulis oblique tuberculatis ſuperne folioſis, foliis oblongis. W.
E. aculeata ſeminuda: angulis oblique tuberculatis. *Hort. cliff.* 196. * *Hort. upſ.* 139. *Fl. zeyl.* 200. *Roy. lugdb* 194. *Amoen. acad.* 3. *p.* 109. * *Mill. n.* 5.

Euphorbium afrum ſpinoſum, foliis latioribus non ſpinoſis. *Seb. theſ.* 1. *p.* 18. *t.* 9 *f.* 1.
Tithymalus aizoides arboreſcens ſpinoſus, caudice angulari, nerii folio. *Comm. prael.* 22. 56. *t.* 6. *Bradl. ſucc.* 3. *p.* 10. *t.* 28.
Tithymalus indicus ſpinoſus, nerii folio. *Comm. hort.* 1. *p.* 25. *t.* 13.
Ligularia. *Rumph. amb.* 4. *p.* 88. *t.* 40.
Ela Calli. *Rheed. mal.* 2. *p.* 83. *t.* 43.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 40.
Oleanderblättrige Wolfsmilch. W.
Habitat in India. ♄ (v. v. ſ. fl.)

*10. EUPHORBIA *Hyſtrix.*
E. caule tereti ſeminudo, ſuperne folioſo, foliis lanceolato-linearibus, pedunculis unifloris perſiſtentibus demum ſpinoſis. W.
E. (*Hyſtrix*) imbricata, tuberculis folio lanceolato-lineari inſtructis, pedunculis unifloris axillaribus ſpineſcentibus. *Jacq. hort. ſchoenb.* 2. *p.* 43. *t.* 207. *
E. (*loricata*) aculeata ſeminuda, caule tereti ſquamoſo baſi ſpinis ſolitariis muricato, foliis lanceolato-linearibus. *Lamarck. encycl.* 2. *p.* 412. *
Tithymalus africanus arboreſcens, ſquamato caule ſpinoſus. *Pluk. alm.* 370. *t.* 230. *f.* 5. *Moriſ. hiſt.* 3. *p.* 344. *Petiv. gazoph. t.* 86. *f.* 5'9.
Vieldornige Wolfsmilch. W.
Habitat ad Cap. b. ſpei. ♄
Planta *primi anni eſt inermis, ſecundi vero ſpinoſa, quum* pedunculi *longiſſimi uniflori poſt floris fructusque caſum perſiſtant, et in ſpinas mutentur, ut in Hedyſaro cornuto.* W.

*11. EUPHORBIA *tribuloides.*
E. aculeata ſeminuda quadrangularis, baſi diphylla, foliis petiolatis ſpathulatis. W.
E. (*tribuloides*) humillima diphylla aculeata quadrangularis, ſpinis geminatis patentibus. *Lamarck. encycl.* 2. *p.* 412. *
Bürzeldornartige Wolfsmilch. W.
Habitat in Canariis ♄
Caulis *e radice fuſiformi ortus, eſt biuncialis craſſitie unciali et paulo ultra, ovatus carnoſus quadrangularis et baſi foliis binis oppoſitis petiolatis ovato-ſpathulatis, breve marceſcentibus ſuffultus.* Aculei

ad angulorum marginem conferti geminati albidi. Lamarck. l. c.

*12. EUPHORBIA *ſtellata.* W. †

E. caulibus digitalibus nudis aggregatis flexuoſis aculeatis, aculeis geminatis. W.

Euphorbe a chenilles. *Vaill. itin. ed. germ. Francof.* 4. *p.* 245. *t.* 11.

Sternförmige Wolfsmilch. W.

Habitat in interioribus regionibus Cap. b. ſpei. ♄

Radix *tuberoſa magnitudine pugni.* Caules *ex apice plures digitales decumbentes flexuoſi cylindracei obtuſi, baſi attenuati, laeves, et ſerie duplici tuberculorum ſuperne inſtructi.* Aculei *bini breves e quolibet tuberculo.* W.

*13. EUPHORBIA *cucumerina.* W. †

E. aculeata elliptica obtuſa ſulcata, aculeis ſubſolitariis, pedunculis ſubtrifloris. W.

Euphorbe a concombre. *Vaill. itin. ed. germ. Francof.* 3. *p.* 410. *t.* 5.

Gurkenförmige Wolfsmilch. W.

Habitat in interioribus Cap. b. ſpei *regionibus.* ♄

Facies totius plantae eſt Cucumeris ſativi. Caulis *eſt ellipticus carnoſus obtuſus, porcis pluribus exaratus.* Aculei *ſolitarii et geminati ſubulati.* Pedunculi *in apice caulis videntur bi- et triflori.* W.

** *Fruticoſae, inermes.* Caulis *nec dichotomus, nec umbelliferus.*

*14. EUPHORBIA *meloformis.*

E. inermis ſubgloboſa multangularis. *Ait. Kew.* 2. *p.* 135.

Euphorbe a cote de melon. *Vaill. itin. ed. germ. Francof* 4. *p.* 245. *t.* 12.

Melonenförmige Wolfsmilch. W.

Habitat in Cap. b. ſpei. *interioribus regionibus.* ♄

Caudex *carnoſus ſubgloboſus, diametro triunciali, glaber multangulatus:* porcae 8, 10 *vel plures baſi latae carinatae:* carinae *florigerae, cicatricibus pedunculorum glandulisque alternatim notatae.* Pedunculi *cylindracei craſſitie pennae columbinae articulati, villis breviſſimis adſperſi, plerumque primum trichotomi, dein dichotomi, raro ſimplices.* Bracteae *ad di-*

viſuras pedunculorum et ad baſin ſinguli floris oppoſitae oblongae acutiuſculae adſperſae lineam longae. Calyx *monophyllus campanulatus bracteis paulo longior, limbo quinquefido*: laciniae *ovatae obtuſae inflexae concavae.* Petala *quinque, calyci inter lacinias inſerta, ſubrotundo-reniformia obtuſiſſima, laciniis calycis duplo majora patentia carnoſa convexa poris raris pertuſa viridia.* Filamenta *viginti et quod excurrit, receptaculo inſerta calyce paulo longiora ſubulata villoſa.* Antherae *flavae didymae.* Piſtillum *nullum.* Act. l. c.

15. EUPHORBIA *Caput meduſae.*

E. inermis imbricata, tuberculis folio lineari inſtructis, floribus ſubpedunculatis petalis palmatis. *Ait. Kew.* 2. *p.* 135.

E. inermis imbricata, tuberculis foliola lineari inſtructis. *Hort. cliff.* 197. *Hort. upſ.* 139. *Roy. lugdb.* 195. *Amoen acad.* 3. *p.* 110. *Mill. dict. n.* 8.

E. inermis ramoſa tuberculata, calycum laciniis externis ſupra planis ſubquadridentatis. *Lamarck. encycl.* 2. *p.* 413. *

Planta lactaria africana. *Comm. hort.* 1. *p.* 33. *t.* 17.

Tithymalus aizoides africanus, ſimplici ſquamato caule, *Comm. prael.* 57. *t.* 7.

β. Euphorbium anacanthum, anguſto Polygoni folio. *Iſnard. Act. pariſ.* 1720. *p.* 586. *Breyn. prodr.* 29. *t.* 19.

γ. Euphorbium humile procumbens, ramis ſimplicibus copioſis, caule craſſiſſimo tuberoſo. *Burm. afr.* 20. *t.* 10. *f.* 1.

δ. Euphorbium procumbens, ramis plurimis ſimplicibus ſquamoſis, foliis deciduis. *Burm. afr.* 17. *t.* 8.

ε. Euphorbium procumbens, ramis geminatis, caule glabro oblongo cinereo. *Burm. afr.* 18. *t.* 9. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 41.

Meduſenhaupt-Wolfsmilch. W.

Habitat in Aethiopia. ♄ (v. v.)

Diſtinctiſſima planta petalis palmatis.

*16. EUPHPORBIA *tuberculata.*

E. inermis imbricata, tuberculis compreſſis acutis foliolo lineari inſtructis, floribus longe pedunculatis, petalis palmatis. W.

E. (*tuberculata*) inermis imbricata, tuberculis patentiſ-

ſimis acutis foliolo lineari inſtructis, floribus pedunculatis, petalis palmatis. *Iacq. hort. Schoenbr.* 2, *p.* 43. *t.* 208.

Knotige Wolfsmilch. W.

Habitat ad Cap. b. ſpei. ♄

*17. EUPHORBIA *Anacantha.*

E. inermis imbricata, tuberculis foliolo ſubrotundo inſtructis, floribus terminalibus ſolitariis ſeſſilibus, petalis palmatis. *Ait. Kew.* 2. *p.* 136. *Mill. dict. n.* 7.

E. (*tridentata*) inermis ramoſa ſubtuberculata, calycum laciniis externis ſupra concavis coloratis tridentatis. *Lamarck. encycl* 2. *p.* 413. *

Euphorbium anacanthum ſquamoſum, lobis florum tridentatis. *Iſnard. Act. pariſ.* 1720. *p.* 387. *n.* 12. *et p.* 392. *tab.* 11.

Euphorbium erectum aphyllum, ramis rotundis, tuberculis tetragonis *Burm afr.* 16. *tab.* 7. *f.* 2.

Dreyzähnige Wolfsmilch. W.

Habitat ad Cap. b. ſpei. ♄

*18. EUPHORBIA *Clava.*

E. inermis imbricata, tuberculis foliolo lanceolato inſtructis, floribus pedunculatis, petalis integerrimis. *Ait. Kew* 2. *p.* 156.

E. (*Clava*) inermis fruticoſa, foliis ſpathulato-linearibus, pedunculis axillaribus unifloris bracteatis. *Iacq. ic.* 1. *t.* 85 *Collect.* 1. *p.* 104. *

E. (*canaliculata*) inermis, caule ſimplici craſſo tuberculato, foliis linearibus acutis canaliculatis, pedunculis unifloris. *Lamarck. encycl.* 2. *p.* 413. *

Euphorbium acaulon erectum tuberoſum, linearibus foliis, flore ac fructu foliaceis. *Burm. afr.* 12. *t.* 6. *f.* 1.

Tithymalus aizoides africanus ſimplici ſquamato caule, chamaenerii folio. *Commel. praelud.* 58. *t.* 8.

Keulenförmige Wolfsmilch. W.

Habitat ad Cap. b. ſpei. ♄

*19. EUPHORBIA *bupleurifolia.*

E. inermis capitata imbricata, tuberculis ſubrhombeis, foliis lanceolatis petiolatis, pedunculis unifloris, petalis integerrimis, bracteis flore longioribus. W.

E. inermis ſubacaulis, foliis lanceolatis. *Iacq. hort. Schoenbr.* 1. *p.* 55. *t.* 106. *

Haſenoehrleinblättrige Wolfsmilch. W.

Habitat ad Cap. b. ſpei. ♄

*20. EUPHORBIA *lophogona.*

E. inermis caule pentagono alato, foliis lanceolato-ſpathulatis, umbella dichotoma terminali. W.

E. (*lophogona*) inermis ſeminuda, caule ſimplici pentagono, angulis membranaceis et quaſi criſtatis, foliis lanceolato-ſpathulatis, umbella dichotoma terminali. *Lamarck. encycl.* 2. *p.* 414.

Flügelſtielige Wolfsmilch. W.

Habitat in Madagaſcar. ♄

Similis Euphorbiae neriifoliae ſed abunde diſtincta: magnitudine, aculeorum *defectu*, caule *alato-angulato*, *et* inflorefcentia. Radix *ramoſa fibroſa.* Caulis *pedalis et ultra, ſimpliciſſimus fruticoſus craſſitie digiti, baſi cylindraceus, ſuperne ſpiraliter quinquangularis, angulis membranaceis laciniatis.* Folia *in ſummitate duntaxat caulis oblonga obtuſa ſpathulata, baſi anguſtata quadri-ſeu ſexpollicaria glabra.* Pedunculus *terminalis bipollicaris bis vel ter dichotome diviſus.* Umbellae *ſex vel octoflorae terminales.* Bracteae *binae ſubrotundae mucronatae albae petaliformes ad cujusque floris baſin.* Lamarck. l. c.

21. EUPHORBIA *mauritanica.*

D. inermis ſeminuda fruticoſa filiformis flaccida, foliis alternis. *Hort. cliff.* 197. *Hort. upſ.* 140. *Roy. lugdb.* 195. *Amoen. acad.* 3. *p.* 111.* *Gron. orient.* 160. *Mill. dict. n.* 16.

Tithymalus aphyllus mauritaniae. *Dill. elth.* 384. *t.* 289. *f.* 373.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 42.

Mauritaniſche Wolfsmilch. W.

Habitat in Africae *maritimis.* ♄ (v. v.)

*22. EUPHORBIA *piſcatoria.*

E. inermis fruticoſa ſtricta, umbellis quinquefidis terminalibus, involucellis oblongis, foliis lanceolatis laevibus. *Ait. Kew.* 2. *p.* 137.

Fiſcher-Wolfsmilch. W.

Habitat in Madera *et* Canariis. ♄ (v. ſ.)

*23. EUPHORBIA *balſamifera.*

E. inermis fruticoſa ſtricta, capitulo terminali, foliis lanceolatis laevibus glaucis. *Ait. Kew.* 2. *p.* 137.

Balſamiſche Wolfsmilch. W.

Habitat in Canariis. ♄

24. EUPHORBIA *Tirucalli.*

E. inermis ſeminuda fruticoſa filiformis erecta, ramis patulis determinate confertis. *Hort. cliff.* 197. *Hort. upſ.* 139. *Fl. zeyl.* 197. *Roy. lugdb.* 195. *Amoen. acad* 3. *p.* 111. * *Mill. dict. n.* 14.

Tithymalus indicus fruteſcens. *Comm. hort.* 1. *p.* 27. *t.* 14.

Tithymalus arboreſcens, caule aphyllo. *Pluk. phyt.* 319. *f.* 6.

Oſſifraga lactea. *Rumph. amb.* 7. *p.* 62. *t.* 29.

Tiru Calli. *Rheed. mal.* 8. *p.* - - . *t.* 44.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 43.

Malabariſche Wolfsmilch. W.

Habitat in India. ♄

25. EUPHORBIA *laurifolia.*

E. inermis foliis lanceolato-oblongis glabris, pedunculis axillaribus trifloris. W.

E. ſeminuda, foliis ſparſis confertis lanceolato-oblongis integerrimis laevibus, pedunculis axillaribus trifloris. *Lamarck. encycl.* 2. *p.* 415. *

Lorbeerblättrige Wolfsmilch. W.

Habitat in Peru. ♄

Fructus *magnitudine et facie Hippomanes biglanduloſae glabri.* Lamarck. l. c.

26. EUPHORBIA *pyrifolia.*

E. inermis, foliis lanceolato-oblongis mucronatis glabris, pedunculis ſubumbellatis. W.

E. inermis, foliis lanceolato-ovalibus integerrimis petiolatis mucronatis laevibus, pedunculis longis nudis ſubumbellatis. *Lamarck. encycl.* 2. *p.* 415. *

Birnenblättrige Wolfsmilch. W.

Habitat in inſula Mauritii. ♄

Pedunculi *bipollicares nudi interdum uniflori, flore uno alterove ad baſin prolifero interdum* 3 *ſ.* 5 *flori umbellati interdum et umbella baſi eſt prolifera.* Capſulae *glabrae.* Lamarck. l. c.

27. EUPHORBIA *Tithymaloides.*

E. inermis fruticoſa, foliis diſtiche alternis ovatis. *Hort. cliff.* 198. * *Roy lugdb.* 95. *Amoen. acad.* 3. *p.* 111. * *Jacq. amer.* 149. *t.* 92.

E. tithymaloides, *myrtifolia.*

E. (*myrtifolia*) inermis fruticosa, foliis alternis ovato-acutis planis ſublucidis. *Lamarck. encycl.* 2. *p.*416.*

Tithymalus (*myrtifolius*) foliis ovatis acuminatis. *Mill. dict. n.* 1.
Tithymalus curaſſavicus myrtifolius, flore coccineo mellifluo. *Herm. par.* 234. *t.* 234.
Tithymalus curaſſavicus myrtifolius, flore papilionaceo coccineo parvo. *Comm. hort.* 1. *p.* 31. *t.* 16. *Pluk. alm.* 369. *t.* 230. *f.* 2.

β. Euphorbia *padifolia.*

E. (*anacampſeroides*) inermis fruticoſa, foliis alternis obovatis obtuſis, ſubtus acute carinatis. *Lamarck. encycl.* 2. *p.* 416. *
Tithymalus (*lauroceraſifolius*) foliis oblongo-ovatis obtuſis ſucculentis. *Mill. dict. n.* 2.
Tithymaloides lauroceraſi folio non ſerrato. *Dill. elth.* 383. *t.* 288. *f.* 372.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 44. *t.* 51. *f. B.*
Saftige Wolfsmilch. W.
Habitat in America *calidiore.* ♄

28. EUPHORBIA *heterophylla.*

E. inermis, foliis ſerratis panduriformibus, ſummis lanceolatis, floribus terminalibus ſubumbellatis. W.
E. inermis, foliis ſerratis petiolatis difformibus ovatis lanceolatis panduriformibus. *Amoen. acad.* 3. *p.* 112. *Mill. dict. n.* 30.
Tithymalus heterophyllus. *Plum. ic.* 250. *t.* 251. *f.* 3.
Tithymalus curaſſavicus, ſalicis et atriplicis foliis variis, caulibus viridantibus. *Pluk. alm.* 369. *t.* 112. *f.* 6.
Tithymalus curaſſavicus, ſalicis et atriplicis foliis hirſutis, caulibus ſubrubentibus. *Moriſ. hiſt.* 3. *p.* 336.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 46.
Verſchiedenblättrige Wolfsmilch. W.
Habitat in America *calidiore.* ♄

*29. EUPHORBIA *cyathophora.*

E. inermis, foliis panduriformibus ovatis, floribus terminalibus ſubumbellatis, involucellis coloratis. W.
E. (*cyathophora*) umbella trifida multifida appendice calycino cyathiformi, foliis pleriſque panduriformibus, infimis caulinis ovatis. *Murr. comment. goett.* 7. *p.* 81. *t.* 1. *.
E. (*cyathophora*) umbella trifida ſubdichotoma, foliis panduraeformibus. *Iacq. ic. rar.* 3. *t.* 480.
Euphorbia heterophylla. *Iacq. Collect.* 1. *p.* 157. *

Geigenblättrige Wolfsmilch. W.

Habitat in America *calidiore.* ♄ (v. v.)

Hanc poſt E. heterophyllam indicavi ob affinitatem et inflorescentiam, revera non exacte umbellatam. Diſtincta a praecedente foliis *ſuperioribus nunquam lanceolatis. Variat involucris rubris et albis.* W.

*30. EUPHORBIA *nudiflora.*

E. inermis fruticoſa, foliis ovatis integerrimis, cyma axillari nuda. W.

E. (*nudiflora*) ſubtrichotoma, foliis integerrimis ovatis, ſubundulatis pedunculis cymoſis ramoſis, caule fruticoſo. *Iacq. ic. rar.* 3. *t.* 479. *Collect.* 3. *p.* 180. *

Nacktblütige Wolfsmilch. W.

Habitat - - - - ♄

Frutex *humanae altitudinis ramis ſaepius trichotomis.* Folia *oppoſita et terna ovata margine ſubundulata glabra.* Cymae *ramoſae in foliorum axillis abſque involucris.* Petala *ſubrotunda tenuiſſima crenata.* Styli *connati.* Capſulae *glabrae.* W.

31. EUPHORBIA *cotinifolia.*

E. foliis oppoſitis ſubcordatis petiolatis emarginatis integerrimis, caule fruticoſo. *Amoen. acad.* 3. *p.* 112. *Mill. dict. n.* 17. *Kniph. cent.* 2. *n.* 24.

E. inermis, caule fruticoſo, foliis oppoſitis ſubcordatis emarginatis, petiolis folio longioribus. *Hort. cliff.* 198. *Roy. lugdb.* 196.

Tithymalus curaſſavicus, folio cotini triphyllos, petalis florum ſerratis. *Pluk. alm.* 369. *t.* 230. *f.* 3.

Tithymalus arboreus curaſſavicus, cotini folio. *Seb. theſ.* 1. *p.* 75. *t.* 46. *f.* 4.

Tithymalus arboreus americanus, cotini folio. *Comm. hort.* 1. *p.* 29. *t.* 15.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 47.

Perrückenbaumblättrige Wolfsmilch. W.

Habitat in Curaſſao. ♄ (v. v.)

*32. EUPHORBIA *mellifera.*

E. foliis ſparſis lanceolatis acutis laevibus, pedunculis dichotomis, capſulis muricatis. *Ait. Kew.* 3. *p.* 493.

Honigbringende Wolfsmilch. W.

Habitat in Madera. ♄ (v. v.)

Caulis *fruticoſus laevis.* Folia *alterna lanceolata utrinque attenuata integerrima glabra, iis Nerii Ole-*

andri ſimillima, coſtata, coſta media ſubtus elevata, in junioribus piloſa. Corymbus *terminalis multiflorus, pedunculis inferioribus et ſummis trifidis, intermediis bifidis.* Flores *aurantii coloris.* Petala *reniformia indiviſa.* Styli *tres, apice bifidi.* Stigmata *capitata.* Capſulae *verrucoſae.* W.

*33. EUPHORBIA *glabrata.*

E. inermis truteſcens ramoſa, foliis oppoſitis cordato-ovatis, acutis glabris integerrimis. *Swartz prodr.* 96. *Fl. ind. occid.* 2. *p.* 870.* *Vahl. ſymb.* 2. *p.* 53.*

E. dichotoma, foliis ovatis integerrimis oppoſitis, caule geniculato, pedunculis capitatis axillaribus. *Houſt. cat* 213.

Peplis fruticoſa maritima geniculata. *Sloan. cat.* 82. *hiſt.* 2. *p.* 198.

Glatte Wolfsmilch. W.

Habitat in littoribus petroſis Jamaicae, *inque aliis inſulis* Caribaeis. ♄ (v. ſ.)

Non lacteſcit. Swartz l. c.

*34. EUPHORBIA *linarifolia.* W.

E. inermis fruteſcens, foliis ſparſis lanceolatis mucronatis, flore ſolitario terminali involucro triphyllo cincto, capſulis muricatis. W.

E. (*linifolia*) fruticoſa inermis, foliis ſparſis lanceolatis mucrone innocuo terminatis, terminalibus ternis. *Iacq ic. rar.* 1. *t.* 86. *Collect.* 1. *p.* 135.*

Löwenmaulblättrige Wolfsmilch. W.

Habitat in - - - - ♄ (v. ſ.)

Folia *oppoſita et alterna lanceolata obtuſa breviſſimo mucrone inſtructa.* Capſulae *e ſquamis truncatis muricatae.* W.

*35. EUPHORBIA *linifolia.*

E. inermis ſuffruticoſa, foliis infimis oppoſitis ovalibus, reliquis alternis linearibus, pedunculis ſubquinis terminalibus, capſulis glabris. W.

E. (*linifolia*) caule ſuffruticoſo, foliis oppoſitis alternisque lineari-lanceolatis. *Vahl. ſymb.* 2. *p.* 53.*

Flachsblättrige Wolfsmilch. W.

Habitat in inſula St. Dominici. ♄

*36. EUPHORBIA *cuneata.*

E. fruticoſa, foliis obovatis, pedunculis lateralibus trifloris. *Vahl. ſymb.* 2. *p.* 53.*

Keilblättrige Wolfsmilch. W.
Habitat in Arabia *felici.* ♄
Capſula *villoſa.* Petala *obovata truncata.* Vahl. l. c.

37. EUPHORBIA *ocymoidea.*
E. inermis herbacea ramoſa, foliis ſubcordatis integerrimis petiolo brevioribus, floribus ſolitariis. *Amoen. acad.* 3. *p.* 112. *Mill. dict. n.* 32.
E. inermis, foliis ſubcordatis obtuſis integerrimis petiolatis, caule ramoſo erecto. *Roy. lugdb.* 199.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 47.
Baſilicumartige Wolfsmilch. W.
Habitat in Campechia. *Houſt.*

*** *Dichotomae (umbella bifida aut nulla.)*

*38. EUPHORBIA *laevigata.*
E. fruteſcens ramoſa, foliis oppoſitis oblongis obtuſis integerrimis. *Vahl. ſymb.* 2. *p.* 54.
Meergrüne Wolfsmilch. W.
Habitat in India *orientali.* ♄
Tota glabra. Caulis *inermis.* Rami *apice dichotomi ter vel quater diviſi.* Folia *breviter petiolato unguicularia articulis ramuorm longiora, altero latere baſeos breviora anguſtiora, apice mucrone minuto terminata laevia avenia glauca viridia.* Pedunculi *ſuperne ex dichotomia ramulorum ſolitarii.* Flores *parvi.* Capſula *glabra laevis magnitudine ſeminis Coriandri.* Vahl. l. c.

39. EUPHORBIA *origanoides.*
E. dichotoma, foliis ſerrulatis ovatis obtuſis trinerviis, panicula terminali, caulibus ſimplicibus. *Amoen. acad.* 2. *p.* 114.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 48.
Doſtartige Wolfsmilch. W.
Habitat in Inſula Aſcenſionis. ♃
Planta refert ita Origanum, ut primo intuitu nequeat non pro eo agnoſci.
Caulis *ſimplex ſpithamaeus teres articulatus.* Folia *rotundo-ovata trinervia ſeſſilia glabra ſerrulata, paginae ſuperioris latere exteriori purpuraſcente.* Rami *nulli.* Panicula *ut in Origano terminat caulem.* Amoen. acad.

*40. EUPHORBIA *Atoto.*
E. dichotoma, foliis ovatis integerrimis umbella terminali. *Forſt. prodr. n.* 207.

Südſee-Wolfsmilch. W.
Habitat in Societatis *inſulis.*

41. EUPHORBIA *hypericifolia.*

E. dichotoma, foliis ſerratis ovali-oblongis glabris, corymbis terminalibus, ramis divaricatis. *Amoen. acad.* 3. *p.* 113.* *Mill. dict. n.* 31. *Swartz obſ.* 194.*

E. inermis, foliis ovalibus oppoſitis hinc ſerratis uniformibus, ramis alternis, caule erectiuſculo. *Hort. cliff.* 198. *Hort. upſ.* 143.* *Roy. lugdb.* 196.

E. minima reclinata, foliolis ovatis denticulatis, floribus ſubumbellatis terminalibus lateralibusque. *Brown. jam.* 235.

Tithymalus americanus, floſculis albis. *Comm. prael.* 60. *t.* 60.

Tithymalus erectus acris, parietariae foliis glabris, floribus ad caulium nodos conglomeratis. *Sloan. jam.* 82. *hiſt.* 1. *p.* 197. *t.* 126.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 48.
Johanniskrautblättrige Wolfsmilch. W.
Habitat in Indiae *occidentalis cultis.* ⊙ (v. v.)

*42. EUPHORBIA *proſtrata.*

E. dichotoma, foliis ovalibus obſolete ſerratis, pedunculis axillaribus ſubtrifloris, caulibus diffuſis glabris. *Ait. Kew.* 2. *p.* 139.

Liegende Wolfsmilch.
Habitat in India *occidentali.* ⊙

Caules *herbacei ſpithamaei procumbentes teretes ramoſi rubicundi.* Folia *oppoſita breviſſime petiolata obtuſiuſcula glabra obſolete trinervia, ſupra viridia ſubtus glaucescentia trilinearia.* Flores *axillares breviter pedicellati ſaepe terni, interdum ſolitarii.* Petala *purpurea.* Ait. l. c.

*43. EUPHORBIA *roſea.*

E. dichotoma diffuſa, foliis obovatis obliquis baſi ſubcordatis apice denticulatis, pedunculis unifloris axillaribus. W.

E. dichotoma, foliis obovatis baſi obliquis apice denticulatis, caule depreſſo diffuſo. *Retz. obſ.* 4. 26.*

Roſenfarbige Wolfsmilch. W.
Habitat in Indiae orientalis *arenoſis.* ⊙ (v. ſ.)

Variat valde magnitudine. Folia *avenia inferiora latiora et majora.* Flores *axillares in ramorum apice.*

Calyx *purpureus.* Petala *lineari-obovata obtuſa roſea.* Germen *papillis minutiſſimis obſitum.* Capſula *junior etiam papilloſa, adulta vero glabra.* W.

44. EUPHORBIA *maculata.*

E. dichotoma, foliis ſerratis oblongis piloſis, floribus axillaribus ſolitariis, ramis patulis. *Iacq. hort t.* 186. *Kniph. cent.* 8. *n.* 39.

Tithymalus ſ. Chamaeſyce altera virginiana, foliis crenatis et macula fuſca eleganter notatis. *Pluk. alm.* 372. *t.* 65. *f.* 8.

Chamaeſyce. *Sloan. jam.* 83. *hiſt.* 1. *p.* 198.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 49.

Gefleckte Wolfsmilch. W.

Habitat in America *ſeptentrionali.* ⊙ (v. v.)

Caules *dichotomi; Ramis alternis, patentibus, ſupra purpuraſcentibus.* Folia *ovali-oblonga, trinervia, ſubpiloſa, ſerrata, altero latere maxima parte integerrima, tenera adhuc planta notata macula fuſca.* Flores *axillares, ſolitarji, parvi, calyce rufo. Fructus laevis. Similis E. hypericifoliae.*

*45. EUPHORBIA *ſcordifolia.*

E. dichotoma, foliis ovatis obliquis obtuſis crenatis hirſutis, pedunculis axillaribus unifloris, petalis ſerratis, capſula hirſuta. W.

E. (*ſcordifolia*) dichotoma hirſuta, foliis crenulatis ſubovatis obtuſis, pedunculis unifloris ſolitariis. *Iacq. ic. rar.* 3. *t.* 476. *Collect.* 113. *.

Gamanderblättrige Wolfsmilch. W.

Habitat in Africa *ad* Senegal? ⊙

Petala *ſubrotundo-truncata apice dentata.* Styli *breviſſimi.* Capſula *hirſuta.* W.

*46. EUPHORBIA *picta.*

E. dichotoma, foliis ovatis hirſutis petiolatis integerrimis, pedunculis axillaribus unifloris, petalis ovatis, capſulis glabris. W.

E. (*picta*) dichotoma villoſa, foliis repandis ovatis pictis acutis, pedunculis unifloris ſolitariis, caule erecto. *Iacq. ic. rar. t.* 477. *Collect.* 3. *p.* 178. *

Habitat in inſula Venezula. ♃ (v. ſ.)

Folia *albo-maculata.* Petala *ovata integerrima alba.* Styli *ſex albi.* W.

47. EUPHORBIA *hirta.*

E. dichotoma, foliis ſerrulatis ovatis acuminatis, pedunculis capitatis axillaribus, caulibus piloſis *Amoen. acad.* 3. *p.* 114.* *Jacq. Collect. ſuppl.* 160. *t.* 11. *f.* 1.

E. inermis, foliis oppoſitis ovalibus ſerratis uniformibus, pedunculis capitatis axillaribus. *Fl. zeyl* 197.*

E. reclinata minor ſubhirſuta, foliis ſerratis. *Brown. jam.* 234.

Tithymalus botryoides zeylanicus, cauliculis villoſis. *Burm. zeyl.* 223. *t.* 104.

Eſula eſculenta. *Rumph. amb.* 6. *p.* 54. *t.* 23. *f.* 2.

Houttuyn Lin. Pfl. Syſt. p. 7 50.

Knopförmige Wolfsmilch. W.

Habitat in India. ⊙

Planta Illuſt. Jacquini non eſt varietas ſequentis, differt enim: caule *proſtrato*, foliis *ſtrigoſo - piloſis non obliquis*, capitulis *florum ſemper ſolitariis.* W.

48. EUPHORBIA *pilulifera.*

E. dichotoma, foliis ſerratis ovali-oblongis, pedunculis bicapitatis axillaribus, caule erecto. *Amoen. acad.* 3. *p.* 115. *Jacq. ic. rar.* 3. *t.* 478 *Collect.* 2. *p.* 361.*

Tithymalus botryoides erectus, florum capitulis conjugatis et longiori pediculo inſidentibus. *Burm. zeyl.* 224. *t.* 105. *f.* 1. *Pet. gaz. t.* 80. *f.* 14.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 50.

Pillentragende Wolfsmilch. W.

Habitat in India. ⊙ (v. ſ.)

*49. EUPHORBIA *braſilienſis.*

E. ramoſiſſima dichotoma, foliis ellipticis oblongis oppoſitis ſerrulatis, racemulis terminalibus. *Lamarck. encycl.* 2. *p.* 420.*

Braſiliſche Wolfsmilch.

Habitat in Braſilia.

Caulis *herbaceus decempollicaris pubeſcens.* Folia *obliqua, ramea multo minora.* Corymbi *terminales.* Calyx *fuſco-ruber campanulatus.* Petala *ſubrotunda alba.* Germen *pubeſcens.* Lamarck. l. c.

50. EUPHORBIA *hyſſopifolia.*

E. dichotoma, foliis ſubcrenatis linearibus, floribus faſciculatis terminalibus, caule erecto. *Sp. pl.* 651.

E. dichotomo erecta tenuis, foliis linearibus, floribus quaſi umbellatis terminalibus. *Brown. jam.* 235.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 51.,
Hopblättrige Wolfsmilch. W.
Habitat in America.

51. EUPHORBIA *thymifolia.*

E. dichotoma, foliis ſerratis ovali-oblongis, capitulis axillaribus glomeratis ſubſeſſilibus, caulibus procumbentibus. *Amoen. acad.* 3. *p* 115.

Tithymalus foliis oppoſitis oblique cordatis obtuſis ſerratis, pedunculis multifloris. *Fl. zeyl.* 198.

Tithymalus indicus annuus dulcis, floribus albis, caulibus viridantibus et rubentibus. *Pluk. alm.* 272. *t.* 113 *f* 2.

Tithymalus humilis ramoſiſſimus hirſutus, foliis thymi ſerratis. *Burm. zeyl.* 225. *t.* 105. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 51. *t.* 52. *f.* 1.
Thymianblättrige Wolfsmilch. W.

β. Euphorbia nodoſa. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 52. *t.* 52. *f.* 2.

Habitat in India. ☉ (v. ſ.)

52. EUPHORBIA *parviflora.*

E. ſubdichotoma, foliis ſerratis oblongis glabris, floribus ſolitariis, caule erectiuſculo alterne ramoſo. *Sp. pl.* 652.

Tithymalus erectus, floribus rarioribus, foliis oblongis glabris integris *Burm. zeyl.* 224. *t.* 105. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 52.
Kleinblütige Wolfsmilch. W.
Habitat in India. ☉ (v. ſ.)

Folia *ſaepe macula fuſca.* Calyces *purpurei.* Petala *alba.*

53. EUPHORBIA *caneſcens.*

E. dichotoma foliis integris ſubrotundis piloſis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Sp. pl.* 652. *Cavan. ic.* 1. *p.* 46. *t.* 63.* *Iacq. Collect. ſuppl.* 115. *t.* 2 *f.* 3.*

Tithymalus exiguus villoſus nummulariae folio. *Tournef. inſt.* 87.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 53.
Graue Wolfsmilch. W.
Habitat in Hiſpania. ☉ (v. ſ.)

Affinis nimis E Chamaeſyces, ſed tota undique albovilloſa. Folia obſolete emarginata, obſolete crenata. In Siphilitide *infuſum.*

54. EUPHORBIA *Chamaeſyce.*

E. dichotoma, foliis crenulatis ſubrotundis glabris, floribus ſolitariis axillaribus, caulibus procumbentibus. *Amoen. acad.* 3. *p.* 115.* *Swartz obſ.* 196.* *Hoffm. germ.* 164. *Roth. germ. I.* 204. *II.* 523.

E. inermis, foliis oblique cordatis ſerrulatis uniformibus, ramis alternis, floribus ſolitariis. *Hort. cliff.* 198.* *Roy. lugdb.* 196. *Gmel ſib.* 2. *p.* 237. *Gron. orient.* 160.

E. minima rufeſcens, foliolis ſubrotundis nitidis oppoſitis, ramulis floriferis foliatis ad alas alternans. *Brown. jam.* 236.

Chamaeſyce. *Bauh. pin.* 293. *Cluſ. hiſt.* 2. *p.* 187.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 53.

Niedrige Wolfsmilch. W.

Habitat in Europae *auſtralis*, Sibiriae, Meſopotamiae et Indiae *occidentalis aridis.* ⊙ (v. ſ.)

Caulis *depreſſus*, *dichotomus.* Folia *oppoſita*, *ſubrotunda*, *laevia*, *ſubcrenulata*, *emarginata*, *baſi antica anguſtiore.* Flores *e dichotomia* Calyx *rufus.* Petala *alba*, *crenata.* Fructus *mihi laevis.*

*55. EUPHORBIA *granulata.*

E. foliis oppoſitis oblongis integerrimis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Vahl. ſymb.* 2. *p.* 54.*

E. (*granulata*) foliis oppoſitis oblongis integerrimis, floribus axillaribus ſolitariis. *Forſk. deſcr.* 94.*

Körnige Wolfsmilch. W.

Habitat in Arabia *felici.*

56. EUPHORBIA *Peplis.*

E. dichotoma, foliis integerrimis ſemicordatis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Amoen. acad.* 3. *p.* 115. *Gron. orient.* 163. *Vahl ſymb.* 2. *p.* 54.*

E. (*dichotoma*) foliis oppoſitis integris lanceolatis ſeſſilibus. *Forſk. deſcr.* 93.*

E. foliis ſubrotundis obtuſis, floribus ſolitariis in foliorum alis. *Guett. ſtamp* 2. *p.* 420.

Tithymalu Peplis. *Scop. carn ed.* 2. *n.* 583.

Peplis maritima, folio obtuſo. *Bauh pin.* 293.

Peplis. *Cluſ. hiſt.* 2. *p* 187. *Cam epit.* 970.

Peplion. *Dalech hiſt* 970.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 54.

Meerſtrands-Wolfsmilch. W.
Habitat in Narbonae, Hiſpaniae, Carniolae *maritimis.* ⊙ (v. ſ.)

57. EUPHORBIA *polygonifolia.*
E. foliis oppoſitis integerrimis lanceolatis obtuſis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Amoen. acad.* 3. *p.* 116.* *Iacq. Collect. ſuppl.* 114. *t.* 13. *f.* 3.*
E. procumbens, ramulis alternis, foliis lanceolato-linearibus, floribus ſolitariis. *Gron. virg.* 58.
E. minima ramoſiſſima anguſtifolia. *Raj. ſuppl.* 431.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 54.
Knötrigblättrige Wolfsmilch. W.
Habitat in Canada, Virginia. ⊙

*58. EUPHORBIA *linearis.*
E. dichotoma, foliis oppſitis linearibus integerrimis, pedunculis ſolitariis. *Retz. obſ.* 3. *p.* 32.
Linienblättrige Wolfsmilch. W.
Habitat in inſula St. Crucis. (v. ſ.)
Caulis *teres nudus articulatus dichotomus.* Folia *oppoſita anguſto-linearia biuncialia integerrima acumine parvo.* Petala *integra.* Capſulae *laeves.* Retz. l. c.

59. EUPHORBIA *graminea.*
E. dichotoma, foliis lanceolato-ellipticis petiolatis integerrimis, caule erecto, pedunculis dichotomis. *Mant.* 74.
E. (*graminea*) dichotoma, foliis oppoſitis integerrimis lanceolatis, pedunculis dichotomis, caule erecto. *Iacq. amer.* 151. *obſ.* 1. *p.* 5. *t.* 31.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 55.
Wieſen-Wolfsmilch. W.
Habitat in Carthagenae *gramineis udis.*
Corollae *albae.*

60. EUPHORBIA *Ipecacuanhae.*
E. dichotoma, foliis integerrimis lanceolatis, pedunculis axillaribus unifloris folia aequantibus, caule erecto. *Amoen. acad.* 3. *p.* 117.*
Tithymalus, flore exiguo viridi apicibus flavis, antequam folia emittit florens. *Gron. virg.* 58.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 55.
Brechenmachende Wolfsmilch. W.
Habitat in Virginia, Canada. ♃

61. EUPHORBIA *portulacoides.*

E. dichotoma, foliis integerrimis ovalibus retuſis, pedunculis axillaribus unifloris folia aequantibus, caule erecto. *Amoen. acad.* 3. *p.* 117. *

Tithymalus perennis, portulacae folio. *Few. peruv.* 707. *t.* 2.?

Houttuyn Lin. Pfl. Syſt. 7. *p.* 56.

Portulacblättrige Wolfsmilch. W.

Habitat in Philadelphia. *Kalm.* ♃

*62. EUPHORBIA *adiantoides.*

E. ramoſiſſima, ramis alternis filiformibus dichotomis, foliis ovalibus et rhombeo-ſubrotundis, petiolis pedunculisque capillaceis. *Lamarck. encycl.* 2. *p.* 422.

Krullfarrenartige Wolfsmilch. W.

Habitat in Peru.

Planta *tenera facie Adianti Capilli Veneris.* Caulis *pedalis tenuis glaber, ramis alternis filiformibus et dichotomis.* Folia *glabra integerrima petiolo longo capillari ſuffulta, caulina alterna elliptica et majora; ramea oppoſita rhombeo-ſubrotunda ſetaceo-mucronata.* Pedunculi *ſolitarii uniflori terminales capillares in ultima ramorum dichotomia.* Calyces *turbinati ciliati.* Capſulae *globoſae trigonae pubeſcentes.* Lamarck. l. c.

63. EUPHORBIA *myrtifolia.*

E. dichotoma, foliis integerrimis ſubrotundis emarginatis ſubtus incanis, floribus ſolitariis, caule erecto. *Sp. pl.* 643. *Swartz obſ.* 197. *

Houttuyn Lin. Pfl. Syſt. 7. *p.* 56.

Myrtenblättrige Wolfsmilch. W.

Habitat ad Jamaicae *montibus frigidioribus.* ♄

*64. EUPHORBIA *imbricata.*

E. umbella dichotoma bifida, involucellis ſubrotundis mucronatis, foliis obovatis imbricatis ſerrulatis, caule fruticuloſo. *Vahl. ſymb.* 2. *p.* 54.

Dachziegelblättrige Wolfsmilch. W.

Habitat in Luſitania. ♄

Fruticulus *palmaris vel parum ultra baſi ramoſus, rami ſimplices teretes foliis tecti, denticulati, poſt caſum foliorum glabri uti tota planta.* Folia *ſeſſilia conferta, undique imbricata cuneata parva bilinearia mucronata: ſuperne denticulata, inferne integerrima.* Umbella *terminalis bis bifida.* Involucra *pentaphylla oblonga*

integerrima. Involucella *diphylla rotundiora.* Petala *integerrima.* Stamina *numeroſa:* Filamenta *albida.* Vahl l. c.

*65. EUPHORBIA *elliptica.*

E. ſubpubeſcens, foliis alternis ovato-ellipticis ſubdentatis petiolatis ſummis oppoſitis, umbella bifida. *Lamarck. encycl. 2. p. 422.**

Elliptiſche Wolfsmilch. W.

Habitat in Peru. ⊙

Caulis *ſesquipedalis et ultra, erectus ſimplex vel ramoſus ſtriatus.* Folia *ſesquipollicaria elliptica ſubdentata pubeſcentia, ſuperiora oppoſita.* Involucella *baſi rubro-maculata.* Umbella *bifida vel potius racemi bini approximati breves, interdum et tertius racemus vel ramus umbellae accedit, tunc rami ſunt parum alterni.* Pedunculi *breviſſimi.* Capſulae *glabrae.* Lamarck. l. c.

*66. EUPHORBIA *rubra.*

E. umbella bifida ſimplici terminali, foliis cuneiformi-obcordatis imbricatis. W.

E. (*rubra*) foliis cuneatis emarginatis imbricatis, umbellis bifidis, corollis pentapetalis. *Cavan. ic. 1. p. 21. t. 34. f. 1.**

Euphorbia exigua. γ. *Sp. pl.* 654.

Tithymalus exiguus ſaxatilis. *Bauh. pin.* 291. *prodr.* 132. *Magn. monſp.* 259. *t.* 258. *Burſ. XVI.* 55.

Rothe Wolfsmilch. W.

Habitat in collibus Hiſpania *et* Galliae *auſtralis.* ⊙

Tota planta rubeſcens. Petala *lunata rubra.* W.

*67. EUPHORBIA *herniariaefolia.* W.

E. umbella trifida ſimplici, foliis ſubrotundo-ovatis acutis pubeſcentibus imbricatis. W.

Tithymalus creticus ſupinus herniariae folio. *Tournef. cor. 1.*

Bruchkrautblättrige Wolfsmilch. W.

Habitat in Creta (v. ſ.)

Caulis *ſemipedalis et ultra ramoſus, ut videtur decumbens.* Rami *alterni teretes pubeſcentes.* Folia *conferta fere imbricata ſuboppoſita, vix manifeſta petiolata, parva Herniariae, ovata acuta utrinque pubeſcentia.* Umbella *ſeſſilis terminalis ſimpliciſſima trifida.* Involucrum *triphyllum foliolis forma foliorum caulinorum ſed majoribus.* Involucella *diphylla.* Capſulae *pubeſcentes.* W.

* * * * *Umbella trifida.*

68. EUPHORBIA *Peplus.*

E. umbella trifida: dichotoma, involucellis ovatis, foliis integerrimis obovatis petiolatis. *Amoen acad.* 3. *p.* 117. * *Gmel. ſib.* 2. *p.* 236. *Pollich. pal n.* 455. *Retz. obſ.* 3. *p.* 32. *Hoffm. germ.* 164. *Roth. germ. I.* 204. *II.* 524.

Tithymalus foliis rotundis, ſtipulis floralibus cordatis, petalis argute corniculatis. *Hall. helv. n.* 1049.

Euphorbia foliis obverſe ovatis integerrimis, umbella univerſali trifida triphylla, partialibus dichotomis, diphyllis. *Hort. cliff.* 199. *Fl. ſuec.* 437, 426. * *Roy. lugdb.* 197. *Dalib. pariſ* 156.

Peplus ſ. Eſula rotunda. *Bauh. pin.* 292. *Lob. ic.* 362.

Peplus. *Fuchſ. hiſt.* 603 *Dod. pempt.* 375. *Tabernaem.* 597.

β. Peplis minor. *Bauh. hiſt.* 3. *p.* 670.

Houttuyn Lin. Pfl. Syſt 7. *p.* 57.

Garten-Wolfsmilch. W.

Habitat in Europae *cultis oleraceis.* ⊙ (v. v.)

Petala *lunata.*

Varietas β. venit in hortis noſtris ſub nomine E. minimae, ſed mihi non diverſa videtur, licet duplo ſit minor. W.

69. EUPHORBIA *falcata.*

E. umbella trifida: dichotoma, involucellis ſubcordatis mucronatis, foliis lanceolatis obtuſiuſculis. *Amoen. acad.* 3. *p.* 118. *Pollich pal n.* 456. *Iacq. auſtr t.* 121. *Hoffm. germ.* 164 *Roth. germ. I.* 204. *II.* 524.

Tithymalus annuus ſupinus, folio rotundiore acuminato. *Tournef. inſt* 87.

Pithyuſa minor ſubrotundis et acutis foliis. *Barrel. ic.* 751.

Tithymalus Eſula exigua chalepenſis ramoſior, foliis latioribus lintriformibus. *Moriſ. hiſt.* 3. *p.* 339.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 58.

Zugeſpitzte Wolfsmilch. W.

Habitat in Europa *auſtrali.* ⊙ (v. ſ.)

70. EUPHORBIA *exigua.*

E. umbella trifida: dichotoma, involucellis lanceolatis, foliis linearibus. *Amoen. acad.* 3. *p.* 118. * *Pollich. pal.* 457. *Flor. dan. t.* 542. *Weig. obſ. bot. p.* 31. *Hoffm. germ.* 164. *Roth. germ. I.* 205. *II.* 525.

Tithymalus foliis linearibus, ſtipulis lanceolatis ariſtatis. *Hall. helv. n.* 1048.

α. Euphorbia *exigua, acuta.*

E. inermis, foliis alternis linearibus acutis, umbella univerſali trifida, partialibus dichotomis diphyllis. *Hort. cliff.* 199. *Hort. upſ.* 143.* *Roy. lugdb.* 197. *Dalib pariſ.* 157. *Gort. gelr.* 304.

Tithymalus ſ. Eſula exigua. *Bauh. pin.* 291.

Eſula minima Tragi. *Dalech. hiſt.* 1656.

β. Euphorbia *exigua, retuſa.*

E. (*retuſa*) foliis lineari-retuſis, involucellis lanceolatis. *Cavan. ic.* 1. *p.* 21. *t.* 34. *f.* 3*

Tithymalus ſ. Eſula exigua, foliis obtuſis. *Buh. pin.* 291. *prodr.* 132.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 59.

Kleine Wolfsmilch. W.

Habitat in Luſatia, Gallia, Helvetia, Hiſpania *inter ſegetes*: β. Monſpelii, Hiſpaniae in ſaxoſis. ⊙ (v. v.)

Petala *lunata.* Fructus laevis.

Umbella *ſaepe* 4-*et* 5-*fida.* Weig. l. c.

*71. EUPHORBIA *obliterata.*

E. umbella trifida, foliis oblongis trapezoideis ſerratis pubeſcentibus, altera baſeos parte obliteratis. *Swartz prodr.* 76. *Fl. ind. occid.* 2. *p.* 872. *

E. (*obliterata*) umbella trifida, foliis oblongis ſerratis, altera baſeos parte obliteratis. *Iacq. amer.* 151.

Schiefe Wolfsmilch. W.

Habitat in ſiccis arenoſis maritimis Jamaicae, Hiſpaniolae.

Planta caulibus *hirſutis lacteſcentibus*, foliis *apice attenuatis.* Iacq. l. c.

Flores *conferti terminales.* Umbella *trifida parva, floribus confertis minutis vireſcentibus.* Involucellum *nullum.* Germina *hirſuta.* Swartz l. c.

*72. EUPHORBIA *ſpathulata.*

E. umbella trifida dichotoma, foliis alternis oblongo-ſpathulatis ſubſerratis. *Lamarck. encycl.* 2. *p.* 424.

Spathelblättrige Wolfsmilch. W.

Habitat in Braſilia *in* Monte Video.

Planta *glabra, videtur herbacea, pedalis et ultra, ramis brevibus alternis inſtructa.* Folia *alterna ſeſſilia oblongo-ſpathulata ſubſerrata glabra vix uncialia.* Umbella *trifida bis vel ter dichotoma.* Involucrum *tri-*

phyllum, foliolis ovalibus obtuſis. Involucella *ut involucrum ſed minora et acuminata.* Capſulae *glabrae.* Lamarck. l. c.

*73. EUPHORBIA *micrantha.*

E. umbella trifida dichotoma, foliis lanceolato-obvatis ſerratis, involucellis cordatis, petalis integris, capſulis verrucoſis. W.

Euphorbia micrantha. *Stephan. in Litt.*

Feinblütige Wolfsmilch. W.

Habitat in Perſia *boreali.* ☉ (v. ſ.)

Caulis *teres ſimpliciſſimus pedalis glaber.* Folia *alterna obovato-lanceolata acuta glabra argute ſerrata.* Umbella *trifida dichotoma.* Involucrum *triphyllum ut folia caulina.* Involucella *cordata mucronata ſerrata.* Flores *parvi.* Petala *integra.* Capſula *punctis elevatis obſita.* W.

*74. EUPHORBIA *dracunculoides.*

E. umbella trifida dichotoma, foliis ſparſis linearibus longis integerrimis, involucris et involucellis foliis ſimilibus. *Lamarck. encycl.* 2. *p.* 424.

Langblättrige Wolfsmilch. W.

Habitat in inſula Mauritii. ☉ (v. ſ.)

Caulis *ſex-vel octouncialis ſimplex undique foliis obſitus.* Folia *ſparſa linearia integerrima glabra erecta bipollicaria et binas lineas lata, obtuſa.* Involucrum *et* involucella *foliis ſimilia ſed breviora et acutiora.* Flores *in dichotomia umbellae ſeſſiles.* Capſulae *glabrae.* Lamarck. l. c.

75. EUPHORBIA *tuberoſa.*

E. umbella trifida, involucro tetraphyllo, caule nudo, foliis oblongis emarginatis. *Amoen. acad.* 3. *p.* 117.

E. inermis, foliis oblongis obtuſis emarginatis. *Roy. lugdb.* 199.

Tithymalus humilis, folio lapathi. *Buxb. cent.* 2. *p.* 27. *t.* 23.

Tithymalus tuberoſus acaulos, foliis oblongis cucullatis et planis. *Burm afr.* 9. *t.* 4.

Tithymalus africanus humilis, foliis latioribus oblongis tuberoſa radice. *Raj. ſuppl.* 933.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 60.

Knollige Wolfsmilch. W.

Habitat in Aegypto *et* Aethiopia. ♄

* * * * * *Umbella quadrifida.*

76. EUPHORBIA *Lathyris.*

E. umbella quadrifida: dichotoma, foliis oppoſitis integerrimis. *Amoen. acad.* 3. *p.* 120.* *Mat. med.* 121. *Gmel. ſib.* 2. *p.* 230. *Mill. dict. n.* 18. *Hoffm. germ.* 165. *Roth. germ. I.* 205. *II.* 526.

Tithymalus foliis ellipticis, ſtipulis umbellaribus, quaternis, petalis bicornibus ſucciferis. *Hall. helv. n.* 1044.

E. inermis, foliis oppoſitis lanceolatis integerrimis, umbella univerſali quadrifida tetraphylla, ulterioribus dichotomis. *Hort. upſ.* 140.

E. inermis, foliis oppoſitis lanceolatis, umbella univerſali trifida polyphylla, partialibus triphyllis reliquis diphyllis. *Hort. cliff.* 198. *Roy. lugdb.* 196. *Dalib. pariſ.* 155.

Lathyris major. *Bauh. pin.* 293.

Lathyris. *Cam epit.* 968. *Fuchſ. hiſt.* 454. *Mathiol. p.* 1259. *Blackw. t.* 123.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 60.

Kreuzblättrige Wolfsmilch. W.

Habitat in Gallia, Italia, Helvetia, Germania, Carniolia *ad agrorum margines.* ♂ (v. v.)

77. EUPHORBIA *terracina.*

E. umbella quadrifida: dichotoma, foliis alternis lanceolatis retuſis mucronatis. *Sp. pl.* 654.

E. umbella quadrifida: bifida, foliolis cuneiformi-linearibus tridentatis. *Allion. corſ.* 209. *t.* 3.?

Tithymalus marinus, folio retuſo, terracinus. *Barr. ic.* 833.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 62.

Spaniſche Wolfsmilch. W.

Habitat in Hiſpania. *Allſtraemer.* ⊙

Caulis *herbaceus, ultra ſemipedalis, teres.* Folia *alterna, lanceolata, laevia, obliqua, margine ſcabriuſcula, apice retuſa et quaſi truncata mucrone reflexo.* Involucrum *ſubquadriphyllum: foliolis oblongo-ovatis, obtuſis, folio latioribus, vix ſerratis.* Involucella *ovata, baſi truncata.* Petala *flaveſcentia, bi-ſ. tridentata.* Fructus *glabri. Ex inferioribus alis foliorum ramuli ſteriles. Figura* Foliorum *ad Euph. exiguam retuſam prope accedit, ſed in illa lineam lata, huic digiti magnitudine.*

*78. EUPHORBIA *diffuſa.*

E. umbella quadrifida dichotoma, involucellis obtuſis, foliis alternis lineari-cuneiformibus emarginatis mucronatis, caule ramoſo diffuſo. W.

H. (*diffuſa*) umbella quadrifida vel quinquefida dichotoma, caule diffuſiſſimo, foliis cuneiformibus integerrimis alternis. *Jacq ic rar.* 1. *t.* 88. *Miſc.* 2. *p.* 311. * *Hoſt. ſynop.* 260. *

Vieläſtige Wolfsmilch. W.

Habitat in Auſtriae *arvis.* ⊙

Valde affinis E. exiguae β. retuſae, ſed excepto numero variante radiorum umbellae diſtincta involucellis obtuſis; a praecedente magnitudine, foliis glabris et habitu diverſa. W.

79. EUPHORBIA *Apios.*

E. umbella quadrifida: bifida, involucellis reniformibus: primis obcordatis. *Syſt. veg.* 375.

E. umbella quinquefida: bifida, involucellis obcordatis. *Amoen. acad.* 3. *p.* 120. *Sp. pl.* 2. *p* 656. *Mill. dict. n.* 26.

Tithymalus tuberoſa pyriformi radice. *Bauh. pin.* 292.

Tithymalus tuberoſa radice. *Cluſ. hiſt.* 2. *p.* 190.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 63.

Cretiſche Wolfsmilch. W.

Habitat in Creta. ♃

Rami *ſteriles*, Foliis *lineari-lanceolatis obtuſis:* Floriferi *vero ſubrotundo-ovatis.* Umbella 4-*fida*, 2-*fida, nec ultra.* Involucrum *univerſale* 4-*phyllum: foliolis ſubrotundis, vix acutis.* Involucella *priora foliolis obcordatis; poſteriora reniformibus, vix mucronatis.*

*80. EUPHORBIA *lacta.*

E. umbella quadri-vel quinquefida, bis dichotoma, involucellis primis oblongis, ſuperioribus rhombeo-ſubrotundis, foliis lineari-lanceolatis ſubemarginatis integerrimis. *Ait. Kew.* 2. *p.* 141.

E. (*divaricata*) umbella trifida vel quadrifida, caule fruticoſo dichotomo. *Jacq. ic. rar.* 1. *t.* 87. *Collect.* 1. *p.* 58 *

Seidelbaſtblättrige Wolfsmilch. W.

Habitat - - - - ♄ (v. ſ)

Tota planta glabra. Caulis *fruticoſus teres laevis.* Folia *ſparſa ſeſſilia ſesquiuncialia.* Involucra *univerſalia foliis ſimillima.* Involucella *primae dichotomiae*

ovali-oblonga ſubemarginata foliis dimidio breviora; involucella ſecundae dichotomiae et floralia elliptico-ſubrotunda emarginata. Ait. l. c.

* * * * * * *Umbella quinqueſida.*

81. EUPHORBIA *geniſtoides.*

E. umbella quinquefida bifida, involucellis ovatis, foliis linearibus erectis, caule frutescente. *Mant.* 564. *Berg. cap.* 146.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 64.

Ginſterartige Wolfsmilch. W.

Habitat ad Cap. b. ſpei ♄

Frutex *erectus: ramis alternis, ſtrictis, ſimpliciſſimis, brevioribus, etiam apice floriferis.* Folia *linearia, confertiuſcula, erecta, glabra, integerrima, parva (minora reliquis, nec majora E. exiguae.)* Umbellae *terminales, ſeſſiles.* Involucrum *teraphyllum: foliolis lanceolatis, longitudine Umbellae quinquefidae.* Involucella *diphylla, ovato-rhombea.* Umbellula *bifida, nec ultra.* Petala *lunata.* Capſulae *glabrae.*

82. EUPHORBIA *ſpinoſa.*

E. umbella ſubquinquefida ſimplici, involucellis ovatis: primariis triphyllis, foliis oblongis integerrimis, caule fruticoſo. *Amoen. acad.* 3. *p.* 120 *

E. inermis fruticoſa, foliis lanceolatis integerrimis, floribus ſolitariis terminalibus, involucris triphyllis. *Hort. cliff.* 201. *Roy. lugdb.* 196.

E. inermis fruticoſa, ramis ſiccis pungens, foliis ad umbellas ternis ſaepius ovatis, floribus ſolitariis. *Sauv. monſp.* 51.

Tithymalus raguſinus, flore luteo pentapetalo. *Herm. lugdb.* 600. *t.* 601.

Tithymalus maritimus ſpinoſus. *Bauh. pin.* 291.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 64.

Dornige Wolfsmilch. W.

Habitat in Cretae *et* Galloprovinciae *montibus maritimis.* ♄ (v. ſ.)

Flores *ſaepius ſolitarii.* Capſulae *verrucoſae.* Petala *rotunda.* Rami *ſeneſcentes ſicci et aridi perſiſtunt. unde quaſi ſpinoſa apparet.*

Raguſinam *Hermanni 5-petalam, a ſpinoſa 4-petala diſtinguit Moriſonus; mihi altera tantum viſa.*

83. EUPHORBIA *epithymoides.*

E. umbella quinquefida: bifida, involucellis ovatis denticulatis, foliis integerrimis lanceolatis obtuſis retuſis ſubtus villoſis, capſulis papilloſo-hiſpidis. W.

E. umbella quinquefida: bifida, involucellis ovatis, foliis lanceolatis obtuſis ſubtus villoſis. *Sp. pl.* 656. *Iacq. auſtr. t.* 344.

Tithymalus epithymi fructu. *Col. ecphr.* 2. *p.* 52. *t.* 51.

Peplios altera ſpecies. *Bauh. pin.* 292.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 65.

Gelbblättrige Wolfsmilch. W.

Habitat in Italia, Auſtria. ♃ (v. ſ.)

Accedit ad E. dulcem, ſed Folia magis margine ſcabra et ſubtus villoſa. Fructus hiſpidi, ſetis ſubulatis purpuraſcentibus adſperſi.

*84. EUPHORBIA *villoſa.*

E. umbella quinquefida: ſubtrifida, involucellis ovatis acutis ſubdenticulatis, foliis lanceolatis acutiuſculis margine ſcaberrimis ſubtus villoſis, capſulis glabris verrucoſis. W.

Euphorbia villoſa. *Waldſtein et Kitaibel pl. rar. hung.*

Vielhaarige Wolfsmilch. W.

Habitat in pratis humidis Hungariae. ♃ (v. ſ.)

Maxime affinis praecedenti, convenit enim habitu, *petalis integerrimis foliisque ſeſſilibus integerrimis, at differt:* foliis *acutiuſculis margine ſcaberrimis, radiis umbellae ſubtrifidis* involucellis *acutis, et* fructu *verrucoſo nec papillis elongatis obſito.* W.

Ab E. amygdaloide pro qua vulgo habetur abunde diverſa, involucellis *omnino diſtinctis nec baſi connatis.* Involucella *primae diviſionis umbellae ſunt terna, ultimae vero bina.* W.

85. EUPHORBIA *dulcis.*

E. umbella quinquefida: bifida, involucellis ſubovatis denticulatis, foliis lanceolatis obtuſis apice denticulatis, capſulis verrucoſis piloſis. W.

E. umbella quinquefida: bifida, involucellis ſubovatis, foliis lanceolatis obtuſis integerrimis. *Amoen. acad.* 3. *p.* 122. *Iacq. auſtr. t.* 213. *Kniph. cent.* 12. *n.* 42. *Hoffm. germ.* 165. *Roth. germ. I.* 205. *II.* 527.

E. (*dulcis*) umbella quinquefida: bifida, involucellis cordatis denticulatis, foliis lanceolatis ſubintegerrimis. *Vahl. ſymb.* 2. *p.* 55?

Tithymalus foliis ellipticis ſerratis, ſtipulis umbellaribus quinis, floralibus fructu aſperrimo. *Hall. helv. n.* 1051. R.
Tithymalus montanus non acris. *Bauh. pin.* 292.
Tithymalus dulcis tuberoſa radice. *Barr. ic.* 909.
Pithyuſa ſ. Eſula minor altera, floribus rubris. *Lob. ic.* 358.
Eſula foliſequa. *Riu. t.* 117. R.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 65.
Süſſe Wolfsmilch. W.
Habitat in Germaniae, Helvetiae, Galliae, Italiae *umbroſis* ♃ (v. v.)

Involucella *ſubtiliſſime ſerrata.* Fructus *ruber, muricatus.* Petala *integra.* Variat *foliis villoſi.* Exſiccatione *nigra evadit.*

Primo vere ante evolutionem umbellae nigreſcit, nec poſtea: Folia *apice denticulata ſubtus pilis raris obſita.* Capſulae *juniores piloſae, adultae verrucoſae interſperſis pilis tectae.* Petala *ut in omnibus affinibus ſunt integra, nec bicornia, ut Clariſſ. Vahl autumat.* W.

*86. EUPHORBIA *ambigua.*

E. umbella quinquefida: bifida, involucellis ovatis obtuſis integerrimis, foliis lanceolatis obtuſis integerrimis, capſulis glabris verrucoſis. W.
Euphorbia ambigua. *Waldſtein et Kitaibel pl. rar. hung.*
Zweifelhafte Wolfsmilch. W.
Habitat in Hungaria. (v. ſ.)

Valde affinis praecedentis ſed differre vedetur: foliis *glabris integerrimis,* involucellis *non denticulatis,* capſulis *glabris.* W.

*87. EUPHORBIA *carniolica.*

E. umbella quinquefida: bifida, radiis nutantibus involucellis ovatis integerrimis, foliis lanceolatis acutis integerrimis, capſulis glabris verrucoſis. W.
E. (*carniolica*) umbella quinquefida, radiis nutantibus, involucris, involucellis et foliis lanceolatis. *Jacq. auſt. t.* 14. *Hoſt. ſynop.* 262.*
Tithymalus piloſus. *Scop. carn. n.* 576. *t.* 21.
Kraienſche Wolfsmilch. W.
Habitat in Carnioliae *ſylvaticis alpinis.* ♃ (v. ſ.)

Ab affinibus ſatis diſtincta radiis *umbellae nutantibus.* Folia *ſubtus ut in E. dulci pilis raris tecta.* W.

*88. EUPHORBIA *angulata.*

E. umbella quinquefida: bifida, involucellis rhombeo-ovatis denticulatis, foliis lanceolatis obtuſis denticulatis, capſulis glabris verrucoſis, caule ſulcato. W.

E. (*angulata*) umbella quinquefida: bifida, involucellis ovatis, caule ſtriato-angulato. *Iacq. ic. rar. t.* 481. *Collect.* 2. *p.* 309. *

Geſtreifte Wolfsmilch. W.

Habitat in Auſtria. ♃ (v. ſ.)

Nimis affinis E. ambiguae, quae et caule angulato gaudet, ſed haec differt: involucellis rhombeo-ovatis denticulatis ut et foliis denticulatis. Folia *ſunt ſubtus pilis ſparſis obſita.* W.

Conveniunt inter ſe ſpecies 82. 83. 84. 85. 86. 87. *ut vix limites obſerventur.* W.

89. EUPHORBIA *Pithyuſa.*

E. umbella quinquefida: bifida, involucellis ovatis mucronatis, foliis lanceolatis: infimis involutis retrorſum imbricatis. *Amoen. acad.* 3. *p.* 122.

Tithymalus Pithyuſa. *Scop. carn. n.* 575.

Tithymalus foliis brevibus aculeatis. *Bauh. pin.* 292.

Pithyuſa. *Dalech. hiſt.* 1652.

Tithymalus maritimus, juniperi folio. *Bocc. ſic.* 9. *t.* 5. *Moriſ. hiſt.* 3. *p.* 337. *ſ.* 10. *t.* 1. *f.* 25.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 66.

Wachholderartige Wolfsmilch. W.

Habitat in arenoſis Hiſpaniae, Italiae, Carnioliae, Maſſiliae. ♃ (v. ſ.)

90. EUPHORBIA *portlandica.*

E. umbella quinquefida: dichotoma, involucellis ſubcordatis concavis, foliis lineari-lanceolatis acutis glabris patentibus. *Sp. pl.* 656. *Hudſ. angl.* 183.

Tithymalus montanus, elulae, folio minor italicus. *Barr. ic.* 822.

Tithymalus maritimus minor. *Raj. ſyn.* 3. *p.* 313. *t.* 24. *f.* 6.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 67.

Engliſche Wolfsmilch. W.

Habitat in Angliae *comitatu Devonſhire.* ♄ (v. ſ.)

Caules *fruticuloſi, palmares, glabri, teretes, hyeme rubri.* Folia *alterna, ſubſeſſilia, lineari-lanceolata, mucronata, glabra, patentia, baſi ſubtus rubra.* Ramuli *laterales: ex alis inferioribus ſteriles, demum*

excreſcentes et caulem inumbrantes. Umbellae *terminales, quinquefidae, dichotomae, patentes.* Involucrum *foliis ſimile.* Involucella *ſubcordata, concava, mucronata.* Flores *ſubſeſſiles, lutei: Primi et ſecundi maſculi: Petalis muticis, obtuſiſſimis; Reliqui hermaphroditi: Petalis bicornibus.* Fructus *glabri angulis muricatis.*

*91. EUPHORBIA *ſaxatilis.*

E. umbella quinquefida: bifida, involucris cordatis, involucellis reniformibus mucronatis, foliis ſpathulato-linearibus emarginatis aveniis. W.

E. (*ſaxatilis*) umbella quinquefida: bifida, involucris et involucellis cordatis, foliis oblongis glabris. *Murr. ſyſt. veg.* 452. *Iacq. auſt. t.* 345. *Hoſt. ſynop.* 262.

Stein-Wolfsmilch. W.

Habitat in ſaxoſis Auſtriae. ♃ (v. ſ.)

Caules *digitales adſcendentes ſimplices plures e radice.* Folia *glauca avenia emarginata glabra, inferiora oblonga vel obovata.* Petala *obtuſe lunata.* Capſulae *glabrae.* W.

92. EUPHORBIA *Paralias.*

E. umbella ſubquinquefida: bifida, involucellis cordato-reniformibus, foliis ſurſum imbricatis. *Amoen. acad.* 3. *p.* 129. *Hudſ. angl.* 183. *Iacq. hort. t.* 188. *Hoffm. germ.* 165. *Roth. I.* 205. *II.* 527. *Ehrh. Beitr.* 2. *p.* 128. *.

E. inermis, foliis ſetaceo-linearibus confertis, umbella univerſali multifida, partialibus ramoſe bifidis. *Hort. cliff.* 200. *Roy. lugdb.* 193. *Gort. gelr.* 306. *Gron. orient.* 162.

Tithymalus floribus linearibus ariſtatis imbricatis, ſtipulis umbellaribus ovato-lanceolatis, floralibus cordatis. *Hall. helv. n.* 1055.

Euphorbia umbella multifida: dichotoma, involucellis cordatis, foliis linearibus, capſulis glabris. *Scop. carn. ed.* 1. *p.* 435. *n.* 11. *ed.* 2. *n.* 581.

Tithymalus maritimus. *Bauh. pin.* 291. *Dod. pempt.* 370. *f.* 1. 2.

Tithymalus paralias. *Cam epit.* 962.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 68.

Dichtblättrige Wolfsmilch. W.

Habitat in Europae *arena maritima.* ♃ (v. ſ.)

Petala *integra.* Planta *glauca.*

*93. EUPHORBIA *juncea.*

E. umbella quinquefida: dichotoma, foliis involucrisque lineari-lanceolatis acutis, involucellis ovato-oblongis acuminatis. *Ait. Kew.* 2. *p.* 142. *Jacq. hort. ſchoenb.* 1. *p.* 56. *t.* 107. *

Schlanke Wolfsmilch. W.

Habitat in inſula Porto Santo *prope* Maderam. ♃ (v. v.)

Folia *anguſta lineari-lanceolata, in ramis vero ſterilibus anguſtiſſime Pini.* Involucella *oblongo-ovata acuminata.* Flores *parvi.* Petala *ſubrotunda.* Styli *bifidi.* Stigmata *capitata.* Capſulae *glabrae.* W.

94. EUPHORBIA *aleppica.*

E. umbella quinquefida: dichotoma, involucellis ovato-lanceolatis mucronatis, foliis inferioribus ſetaceis. *Amoen. acad.* 3. *p.* 122. *Mill. dict. n.* 27.

Tithymalus foliis inferioribns capillaceis: ſuperioribus myrto ſimilibus. *Moriſ. hiſt.* 3 *p.* 338.

Tithymalus cypariſſius. *Alp. exot.* 65. *t.* 64.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 68.

Aleppiſche Wolfsmilch. W.

Habitat in Creta, Aleppo. ♃ (v. ſ.)

Folia *inferiora confertiſſima lineari-lanceolata reflexa, ſuperiora lanceolata acuta patentia.* Involucra *et* Involucella *ovato-lanceolata mucronata.* Petala *obtuſe lunata.* Styli *ſimplices.* Stigmata *capitata.* Capſulae *glabrae.* W.

95. EUPHORBIA *pinea.*

E. umbella quinquefida: dichotoma, involucellis cordatis, foliis linearibus acuminatis confertis, capſulis laeviuſculis. *Syſt. veg.* 376.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 69.

Fichtenblättrige Wolfsmilch. W.

Habitat

96. EUPHORBIA *ſegetalis.*

E, umbella quinquefida: dichotoma, involucellis cordatis acutis, foliis lineari-lanceolatis: ſuperioribus latioribus. *Ger. prov.* 538. *Pall. it.* 1. *p.* 145. *Hoffm. germ.* 165. *Roth. germ. I.* 205. *II.* 528.

E. umbella quinquefida: dichotoma, involucellis cordatis acutis, foliis lineari-lanceolatis, ramis floriferis *Amoen. acad.* 3. *p.* 121. * *Hudſ angl.* 182. *Sp. plant.* 2. *p* 657.

E. inermis, foliis alternis linearibus acutis, partialibus umbellae ovato-rhombeis, petalis bicornibus. *Hort. upſ.* 142. *

E. inermis, foliis linearibus acutis ad umbellam quinis iſoſcelibus, ad umbellulas ter dichotomas ovato-trigonis. *Sauv. monſp.* 46.

Tithymalus annuus, lunato flore, linariae folio longiore. *Moriſ. hiſt.* 3. *p.* 339. *ſ.* 10. *t.* 2. *f.* 3.

Tithymalus ſegetum longifolius. *Raj. angl.* 3. *p.* 312.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 69.

Acker-Wolfsmilch. W.

Habitat in Mauritania, Ruſſia, Anglia, Gallia, Germania. ⊙ (v. v.)

Petala *lunata.* Fructus *laevis angulis ſcabris, umbella ſaepe longitudine totius plantae.* Involucella *fere ſemiorbiculata.* Rami *etiam floriferi infra umbellam.*

*97. EUPHORBIA *provincialis.* W.

E. umbella quinquefida: bifida, involucellis cordatis mucronatis, foliis oblongis. W.

Euphorbia ſeticornis. *Poiret. it. ed. germ. p.* 195.?

Provencer Wolfsmilch. W.

Habitat in Gallo-provinciae *collibus* ⊙ (v. v.)

Saepius hujus ſemina ſub nomine E ſegetalis accepi, ſed pluribus notis diverſa eſt, nec cultura mutatur. Caulis *ſimplex pedalis et ultra, ramis aliquot ſterilibus in foliorum axillis ſuperioribus inſtructus.* Folia *ſeſſilia oblonga glabra integerrima obtuſa ſubmucronata.* Umbella *quadri-vel quinquefida tantum bifida nec dichotoma.* Involucrum *pentaphyllum, foliis ovatis acutis.* Involucella *cordata mucronata, lobis baſi ſaepius mucrone donatis.* Petala *roſtrato-lunata.* Styli *bifidi.* Capſulae *omnino laeves, nec angulis ſcabrae.* W.

98. EUPHORBIA *helioſcopia.*

E. umbella quinquefida: bifida: dichotoma, involucellis obovatis, foliis cuneiformibus ſerratis glabris, capſulis laevibus. *Vahl. ſymb.* 2. *p.* 55.

E. umbella quinquefida: trifida: dichotoma, involucellis obovatis, foliis cuneiformibus ſerratis. *Amoen. acad.* 3. *p.* 1?4.* *Pollich pal. n.* 458. *Kniph. cent.* 12. *n.* 43. *Flor. dan. t.* 725. *Roth. germ. I.* 206. *II.* 520. *Hoff. germ.* 166.

E. foliis crenatis, umbella univerſali quinquefida pentaphylla, partialibus trifidis, propriis triphyllis. *Hort. cliff.* 198. *Fl. ſuec.* 436. 425. *Roy. lugdb.* 197. *Dalib. pariſ.* 156.

E. inermis, foliis ſubrotundis crenatis. *Fl. lapp.* 220.

E. umbella quinquefida, trifida: dichotoma, involucellis

ovatis, petalis ſubrotundis, capſulis glabris. *Scop. carn. ed.* 1. *p.* 434. *n.* 9. *ed.* 2. *n.* 579 *ſub* Tithymalo.
Tithymalus foliis petiolatis, ſubrotundis, ſerratis, ſtipulis rotundis ſerratis. *Hall. helv. n* 1050.
Tithymalus helioſcopius. *Bauh pin.* 291. *Fuchſ. hiſt.* 811. *Camer. epit.* 963 *Dodon purg.* 145.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 70.
Sonnenzeiger-Wolfsmilch. W.
Habitat in Europae *cultis.* ⊙ (v. v.)
Umbellae radii bis trifidi, dichotomi et involucra foliis ſimila. Petala *integra.* Fructus *laevis.*

99. EUPHORBIA *pubeſcens.*
E. umbella quinquefida: trichotoma, involucellis ſemicordatis, foliis cuneiformibus piloſis ſerrulatis, capſulis muricatis. *Vahl ſymb.* 2. *p.* 55. *
Tunetaniſche Wolfsmilch. W.
Habitat in cultis Tuneti. (v. ſ.)
Primo intuitu ita ſimilis E. helioſcopiae, ut eandem diceres, folia *enim et modus creſcendi omnino idem; differt vero:* caule *ſuperne,* foliis, pedunculis *et* pedicellis *villoſis,* capſulisque *e tuberculis parvis muricatis.* Vahl. l. c.

100. EUPHORBIA *ſerrata.*
E. umbella quinquefida: trifida: dichotoma, involucellis diphyllis reniformibus, foliis amplexicaulibus cordatis ſerratis. *Amoen. acad.* 3. *p.* 125. * *Vahl. ſymb.* 2. *p.* 55. * *Jacq. ic.* 3. *t.* 384. *Collect.* 2. *p* 310. *
E. inermis foliis denticulatis caulinis lanceolatis, umbellarum cordatis. *Hort. cliff.* 200. *Hort. upſ.* 141. * *Roy lugdb.* 197.
Tithymalus characias, folio ſerrato *Bauh. pin.* 290.
Tithymalus myrtites valentinus. *Cluſ. hiſt.* 2. *p* 189.
β. E. (*retuſa*) umbella trifida, dichotoma, involucris lanceolatis, foliis linearibus retuſis denticulatis, inferioribus minoribus *Forſk. deſcr.* 93. *Icon. t.* 13.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 71.
Geſägte Wolfsmilch. W.
Habitat in Narbona, Hiſpania, Italia, Oriente. ⊙ (v. ſ.)

101. EUPHORBIA *verrucoſa.*
E. umbella quinquefida: ſubtrifida: bifida, involucellis ovatis, foliis lanceolatis ſerrulatis villoſis, capſulis verrucoſis. *Amoen. acad.* 3. *p.* 120. * *Hudſ. angl.* 183. *Hoffm. germ.* 166. *Roth. germ. I.* 206. *II.* 630.

Tithymalus foliis ellipticis ſerratis, ſtipulis umbellaribus quinis, floralibus obtuſe quadrangulis, fructibus undique exaſperatis. *Hall. helv. n.* 1052.

E. inermis, foliis linearibus, umbella univerſali trifida, triphylla, partialibus trifidis, propriis dichotomis diphyllis, foliolis ſubrotundis. *Guett. ſtamp.* 269. *Dalib. pariſ.* 154.

Tithymalus myrſinites, fructu verrucae ſimili. *Bauh. pin.* 291. *Moriſ. hiſt. ſect.* 10. *t.* 3. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 72.

Warzige Wolfsmilch. W.

Habitat in Gallia, Helvetia, Italia, Oriente. ♂

Fructus verrucoſus ſimulque villoſus. Petala *integra.* Caules *decumbentes.*

*102. EUPHORBIA *glauca.*

E. umbella ſubquinquefida: trifida, involucris et involucellis ovatis, foliis ſparſis oblongo-lanceolatis integerrimis, caule fruteſcente. *Forſt. prodr. n.* 208.

Meergrüne Wolfsmilch. W.

Habitat in Nova Zeelandia. ♄

*103. EUPHORBIA *punicea.*

E. umbella quinquefida; trifida, involucellis ovalibus acuminatis coloratis, capſulis glabris, foliis obovato-lanceolatis ſubtus glaucis. *Swartz prodr.* 76. *Fl. ind. occid.* 2. *p* 873. * *Ait.* 2. *p.* 143. *Smith. ic. pictae.* 3. *tab.* 3.* *Jacq. ic. rar.* 3. *t.* 484. *Collect.* 2. *p.* 179. *

Schoengefärbte Wolfsmilch. W.

Habitat in petroſis montoſis Jamaicae *interioris.* ♄ (v. ſ.)

104. EUPHORBIA *corollata.*

E. umbella quinquefida: trifida: dichotoma, involucellis foliisque oblongis obtuſis, petalis membranaceis. *Amoen. acad.* 3. *p.* 122.

E. inermis, foliis lanceolatis obtuſis alternis, ramis floriferis dichotomis, petalis maximis ſubrotundis. *Gron. virg.* 58.

Tithymalus marianus, anguſtis rigidis foliis, ſummo caule ramoſus, foliis atque ramulis ad divaricationes ternis, *Pluk. mant.* 182. *t.* 446. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 73.

Groſ-blütige Wolfsmilch. W.

Habitat in Virginia, Canada.

105. EUPHORBIA *corallioides.*

E. umbella quinquefida: trifida: dichotoma, involucel-

lis ovatis, foliis lanceolatis, capſulis lanatis. *Amoen. acad.* 3. *p.* 123. *
E. inermis fruticoſa, foliis lanceolatis, involucro univerſali quinquefido, partialibus trifidis, reliquis bifidis. *Hort. upſ.* 142. * *Roy. lugdb.* 198.
Tithymalus arboreus, caule corallino, folio hyperici, pericarpio barbato. *Boerh. lugdb.* 1. *p.* 256.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 73.
Rothſtielige Wolfsmilch. W.
Habitat in Sicilia, Mauritania, Oriente. ♃

106. EUPHORBIA *piloſa.*

E. umbella quinquefida: trifida: bifida, involucris ovatis, petalis integris, foliis lateolatis ſupiloſis apice ſerrulatis. *Sp. pl.* 659.
E. foliis alternis ex ovali-lanceolatis, umbellis diphyllis ſubtrifloris, capſulis erectis muricatis, caule ſimplici. *Gmel. ſib.* 2. *p.* 226. *t.* 93.
Tithymalus paluſtris villoſus mollior erectus. *Bar. rar.* 41. *t.* 885.
Tithymalus incanus hirſutus. *Bauh. pin.* 262? *prodr.* 135?
Tithymalus Characias pratenſis incanus. *Magn. monſp.* 255.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 74.
Rauhfrüchtige Wolfsmilch. W.
Habitat in Sibiria. ♃ (v. v.)

Habitus exacte E. paluſtris, ut facile pro eadem ſumeretur, eodemque tempore floret, paulo tamen major. Folia *lato-lanceolata, alterna, utrinque vix manifeſte piloſa, apice ita tenuiſſime ſerrata, ut vix obſerventur ſerraturae.* Umbellae *cum umbellulis lateralibus ita coacervatae, ut primaria difficilius eruatur; luteae petalis et involucris.* Flores *primarii maſculi pentapetali; reliqui hermaphroditi tetrapetali: petalis transverſe ovalibus.* Fructus *verrucoſi et pilis albis ſubtiliſſimis adſperſi.* Rami *ſteriles ex alis foliorum inferiorum, ut ex ſummis alis pedunculi umbelluliferi.*

*107. EUPHORBIA *orientalis.*

E. umbella quinquefida: quadrifida: dichotoma, involucellis ſubrotundis acutis, foliis lanceolatis. *Amoen. acad.* 3. *p.* 123. *Mill. dict. n.* 23.
E. inermis, foliis lanceolatis, involucro univeſali quin-

quefido lanceolato, partiali tetraphyllo ſubrotundo, propriis diphyllis. *Roy. lugdb.* 198. •

Tithymalus orientalis, ſalicis folio, caule purpureo, flore magno. *Tourneſ. cor.* 2.

Houttuyn Lin. Pfl. Syſt 7. *p.* 75.

Orientaliſche Wolfsmilch. W.

Habitat in Oriente.

*108. EUPHORBIA *ſquamoſa.* W.

E. umbella quinquefida, ſubtrifida: bifida, involucellis ſubrotundis mucronatis, foliis oblongis acutis ſubrepandis ſubtus pubeſcentibus, capſulis ſquamoſis. W.

Tithymalus orientalis latiſſimo folio, villoſus, flore lunato. *Tourneſ. cor.* 2.

Schuppige Wolfsmilch. W.

Habitat in Cappadocia. (v. ſ.)

Caulis *tenuiſſime pubeſcens laevis.* Folia *breviſſime petiolata oblongo-ovata acuta, margine ſubrepanda, ſubtus pubeſcentia.* Umbella *ampla divaricata quinquefida, radiis bi-ſ. trifidis dichotomis.* Involucri *folia iis caulis ſimilia ſed minora.* Involucella *primaria ſubrotundo-ovata acuta, ſecundae diviſionis vero ſubrotunda mucronata.* Petala *lunata.* Capſulae *ſquamis deciduis tectae, glabrae.* W.

109. EUPHORBIA *platyphyllos.*

E. umbella quinquefida: trifida: dichotoma, involucris carina piloſis, foliis ſerratis lanceolatis, capſulis verrucoſis. *Syſt. nat.* 10. *p.* 1049 *n* 42. 47. *Iacq. auſtr. t.* 376. *Schreb. ſpicil. p* 8. *Pollich. pal. n.* 459. *Kniph. cent.* 9. *n.* 38. *Hoffm germ.* 166. *Roth. germ. I.* 206. *II* 531.

E. umbella quinquefida: ſubquadrifida: dichotoma, involucellis primariis tetraphyllis, foliis ſerratis lanceolatis ſeſſilibus. *Amoen. acad.* 3. *p.* 124.

E. inermis foliis alternis lanceolatis amplexicaulibus ſubſerratis, umbella univerſali quinquefida pentaphylla, partialibus bifidis. *Hort. upſ.* 141. *

E. inermis, foliis lanceolatis, umbella univerſali partialiumque prima quinquefidis: ſecunda trifida: reliquis bifidis. *Roy. lugdb* 198. *Dalib. pariſ.* 156.

Tithymalus foliis lanceolatis, ſerratis, ſtipulis floralibus cordatis, fructibus aſperis, lineis laevibus diviſis. *Hall. helv. n.* 1053.

Tithymalus arvenſis latifolius germanicus. *Bauh. pin.* 291. *Moriſ. hiſt. ſect.* 10. *t.* 3. *f.* 1.

Tithymalus platyphyllos. *Fuchſ hiſt.* 813. *Cauſ. hiſt.* 3. *p.* 670.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 76.
Flachblättrige Wolf milch. W.
Habitat in agris Galliae, Angliae, Germaniae. ⊙ (v. v.)

Caulis *erectus, pedalis, glaber.* Folia *alterna, remota, patentia, lanceolata, glabra, ſerrulata, carina villis rariſſimis: inferiora ſubpetiolata, verſus apicem latiora; ſuperiora baſi emarginata et ſupra baſin ad latera deflexa.* Involucrum *lanceolatum.* Involucella *partialia ovato-oblonga; reliqua ovato-cordata, omnia ſerrulata.* Petala *integra, ſuborbiculata, lutea.* Germina *obſcure verrucoſa. Ex alis foliorum umbellula trifida. Planta ſuperior viridi-flaveſcens.*

Tenellae *adhuc plantae foliis macula rubra e lineolis inſidet.*

*110. EUPHORBIA *literata*

E. umbella quinquefida: trifida: dichotoma, involucellis lanceolatis, foliis lanceolatis denticulatis pubeſcentibus, capſulis glabris verrucoſis. W.

E. (*literata*) umbella quinquefida: trifida: bifida, involucellis ovatis, foliis lanceolatis ſubpiloſis obſolete ſerrulatis, petalis integris, radice annua. *Iacq. ic. rar.* 3. *t.* 482. *Collect* 2 *p* 340. *

Rothſtreifige Wolfsmilch. W.
Habitat - - - - ⊙ (v. v.)

Folia *junioris plantae lineis rubris picta. Pro varietate praecedentis antea habui, ſed diverſa videtur,* foliis *anguſtioribus pubeſcentibus,* habituque *teneriore. Ab E. piloſa diſtinctiſſima,* radice *annua,* foliis *anguſtioribus maculatis,* fructibusque *non piloſis.* W.

* * * * * * * *Umbella multifida.*

111. EUPHORBIA *Eſula.*

E. umbella multifida: bifida, involucellis ſubcordatis, petalis ſubbicornibus, ramis ſterilibus, foliis uniformibus. *Amoen. acad* 3. *p.* 127. *Pollich. pal n.* 460. *Hoffm. germ.* 166. *Roth germ I.* 206. *II.* 532.

Tithymalus foliis linearibus, petalis emarginatis. *Hall. helv. n.* 1046.

E. inermis, foliis lanceolato-linearibus involucri univerſalis foliis quinis ovato-acutis, partialis ſemiorbiculatis. *Hort. upſ.* 141. *Sauv. monſp.* 47.

E. inermis, foliis ligulatis ad umbellam quinis ovato-oblongis acutis, bracteis trigonis. *Sauv. monſp.* 51.?
Tithymalus foliis pini, forte Doſcoridis Pithyuſa. *Bauh. pin.* 292.
Eſula *Riu. tetr.* 227.
Eſula minor *Dalech. hiſt.* 1653. *Dod. pempt.* 374. *Blackw t.* 163. *f.* 1 2.
Houttuyn Lin. Pfl Syſt. 7. *p.* 77.
Gemeine Wolfsmilch. W.
Habitat in Germania, Belgio, Gallio, Helvetia, Carniolia. ♃ (v. v.)

Flores *omnes fertiles.* Petala *flaveſcentia*, 4, *obſolete bicornia.* Fructus *glaber.* Umbellae *caulis laterales dichotomae.*

*112. EUPHORBIA *Gerardiana.*

E. umbella multifida dichotoma, involucellis ſubrotundis, petalis integerrimis ramis nullis. *Murr. ſyſt.* 454. *Jacq. auſt.* 5. *t.* 436. *Hoſt ſynop.* 265. *Hoff. germ.* 167.
E. (*Cajogala*) radice perenni, caule herbaceo, foliis alternis lineari-lanceolatis acutis mucronatis trinerviis, umbella multifida, involucri foliolis lato-lanceolatis, umbellulis dichotomis, involucellis diphyllis, foliolis lato-cordiformibus, petalis integris, fructu glabro. *Ehrh. Beitr.* 2. *p.* 102* *Hoffm. germ.* 167.
E. umbella multifida: bifida, involucellis triangulari-cordatis, foliis ſuperioribus latioribus. *Ger. proo.* 540.
Tithymalus amygdaloides anguſtifolius. *Tabern. ic.* 541. *Vaill. pariſ.* 192.
Tithymalus maritimo affinis, lineariae folio. *Bauh. pin.* 291.
Gerardiſche Wolfsmilch. W.
Habitat ad fluviorum et lacuum ripas Galliae, Hannoverae, Mansfeldiae, Bohemiae, Auſtriae. ♃ (v.v.)

113. EUPHORBIA *Cypariſſias.*

E. umbella multifida: dichotoma, involucellis ſubcordatis, ramis ſterilibus, foliis ſetaceis, caulinis lanceotis. *Amoen. acad.* 3. *p.* 127. *Weig. obſ. bot. p.* 31. *Hoff. germ.* 167. *Roth. germ. I.* 207. *II.* 532. *Pollich. pal. n.* 461. *Kniph. cent.* 12. *n.* 41. *Blackw. t.* 163. *f.* 3.
E. inermis, folius confertis linearibus, umbella univerſali multifida, partialibus dichotomis: foliolis ſubro-

tundis. *Hort. cliff.* 199.* *Hort. upſ.* 142.* *Roy. lugdb.* 197. *Dalib. pariſ.* 155 *Gort. gelr.* 105.

Tithymalus foliis linearibus, ſtipulis numeroſis, floralibus cordiformibus, petalis lunatis. *Hall. helv. n.* 1047.

Tithymalus cypariſſius. *Bauh. pin.* 291. *Dalech. hiſt.* 1644.

Tithymalus cypreſſinus. *Tabern. hiſt.* 990.

β. Eſula degener. *Riu. Hoppe ectyp.* 560. *Hoffm. germ.* 168.

Tithymalus cypariſſias, foliis punctis croceis notatis. *C. B. pin.* 291.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 79.

Cypreſſen-Wolfsmilch. W.

Habitat in Germaniae, Bohemiae, Helvetiae, Gall. Narbonenſis *collibus, viis, ſiccis.* ♃ (v. v.)

Primo *vere umbella ſimilis E. Eſulae; at umbella marceſcente ramuli laterales foliis anguſtiſſimis filiformi-planis excreſcunt Pinum referentes.* Petala *lunata.* Fructus *loevis.* (*Multi capſulas verrucoſas dicunt, quas et ego in meis video.* R.)

*114. EUPHORBIA *nicaeenſis.*

E. umbella quinquefida: bifida, involucellis cordato-ſubrotundis integerrimis, foliis lanceolatis mucronatis ſubcoriaceis. *Vahl. ſymb.* 3. *p.* 63.*

E. umbella multifida; dichotoma, involucellis ſubrotundis cordatis mucronatis, foliis cuneiformibus integerrimis glabris. *Hoſt ſynop.* 267.

E. umbella quinquefida ad decemfidam: dichotoma, involucellis cordato-ſubrotundis, foliis lanceolato-ſpathulatis glaucis. *Iacq. ic. rar.* 3. *t.* 485. *Collect.* 2. *p.* 324.*

E. glabra, caule ſubſimplici, umbella quinquefida: bifida, foliis ovatis, capſulis laevibus non ſulcatis, *Allion. ped. n.* 1039 *tab.* 69. *f.* 1.*

Nicaeenſiſche Wolfsmilch. W.

Habitat in comitatu Nicaeenſi, *in* Hiſpania *circa* Pampelonam, *inque* Auſtria *calidiore.* ♃ (v. ſ.)

Umbella *quinquefida, ſaepius vero multifida.* W.

115. EUPHORBIA *myrſinites.*

E. umbella ſuboctifida: bifida, involucellis ſubovatis, foliis ſpathulatis patentibus carnoſis mucronatis margine ſcabris. *Amoen. acad.* 3. *p.* 128.* *Mill. dict. n.* 19.

E. inermis, foliis ſuperioribus reflexis latioribus lanceolatis, umbella univerſali trifida, partialibus bifidis. *Hort. cliff.* 199. *Hort. upſ.* 141.
E. inermis, foliis ligulatis ſpinula terminatis, ad umbellam duodenis, bracteis trigonis ſpinula terminatis. *Sauv. monſp.* 51.

Tithymalus myrſinites latifolius. *Bauh. pin.* 290.
Tithymalus myrſinites legitimus. *Cluſ. hiſt.* 2. *p.* 189.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 80.
Südliche Wolfsmilch. W.
Habitat in Calabria, Monſpelii. ♃ (v. ſ.)

Caules *multi, pedales, deflexi, virides, inferne a caſu foliorum citatriſati.* Folia *alterna, ſpathulata, coriacea, concava, glauco-viridia, patentia, mucronata, margine ſubſcabra, ſuperiora reflexa.* Umbella 7-9 *fida, radiis ſemel bifidis.* Involucrum *univerſale foliolis* 7-9, *ovatis, tenuioribus, acutis;* Involucella *diphylla, ſubcordata latiora, concava, acuta, margine ſubſcabra.* Flores *intra involucella primaria et ſecundaria maſculi, reliqui hermaphroditi.* Calyces *ore ſerrati.* Petala 4, *flava, bicornia apicibus teretibus, nitentia.* Capſulae *glabrae.*

116. EUPHORBIA *paluſtris.*

E. umbella multifida, ſubtrifida: bifida, involucellis, ovatis, foliis lanceolatis, ramis ſterilibus *Amoen. acad.* 3. *p.* 126.* *Mat. med.* 121. *Mill. dict. n* 22. *pall. it.* 1. *p.* 35 *Pollich. pal. n.* 462. *Hoff. germ.* 167. *Roth. germ. I.* 207. *II.* 534.
E. foliis lanceolatis, umbella univerſali multifida polyphylla: partialibus trifidis triphyllis; propriis dichotomis. *Hort. cliff* 200. *Fl. ſuec.* 438, 427.* *Roy. lugdb.* 108. *Dalib. pariſ.* 153 *Gort. gelr.* 307.

Tithymalus foliis ellipticis, ramis umbellam ſuperantibus, capſulis exaſperatis. *Hall. helv. n.* 1054.
Tithymalus paluſtris fruticoſus. *Bauh. pin.* 292. *Moriſ. hiſt. ſect.* 10. *t.* 2. *f.* 1.
Eſula major. *Dalech. hiſt.* 1653. *Dodon. purg. p.* 158.
Eſula paluſtris. *Riu. tetr.* 230.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 81.
Sumpf-Wolf-milch. W.
Habitat in Suecia *auſtrali,* Germania, Belgio. ♃ (v. v.)
Petala *integra.* Fructus *verrucoſus.*

*117. EUPHORBIA *pallida.*

E. umbella multifida: trifida, involucellis ſubrotundis, foliis lanceolatis attenuatis, caule ſimplici, petalis bicornibus, capſulis glabris. W.

Euphorbia amygdaloides. *Lumnitz poſon n.* 468. *deſcriptio. nec diagnoſis.*

Bleiche Wolfsmilch. W.

Habitat in Hungariae *ſylvaticis ſiccis.* ♃ (v. v.)

Caulis *ſimplex abſque ramis, in planta culta bipedalis glaber, in ſpontanea pedalis pubeſcens.* Folia *lanceolata utrinque praecipue apicem verſus attenuata, obtuſiuſcula, integerrima, ſubtus glauceſcentia, in culta planta omnino glabra, ſpontanea vero utrinque tenui pube tecta.* Umbella *multifida, radiis ſimpliciter trifidis brevibus, ſaepius vero in ſpontanea quadrifidis.* Involucri *foliola ovata.* Involucella *ſubrotunda acuta fere ad cordatam formam accedentia.* Petala *lunata.* Capſulae *glabrae.* W.

*118. EUPHORBIA *emarginata.*

E. umbella multifida: bifida, involucellis late cordatis, foliis oblongis emarginatis glabris, caule ramoſo, capſulis ſubverrucoſis. *Ait. Kew.* 2. *p.* 145.

Ausgerandete Wolfsmilch. W.

Habitat in Italia. ♃ (v. ſ.)

119. EUPHORBIA *hiberna.*

E. umbella ſexfida: dichotoma involucellis ovalibus, foliis integerrimis, ramis nullis, capſulis verrucoſis. *Amoen. acad.* 3. *p* 128. *Hudſ. angl.* 185. *Mill. dict. n.* 25. *Lepech. it.* 1. *p.* 221. *Kniph. cent.* 9. *n.* 37.

Tithymalus latifolius hiſpanicus. *Bauh. pin.* 291. *Tabern. hiſt.* 987.

Tithymalus hibernicus, vaſculis muricatis erectis. *Dill. elth.* 287. *t.* 290. *f.* 374.

Houttuyn Lin. Pfl Syſt. 7. *p.* 82.

Irrländiſche Wolfsmilch. W.

Habitat in Hibernia, Sibiria, Auſtria, Pyrenaeis. ♃

*120. EUPHORBIA *ſalicifolia.* †

E. umbella multifida: dichotoma, involucellis reniformi-cordatis, foliis lanceolatis villoſis. *Hoſt. ſynop.* 267.

Weidenblättrige Wolfsmilch. W.

Habitat ad ſylvarum margines Hungariae. ♃

Caulis *erectus ſtriatus.* Folia *ſeſſilia alternatim ſpar-*

ſa lanceolata, aut ovato-lanceolata, utrinque villoſa, obtuſa glauca. Involucri *foliola foliis aemula ſunt, ſed breviora.* Involucella *ſunt reniformi-cordata, apice compreſſa et uti tota pars ſuperior plantae, flaveſcentia.* Calyces *ventricoſi villoſi.* Petala *in omnibus floribus quatuor bicornia glabra.* Capſulae *glabrae rugoſulae.* Hoſt l. c.

121. EUPHORBIA *dendroides.*

E. umbella multifida; dichotoma, involucellis ſubcordatis: primariis triphyllis, caule arboreo. *Amoen. acad.* 3. *p.* 128. *Mill. dict. n.* 20.

Tithymalus myrtifolius arboreus. *Bauh. pin.* 290.

Tithymalus dendroides. *Cam. epit.* 965.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 83.

Baumartige Wolfsmilch. W.

Habitat in Italia, Creta, Inſulis Stoechadum. ♄ (v. ſ.)

Folia *ſparſa anguſtiora lanceolata.* Involucrum *univerſale polyphyllum, partialia prima ſaepius triphylla, reliqua diphylla, obtuſe cordata.* Umbellulae *laterales caulis nullae.* Capſula *glabra*

122. EUPHORBIA *amygdaloides.*

E. umbella multifida: dichotoma, involucellis perfoliatis orbiculatis, foliis obtuſis. *Amoen. acad.* 3. *p.* 126. *Mill. dict. n.* 21. *Hoſt. ſynop.* 266. *Hoffm. germ.* 167.

Euphorbia ſylvatica. *Iacq. auſt. t.* 275. *Scop. carn. n.* 570. *Roth. germ. I.* 206. *II.* 531.

Tithymalus characias amygdaloides. *Bauh. pin.* 290.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 83.

Mandelblättrige Wofsmilch. W.

Habitat in Gallia, Anglia, Germania. ♃ (v. ſ.)

Folia *alterna, remota, tenuia (non coriacea rigida).* Caulis *herbaceus, nec lignoſus.* Umbellae *etiam laterales.*

Vaillantius *negat hanc ſeparandam ab E. ſylvatica.*

Hujus umbella etiam quinquefida, ſed a ſequenti abunde diſtincta, caule *herbaceo; a reliquis Euphorbiae ſpeciebus* involucellis *connatis diverſiſſima. E. ſylvatica et E. Characias non creſcunt in Germania.* W.

123. EUPHORBIA *ſylvatica.*

E. umbella quinquefida: bifida, involucellis perfoliatis ſubcordatis acutiuſculis, foliis lanceolatis integerrimis. *Mill. dict. n.* 29.

Tithymalus foliis ellipticis, hirſutis, ſtipalis umbella-

ribus obſcuris, floralibus connatis, petalis lunatis. *Hall. helv. n.* 1045.

Tithymalus toto anno folia retinens. *Bauh. hiſt.* 3. *p.* 671.

Tithymalus lunato flore. *Col. ecphr.* 2. *p.* 56. *t.* 57.

Tithymalus ſylvaticus, lunato flore. *Bauh. pin.* 290. *Moriſ. hiſt.* 3. *p.* 335. *ſ.* 10 *t.* 1. *f.* 3.

Eſula caule craſſo. *Riv. tetr.* 227.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 84.

Wald-Wolfsmilch. W.

Habitat in Europa *auſtraliori.* ♄ (v. f.)

Caules *fruticoſi, proliferi, craſſi.* Folia *piloſa, perennia.* Petala *lunata, eroſa* (*E. Characiae vero integra*).

124. EUPHORBIA *Characias.*

E. umbella multifida: bifida, involucellis perfoliatis emarginatis, foliis lanceolatis integerrimis, caule fruteſcente. *Amoen. acad.* 3. *p.* 126.* *Mill. dict. n.* 24. *Kniph. cent.* 1. *n.* 29. *Iacq. ic. rar.* 1. *t.* 89. *Collect.* 1. *p.* 57.*

E. inermis, foliis lanceolatis, umbella univerſali multifida partialibus dichotomis, involucris ſemibifidis perfoliatis. *Hort. cliff.* 199. *Hort. upſ.* 142.* *Roy. lugdb.* 197.

Tithymalus characias rubens peregrinus. *Bauh. pin.* 290.

Tithymalus characias. *Cluſ. hiſt.* 2. *p.* 188.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 85.

Thal-Wolfsmilch. W.

Habitat in Gallia, Hiſpania, Italia. ♄ (v. v.)

Nimis *affinis E. ſylvaticae et amygdaloidi; varietates a ſpeciebus in his ulterius diſtinguendae a popularibus, mihi tales fuere:*

α. Caulis *fruticoſus, pubeſcens, ramis aliquot ex apice.* Folia *vix manifeſte pubeſcentia.* Involucella *prima bipartita; reliqua perfoliata, bifida.* Petala *integra, retuſa.* Frutex *hirti; huc.*

Tithymalus Characias folio lanuginoſo. *Bauh. pin.* 290.

β. Caulis *fruticoſus, obſolete pubeſcens, ſimplex.* Folia *patentia.* Pedunculi *pubeſcentes.* Petala *lunata, rubra.* Flores *parvi.*

γ. Caulis *fruticoſus, laevis, ſimplex.* Folia *reflexa, glabra, lucida.* Racemus *communis oblongus, ante antheſin nutans.* Pedunculi *glabri.* Involucella *omnia*

perfoliata, bifida. Petala *lunata, denticulata.* Flores *majuſculi.* Fructus *laeves.*

960. VISNEA.

Gen. plant. ed. Schreb. n. 833.

Cal. 5-phyllus inferus. *Cor.* 5-petala. *Stigmata* 3. *Nux* 2-ſ. trilocularis ſemiinfera.

1. VISNEA *Mocanera.*
V. Mocanera. *Lin. ſuppl.* 251.
Mocanera *Canar.*
Canariſche Viſnea. W.
Habitat in ſylvis montoſis inſularum Canarienſium. ♄ *Franc. Maſſon.*

Frurex *parvus.* Caulis *ſubverrucoſus teres.* Folia *alterna erecta breve petiolata elliptica glaberrima venoſa ſerrata, conſiſtentia Lauri.* Peduuculi *axillares ſolitarii nutantes vix petiolo longiores nudi uniflori.* Flores *parvi.* Corolla *lutea. Flore foecundato erigitur pedunculus, calyx clauditur et incraſſatur, ejusque foliola tria exteriora fuſca piloſa evadunt.*

TETRAGYNIA.

961. CALLIGONUM.

Gen. pl. ed. Schreb. n. 834.

Cal. 5-partitus. *Cor.* 0. *Filam.* circiter 16, baſi ſubcoalita. *Germen* ſuperum tetraëdrum. *Styli* 4. *Nux* cruſta polyplera ſ. potychaeta unilocularis.

1. CALLIGNUM *Polygonoides.*
C. fructibus cancellatis, ſetis ramoſis rigidis. *l'Herit. in Act. ſoc. Lin. Lond.* 1. *p.* 179.
C. polygonoides. *Sp. pl.* 748 *Hort. cliff.* 212.
Polygonoides orient. le Ephedrae facie. *Tournef. cor.* 47. *Itin.* 2. *p.* 356. *t.* 356.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 38.
Armeniſche Hackenknopf. W.
Habitat in monte Ararat. ♄

*2. CALLIGONUM *comoſum.* †
E. fructibus cancellatis, ſetis ramoſis mollibus. *l'Herit. in Act. ſoc. Lin. Lond.* 1. *p.* 180.
Aegyptiſcher Hackenknof. W.
Habitat in Aegypto, Barbaria. ♄
Varietas forte praecedentis. Plantae *in omnibus ſimillimae ſed in planta Aegyptiaca* fructus *comoſior, ſetis mollioribus, dum* ſetae *diſtinctiſſimae rigidiores in planta Tornefortiana, cujus unicum vidi fructum.* l'Herit. l. c.

2. CALLIGONUM *Pallaſia.*
E. fructibus alatis, alis membranaceis criſpis dentatis. *l'Herit. in Act. ſoc. Lin. Lond.* 1. *p.* 180. * *Ait. Kew.* 2. *p.* 242.
Pallaſia caſpica. *Lin. ſuppl.* 252.
Pallaſia Pteroccus. *Pall. fl. roſſ.* 2. *p.* 70. *t.* 77. 78. *
Pteroccus aphyllus. *Pall. itin.* 2. *p.* 738. *t. S.* *
Calligonum polygonoides. *Pall. itin.* 3. *p.* 536.
Caſpiſcher Hackenknopf W.
Habitat in deſerto inter Volgam *et* Rhymnum *verſus mare* Caſpium. ♄ (v. ſ.)
Frutex *quadripedalis ramoſiſſimus aphyllus articulatus, articulis interceptis ut Ephedrae.* Folia *nulla.* Flores *laterales ad genicula glomerati albi parvi.*

962. APONOGETON.

Gen. plant. edit. Schreb. n. 835.

Amentum e ſquamis compoſitum. *Cal.* 0. *Cor.* 0. *Capſulae* 4, triſpermae.

1. APONOGETON *monoſtachyon.*
A. ſpica ſimplici, foliis cordato-ovalibus. *Lin. ſuppl.* 214. *Thunb. diſſ. Nov. Pl. gen.* 4. *p.* 73. *c. icone.* *
Saururus (*natans*) acaulis, ſcapo monoſtachyo. *Mant.* 227. *
Parya Kelanga. *Rheed. mal.* 11. *p.* 31. *t.* 15.
β. A. (*lineare*) ſpica ſimplici, foliis linearibus. *Vahl. ſymb.* 3. *p.* 51.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 403.
Einähriges Aponogeton. W.
Habitat in Indiae *orientalis paludoſis inundatis, frequens inter Oryzam.* ♃ (v. ſ.)
Radix *bulboſa.* Folia *longiſſima petiolata radicalia*

cordata ovalia integerrima laevia natantia fere Potamogetonis natantis. Scapi *flaccidi obſolete trigoni.* Spica *ſimplex ſexangularis ſpatha caduca tecta.* Bracteae *corollae loco duae infra ſingulos floſculos parvae clavatae obtuſiuſculae carnoſae perſiſtentes.* Stamina *ſex ſubulata bracteis longiora.* Germina 3 *ſ.* 4 *oblonga acuta.* Stylus *germinibus vix brevior.* Stigma *obtuſiuſculum.*

*2. APONOGETON *criſpum.*

A. ſpica ſimplici, foliis enſiformibus criſpis. *Thunb. Diſſ. Nov. Pl. gen. p.* 73.

Krauſes Aponogeton. W.

Habitat in locis depreſſis Zeylona.

Radix *fibrilloſa.* Scapi *lineares foliis triplo longiores flexuoſi erecti glabri.* Folia *petiolata enſiformia acuminata criſpa glabra, petiolis longiora ſpithamaea.* Flores *ſpicati, ſpica interrupta pollicari.* Thunb. l. c.

3. APONOGETON *diſtachyon.*

A. ſpica bifida, foliis lineari-oblongis natantibus, bracteis integris, floribus polyandris. *Ait, Kew.* 1. *p.* 495.

A. ſpica bipartibili, foliis elliptico-lanceolatis. *Lin. ſuppl.* 215. *Thunb. Diſſ. Nov. Pl. gen.* 4. *p.* 74. *c. icone.* *

Zweyähriges Aponogeton. W.

Habitat in rivulis Cap. b. ſpei. *juxta Paarl.* ♃

Radix *bulboſa.* Folia *longiſſime petiolata radicàlia, lanceolato-ovata integerrima laevia natantia Potamogetonis natantis.* Spica *bipartibilis introrſum imbricata.* Flores *albi alterni erecti, bractea ovata excepti.* Stamina *multum variant a* 6 *ad* 12. Piſtilla 3 *ſ.* 4. Flores *odore fragrantiſſimo.* Bulbi *aſſati eduntur.*

Linnaeus in ſupplemento hanc ſpeciem cum ſequenti confundit, dum Stamina 6. *ad* 12 *variare dicit.* Ait. l. c.

*4. APONOGETON *anguſtifolium.*

A. ſpica bifida, foliis lineari-lanceolatis erectis, bracteis bipartitis, floribus hexandris. *Ait. Kew.* 1. *p.* 495.

Schmalblättriges Aponogeton. W.

Habitat ad Cap. b. ſpei. ♃

Folia *anguſtiora quam in praecedenti utrinque attenuata.* Raches *pallide rubrae.* Flores *pauci.* Bracteae *albae baſi rubrae, ad baſin fere bipartitae (vel ſi mavis binae) laciniae lineari-oblongae.* Stamina 6, *bracteis ter vel quater breviora.* Styli *tres.* Ait. l. c.

PENTAGYNIA.

963. GLINUS.

Gen. pl. ed. Schreb. n. 836.

Cal. 5-phyllus. *Cor.* 0. *Nectaria* ſetis bifidis. *Capſ.* 5-angularis, 5-locularis, 5-valvis, polyſperma.

1. GLINUS *lotoides.*

G. caule piloſo, foliis obovatis. *Loefl. it.* 145. *Gron. orient.* 143. *Burm. ind. n.* 2. *t.* 36 *f.* 1.

Alſine lotoides ſicula. *Bocc. ſic.* 21. *t.* 11. *Moriſ. hiſt.* 2. *p.* 552.

Alſine heliotropio cognata. *Cup. cath.*

Portulaca baetica, luteo flore, ſpuria aquatica. *Barr. rar.* 478. *t.* 336.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 87. *t.* 52. *f.* 3.

Schotenkleeartiger Glinus. W.

Habitat ab Hiſpania *in* Aſiam, *ad foſſas, locis inundatis.* ⊙

*2. GLINUS *ſetiflorus.*

G. foliis obovatis plicatis, caule fruticoſo, foliolis calycinis ovatis. *Vahl. ſymb.* 3. *p.* 64.

G. (*ſetiflorus*) caulibus diffuſis, floribus capitatis axillaribus ſeſſilibus. *Forſk. deſcr.* 95.*

Borſtenblütiger Glinus. W.

Habitat in ſylvis tempore pluvioſo inundatis Arabiae *felicis.* ♄

Habitus reliquorum. Tota planta e pilis denſiſſimis incana, pili longiores quam in G. lotoide. Vahl. l. c.

3. GLINUS *dictamnoides.*

G. foliis orbiculatis tomentoſis planis, caule fruteſcente proſtrato, foliolis calycinis lanceolatis. *Vahl. ſymb.* 3. *p.* 64.

G. caule rigoſo, foliis orbiculatis tomentoſis. *Mant.* 243. *Burm. ind.* 113. †

Alſine lotoides, dictamni cretici facie, maderaſpatana. *Pluk. amal.* 10 *t.* 256. *f.* 6.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 88. *t.* 52. *f.* 4.

Diptamartiger Glinus. W.

Habitat in India. ♄ v. ſ.)

Conf *Pluk. t.* 12 *f.* 3.

*964. BLACKWELLIA.

Juſſ. gen. ed. Uſteri p. 379.

Cal. 5-fidus ſemiſuperus. *Cor.* 15-petala. *Capſ.* 1-locul. polyſperma.

*1. BLACKWELLIA *integrifolia.*
B. foliis ovatis obtuſis pleriſque integerrimis, floribus terminalibus paniculatis. *Lamarck. encycl.* 1. *p.* 423.* *Illuſt. gen. t.* 412. *f.* 2.
Ganzblättrige Blackwellie. W.
Habitat in inſula Mauritii. ♄

*2. BLACKWELLIA *paniculata.*
B. foliis ovato-ſubrotundis dentatis, floribus paniculatis terminalibus. *Lamarck. encycl.* 1. *p.* 423.*
Vermontea decandra. *Comerſ. herb.*
Riſpenblütige Blackwellie. W.
Habitat in inſula Borbonia. ♄
Valde ſimilis praecedenti, ſed folia *multo minora fere ſubrotunda, dentibus diſtantibus inſtructa,* florum *panicula in ramorum apice magis compoſita et divaricata.* Lamarck l. c.

*3. BLACKWELLIA *axillaris.*
B. foliis ovatis ſubcrenatis, ſpicis axillaribus longis ſimpliciſſimis nutantibus. *Lamarck. encycl.* 1. *p.* 423.* *Illuſt. gen. t.* 412. *f.* 1.
Winkelriſpige Blackwellie. W.
Habitat in Madagaſcar. ♄
Spicae *floribus parvis confertis fere ſeſſilibus tomentoſis vel plumoſis, tectae.* Lamarck. l. c.
Affinitas huic generi eſt magna cum Homalio. W.

DODECAGYNIA.

965. SEMPERVIVUM.

Gen. plant. edit. Schreb. n. 837.

Cal. 12-partitus. *Petala* 12. *Capſ.* 12, polyſpermae.

1. SEMPERVIVUM *arboreum.*
S. caule arboreſcente laevi ramoſo, foliis cuneiformibus glabriuſculis ciliatis, ciliis patulis mollibus. *Ait. Kew.* 2. *p.* 147.

S. caule arboreſcente laevi ramoſo. *Sp. pl.* 664.
S. caule inferne nudo laevi ramoſo. *Hort. cliff.* 179. *Hort. upſ.* 118. *Roy. lugdb.* 456. *Mill. dict. n.* 5
Sedum majus arboreſcens, floſculis candidis. *Bauh. pin.* 282. *Bradl ſucc.* 4. *p.* 1. *t.* 31.
Sedum majus legitimum. *Cluſ. hiſt* 2. *p.* 58.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 89.
Baumartige Hauswurz. W.
Habitat in Luſitania, Creta, Corcyra, Zacintho. ♄ (v. v.)
Caulis *ſaepe radices filiformes dimittit.* Thyrſus *terminalis:* Racemis *bipartitis: Racemi brachia ramoſa, ſimplicia, recurva:* Floribus *alternis, oppoſitifoliis, pedicellatis, flaviſſimis.* Calyx. 10 *partitus.* Petala 10. Piſtilla 10. Nectarii *ſquamae totidem, ad filamentorum baſin.* Stamina 20.

2. SEMPERVIVUM *canarienſe.*

S. caule fruteſcente, foliis orbiculato-ſpathulatis villoſis, nectariis ſubquadratis truncatis. *Ait. Kew.* 2. *p.* 147.
S. caule foliorum ruderibus lacero, foliis retuſis. *Sp. pl.* 664. *Mill. dict. n.* 6.
S. caule infra folio lacero, foliis cuneiformibus: ſummis ſubrotundis. *Hort. cliff.* 179. *Roy. lugdb* 457.
Sedum canarium, foliis omnium maximis. *Comm. hort.* 2. *p.* 189. *t* 95.
Sedum majus canarium acaulon, pilis ad oras foliorum hiſpidis. *Pluk alm* 340. *t.* 314. *f.* 1. *Raj. ſuppl.* 362.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 98.
Canariſche Hauswurz W.
Habitat in Canariis. ♄ (v. v. ſ. fl.)
Caulis *brevis, ſimplex.*
Petala 9. Stamina 18. Piſtilla 9. Ait. l. c.

*3. SEMPERVIVUM *glutinoſum.*

S. caule fruteſcente, foliis cuneiformibus viſcidis ciliatis, ciliis cartilagineis adpreſſis. *Ait. Kew.* 2. *p.* 147.
Klebrige Hauswurz. W.
Habitat in Madera ♄ (v. v.)
Petala *lutea* 8 *vel* 9. Stamina 16 *ſeu* 18. Piſtilla 8 *vel* 9. Ait. l. c.

*4. SEMPERVIVUM *glanduloſum.*

S. caule fruteſcente, foliis orbiculato-ſpathulatis margine glutinoſis, glandulis globoſis, nectariis cuneiformibus truncatis. *Ait. Kew.* 2. *p.* 148.

Drüſige Hauswurz. W.
Habitat in Madera. ♄

5. SEMPERVIVUM *tectorum.*
S. foliis ciliatis, propaginibus patulis, floribus dodecandris dodecagyniis, nectariis cuneiformibus carunculatis. *Ait. Kew.* 2. *p.* 148.
S. foliis ciliatis, propaginibus patentibus. *Mat. med.* 121. *Mill. illuſtr. Syſt. Pollich. pal. n.* 464. *Flor. dan. t.* 601. *Blackw. t.* 366. *Hoffm. germ.* 168. *Roth. germ. I.* 208. *II.* 536.
Sedum roſulis glabris, ciliatis, petalis quatuordecim conglutinatis, lanceolatis hirſutis. *Hall. helv. n.* 949.
Sempervivum majus. *Neck. gallob. p.* 213.
Sedum tectorum. *Scop. carn. ed.* 2. *n.* 559.
Sempervivum foliis radicalibus carnoſis ciliatis, caulinis imbricatis membranaceis, corymbo racemoſo reflexo. *Hort. cliff.* 179. *Fl. ſuec.* 395, 418. *Roy. lugdb.* 457.
S. majus alterum ſ. Barba jovis. *Dod. pempt.* 127.
Sedum majus vulgare. *Bauh. pin.* 283.
Houttuyn Lin. Pſl. Syſt. 7. *p.* 91.
Gemeine Hauswurz. W.
Habitat in Europae *tectis et collibus.* ♃ (v. v.)

6. SEMPERVIVUM *globiferum.*
S. foliis ciliatis, propaginibus globoſis. *Mill. dict. n.* 2. *Jacq. auſtr.* 5. *p.* 50. *tab app.* 40. *Knorr. del.* 2. *t.* S. 4. *Doerr. naſſ. p.* 212. *Hoffm. germ.* 168. *Roth. germ. II.* 536.
S. foliis radicalibus in globum congeſtis ciliatis, propaginibus globoſis. *Hort. cliff.* 180. *Roy. lugdb.* 457.
S. petalis fimbriatis, ſobole compacta. *Schmid. ic.* 95. *t.* 26 *
Sedum roſulis hirſutis ciliatis, petalis ſubulato-lanceolatis duodenis. *Hall. helv. n.* 950.
Sedum majus vulgari ſimile, globulis decidentibus. *Moriſ.* 3. *p.* 472. *ſ.* 12. *t.* 7. *f.* 18.
Sedum vulgari magno ſimile. *Bauh. hiſt.* 3. *p.* 688.
Houttuyn Lin. Pſl. Syſt. 7. *p.* 92.
Kugelknoſpige Hauswurz. W.
Habitat in Rutheno. *D. Gmelin*; Auſtria, Germania. ♃ (v. v.)

Petala 6. Stamina 6. Piſtilla 6. *Jacq. Mygind.* Petala 12, *ſubulato-lanceolata, flava, hirſuta, ciliata. Haller.*

*7. SEMPERVIVUM *villoſum.*

S. foliis ſpathulato cuneiformibus obtuſis villoſis, nectariis palmatis, laciniis ſubulatis. *Ait. Kew.* 2. *p.* 148.

Haarige Hauswurz. W.

Habitat in Madera. ⊙.

Petala 8 *flava.* Stamina 12-16. Piſtilla 8. Ait. l. c.

*8. SEMPERVIVUM *tortuoſum.*

S. foliis obovatis ſubtus gibbis villoſis, nectariis bilobis. *Ait. Kew.* 2. *p.* 148.

Gedrehte Hauswurz. W.

Habitat in Canariis. ♄

Petala 8 *flava.* Stamina 16. Piſtilla 8. Ait. l. c.

*9. SEMPERVIVUM *ſtellatum.*

S. caule herbaceo pubeſcente, foliis ſpathulatis ſparſis. *Smith. Act. ſoc. Lin. Lond.* 1. *p.* 251.

Sedum petraeum rotundifolium, flore luteo ſtellato montis Baldi. *Segn. veron.* 2. *p.* 360. *t.* 17.

Sternförmige Hauswurz. W.

Habitat in rupibus montis Baldi. ⊙

Tota planta in ſtatu minimo repraeſentat S. arboreum; ſed eſt laxior et magis diffuſa. Ab hoc vero ſatis diſtinctum; radice *annua,* caule *herbaceo pubeſcente, et* panicula *divaricata.* Flores *flavi cum charactere Sempervivi exacte conveniunt.* Smith l. c.

10. SEMPERVIVUM *arachnoideum.*

S. foliis pilis intertextis, propaginibus globoſis. *Mill. dict. n.* 4. *Knorr. del.* 2. *t. S.* 8. *Iacq. auſt.* 5. *p.* 51. *tab. app.* 42.

S. foliis radicalibus in globum congeſtis: villis reticulatim connexis. *Hort. cliff.* 80. * *Roy. lugdb.* 457.

S. petalis integris, ocellis turuloſis. *Schmidel. icon. p.* 149. *t.* 38. *

S. rubrum montanum gnaphaloides. *Col. ecphr.* 1. *p.* 292. *t.* 291.

Sedum roſulis araneoſis, calveſcentibus, petalis connatis nervoſis. *Hall. helv. n* 952.

Sedum montanum tomentoſum. *Bauh. pin.* 294.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 93.

Spinnewebenartige Hauswurz. W.

Habitat in alpibus Italiae, Helvetiae, Pyrenaeis. ♃ (v. v.)

Petala 9, *purpurea;* Varietas *S. montani. Haller.* Foliorum *faſciculi contracti, obtecti pilis longis. Gerard.*

11. SEMPERVIVUM *hirtum.*

S. foliis caule petalorumque apicibus hirtis. *Amoen. acad.* 4. *p* 273. * *Hoffm. germ.* 168.

Sedum majus montanum, foliis dentatis. *Bauh. pin.* 283. *Raj. hiſt.* 1. *p.* 688.

Cotyledon altera montana. *Cluſ. hiſt.* 2. *p,* 63.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 94.

Rauhblütige Hauswurz. W.

Habitat in Helvetia, in Tauero *raſtadienſi.* ♃

Petala 6; Stamina 12; Piſtilla 6 *Mygind.*

12. SEMPERVIVUM *montanum.*

S. foliis integerrimis, propaginibus patulis. *Mill. dict. n.* 3. *Mattuſch. ſil.* 345. *Hoffm. germ.* 169. *Iacq. auſt.* 5. *p.* 5. *tab. app.* 41.

Sedum foliis ſemipatulis hirſutis, petalis novenis ſubhirſutis. *Hall helv. n.* 951.

Sedum alpinum, rubro magno flore. *Bauh. pin.* 284.

Sedum majus montanum, foliis non dentatis, floribus rubentibus. *Bauh. pin.* 283.

Sedum minus, flore rubente. *Beſl. eyſt. vern.* 6. t. 8. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 94.

Berg-Hauswurz. W.

Habitat in rupibus Helvetiae, Sileſiae. ♃ (v. ſ.)

Foliorum *faſciculi expanſi, pilis diſtituti. Gerard.*

13. SEMPERVIVUM *ſediforme.*

S. foliis ſparſis, inferioribus teretibus; ſuperioribus depreſſis. *Syſt. Veg.* 379. *Iacq. hort.* 1. *p.* 35. *t.* 81.

β. Monſtroſum. *Iacq. miſc.* 1. *p.* 133. *t.* 5. *

Houttuyn Lin. Pfl. Syſt. 7. *p.* 95.

Sedumartige Hauswurz. W.

Habitat in Europa *auſtrali.* ♃

Petala *albida,* Caules *omnes perpetuo et conſtanter ſtrictiſſime ſtant erecti.* Iacq.

*14. SEMPERVIVUM *monanthos.*

S. foliis teretibus clavatis confertis, pedunculis nudis ſubunifloris, nectariis obcordatis. *Ait. Kew.* 2. *p.* 149.

Einblütige Hauswurz. W.

Habitat in Canariis. ♃ (v. v.)

Numerus partium fructificationis variat a quinario ad octonarium. Ait l. c.

Claſſis XII.

I C O S A N D R I A.

M O N O G Y N I A.

966. CACTUS. *Cal.* ſuperus, 1-phyllus. *Cor.* multifida. *Bacca* 1-locul. polyſperma.
972. EUGENIA. *Cal.* ſuperus, 4-partitus. *Cor.* 4-petala. *Bacca* 1-locularis, 1-ſperma.
967. PHILADELPHUS. *Cal.* ſuperus, 5-ſ. 4-partitus. *Cor.* 5-ſ. 4-petala. *Stigma* 4-fidum. *Capſ.* 5-4-locul. polyſperma.
968. LEPTOSPERMUM. *Cal.* ſuperus 5-fidus. *Petala* 5 unguiculata ſtaminibus longiora. *Stigma* capitatum. *Capſ.* 4-ſ. 5-loculaais.
969. FABRICIA. *Cal.* ſuperus 5-fidus. *Petala* 5 ſeſſilia. *Stigma* capitatum. *Capſ.* multilocularis.
970. METROSIDEROS. *Cal* ſuperus 5-fidus. *Petala* 5. *Stam* longiſſima exſerta. *Stigma* ſimplex. *Capſ.* 3-*ſ.* 4-locularis.
971. PSIDIUM. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Bacca* 1 locularis, polyſperma.
973. MYRTUS. *Cal.* ſuperus, 5-fidus. *Cor.* ſub 5-petala. *Bacca* 3-locularis, polyſperma.
980. PUNICA. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Pomum* 10-loculare, polyſpermum.
985. ROBINSONIA. *Cal.* ſuperus 5-dentatus. *Petala* 5. *Bacca* ſtriata 7-locularis.
974. CALYPTRANTHES. *Cal.* ſuperus truncatus operculo tectus. *Cor.* o. *Bacca* 1-locul. 1-4-ſperma.
975. EUCALYPTUS. *Cal.* ſuperus truncatus operculo tectus. *Cor.* o. *Capſ.* 4-locularis polyſperma.
978. FOETIDIA. *Cal.* ſuperus 4-fidus. *Cor.* o. *Capſ.* 4-locularis lignoſa.
986. SONNERATIA. *Cal.* inferus 6-fidus. *Petala* 6. *Bacca* multilocularis, loculis polyſpermis.
981. AMYGDALUS. *Cal.* inferus 5-fidus. *Cor.* 5-petala. *Drupa* nucleo foraminoſo.
982. PRUNUS. *Cal.* inferus, 5-fidus. *Cor.* 5-petala. *Drupa* nucleo integro.

984. CHRYSOBALANUS. *Cal.* inferus, 5-fidus. *Cor* 5-petala *Drupa* nucleo ſulcato.

983. PLINIA. *Cal.* inferus 4-ſ. 5-partitus. *Cor.* 4-ſ. 5-petala. *Drupa* ſulcata.

976. BANARA. *Cal.* inferus 4-fidus. *Petala* 4. *Bacca* 1-locularis polyſperma.

977. ANTHERYLIUM. *Cal* inferus 4-partitus. *Petala* 4. *Capſ.* 1-locularis 3-valvis polyſperma.

979. SCOLOPIA. *Cal.* inferus 3-ſ. 4-partitus. *Petala* 3. ſ. 4. *Bacca* 1-locularis. *Semina* arillata.

† *Cleome icoſandra.*

DIGYNIA.

987. CRATAEGUS. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Bacca* 2-ſperma.

988. WALDSTEINIA. *Cal.* 10-fidus, laciniis alternis. *Petala* 5. *Semina* 2 obovata.

TRIGYNIA.

989. SORBUS. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Bacca* 3-ſperma.

990. SESUVIUM. *Cal.* inferus, 5-fidus. *Cor.* nulla. *Capſ.* 3-locularis, circumſciſſa.

† *Spiraea opulifolia.*

TETRAGYNIA.

† *Tetragoniae aliquot. Meſembryanthemum.*

PENTAGYNIA.

993. TETRAGONIA. *Cal.* ſuperus, 5-ſ. 4-fidus. *Cor.* o. *Peric.* nucleo 5. ſ. 4-loculari.

991. MESPILUS. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Bacca* 5-ſperma.

992. PYRUS. *Cal.* ſuperus, 5-fidus. *Cor.* 5-petala. *Pomum* 5-loculare, polyſpermum.

994. MESEMBRYANTHEMUM. *Cal.* ſuperus, 5-fidus. *Cor.* multifida. *Capſ.* carnoſa locularis, polyſperma.

995. AIZOON. *Cal.* inferus, 5-fidus. *Cor.* o. *Capſ.* 5-locularis, polyſperma.

996. SPIRAEA. *Cal.* inferus, 5-fidus. *Cor.* 5-petala. *Capſ.* plures congeſtae.

O C T O G Y N I A.

† *Meſembryanthemum calamiforme.*

D E C A G Y N I A.

† *Meſembryanthema aliquot.*

P O L Y G Y N I A.

997. ROSA. *Cal.* 5-fidus. *Cor.* 5-petala. *Cal.* baccatus, polyſpermus.

998. RUBUS. *Cal.* 5-fidus. *Cor.* 5-petala. *Bacca* compoſita.

1001. TORMENTILLA. *Cal.* 8-fidus. *Cor.* 4-petala. *Sem.* 8, mutica.

1003. DRYAS. *Cal.* 8-fidus. *Cor.* 8-petala. *Sem.* plurima: ariſta lanata.

999. FRAGARIA. *Cal.* 10-fidus. *Cor.* 5-petala. *Sem.* plurima ſupra receptaculum baccatum, decidua.

1000. POTENTILLA. *Cal.* 10-fidus. *Cor.* 5-petala. *Sem.* plurima, mutica.

1002. GEUM. *Cal.* 10-fidus. *Cor.* 5-petala. *Sem.* plurima: ariſta geniculata.

1004. COMARUM. *Cal.* 10-fidus. *Cor.* 5-petala. *Sem.* plurima ſupra receptaculum carnoſum perſiſtens.

1005. CALYCANTHUS. *Cal.* ſquamoſus, corollinus. *Cor.* 0. *Sem.* caudata calyce ſucculento.

† *Spiraea Filipendula, Ulmaria. Phytolacca icoſandra. Meſembryanthema aliquot.*

MONOGYNIA.

966. CACTUS.

Gen. plant. edit. Schreb. n. 838.

Cal. 1-phyllus, ſuperus, imbricatus. *Cor.* multiplex. *Bacca* 1-locularis, polyſperma.

* *Echinomelocacti ſubrotundi.*

1. CACTUS *mammillaris.*
C. ſubrotundus tectus tuberculis ovatis barbatis. *Hort. cliff.* 181. *Hort. upſ.* 119. *Roy. lugdb.* 278. *Mill. dict. n.* 5.
Echino-Melocactus minor lacteſcens; tuberculis ſ. mammillis majoribus. *Herm. par.* 136. *t.* 136.
Ficoides ſ. Melocactus mammillaris glabra ſulcis carens fructum ſuum undique fundens. *Pluk. alm.* 184. *t.* 29. *f.* 1.
Ficoides ſ. Ficus americana ſphaerica tuberculata lacteſcens, flore albo. *Comm. hort.* 1. *p.* 105. *t.* 55. *Bradl. ſucc.* 3. *p.* 11. *t.* 19.
β. Melocactus minimus lanuginoſus et tuberoſus. *Plum. ſpec.* 19. *ic.* 201. *f.* 1.
γ. Ficoides ſ. Melocactus minima lanuginoſa, ſpinis mitioribus, fructu ſparſim egrediente. *Pluk. alm.* 284. *t.* 29. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 622.
Warzentragende Fackeldiſtel. W.
Habitat in Americae *calidioris rupibus.* ♄ (v. v.)

2. CACTUS *Melocactus.*
E. ſubrotundus quatuordecim-angularis. *Hort. cliff.* 181. *Hort. upſ.* 119. *Roy. lugdb.* 279. *Mill. dict. n.* 1. *Swartz. obſ.* 198. *
Cactus humilis ſubrotundus ſulcatus et coronatus, ſpinis confertis. *Brown. jam.* 238.
Melocactus indiae occidentalis. *Bauh. pin.* 238.
Echino Melocactus. *Cluſ. exot.* 92. *t.* 92. *Bradl. ſucc.* 4. *p.* 9. *t.* 32.
β. Cactus ſubrotundus 14-angularis: ſpinis longis recurvis albidis. *Mill. dict.*
Houttuyn Lin. Pfl. Syſt. 3. *p.* 623.
Melonenartige Fackeldiſtel. W.
Habitat in Jamaica, America *calidiore.* ♄

3. CACTUS *nobilis.*
C. ſubrotundus quindecim-angularis, ſpinis latis recurvis. *Mant. p.* 243. *Mill. dict.* †
C. recurvus. *Mill. dict. p.* 490.
Edele Fackeldiſtel. W.
Habitat in Mexico. ♄
Corpus *ſubrotundum angulis* 14 (*nonne* 15? R). Puncta *dorſo angulorum inſident plura, faſciculata ſpinis aceroſis.* Spina, *cuiusvis faſciculi aceroſi medio, magna, recurvata.*

* * Cerei *erecti, ſtantes, per ſe.*

*4. CACTUS *cylindricus.*
C. erectus debilis cylindricus non angulatus, ſuperficie ſulcis decuſſatis reticulata. *Lamarck. encycl.* 1. *p.* 533. *
Walzenförmige Fackeldiſtel. W.
Habitat in Peru. ♄ (v. v. ſ. fl.)

5. CACTUS *Pitaiaya.*
D. erectus triangularis. *Iacq. amer.* 151.
Houttuyn Lin. Pfl. Syſt. 3 *p.* 624.
Aufrechte Fackeldiſtel. W.
Habitat Carthagenae *in fruticetis maritimis.* ♄

6. CACTUS *heptagonus.*
C. erectus oblongus ſeptemangularis. *Hort. cliff.* 181. * *Roy. lugdb.* 279.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 625.
Siebeneckige Fackeldiſtel. W.
Habitat in America. ♄ (v. ſ.)
Hiſtoria Cactorum angulatorum, *quorum plurimas ſpecies Plumier in iconibus ſiſtit, etiamnum obſcura permanſit, quam itaque attenti per Indias peregrinatores graphice deſcribant.*

7. CACTUS *tetragonus.*
C. quadrangularis longus erectus: angulis compreſſis. *Hort. cliff.* 181. *Hort. upſ.* 119. *Roy. lugdb.* 280. *Mill. dict. n.* 2. *ſub.* Cereo.
Cereus erectus minor, fructu ſpinoſo, coſtarum numero varians. *Herm. par.* 117.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 626.
Viereckige Fackeldiſtel. W.
Habitat in Curaſſao, America *calidiore.* ♄ (v. v. ſ. fl.)

8. CACTUS *hexagonus.*

C. erectus ſexangularis longus: angulis diſtantibus. *Hort. cliff.* 181. *Hort. upſ.* 119. *Roy. lugdb.* 279. *Mill. dict. n.* 1. *ſub* Cereo. *Fabric. helmſt.* 339. *Knorr. del.* 1. *t. F.* 7.

Cereus ſurinamenſis. *Eph. N. C.* 3. *p.* 349. *t.* 7, 8.

Cereus erectus altiſſimus ſurinamenſis. *Herm. par.* 116. *Raj. dendr.* 23.

Cereus erectus maximus americanus hexangularis, flore albo radiato. *Bradl. ſucc.* 1. *p.* 1. *t.* 1.

Melocactus monoclonos, flore albo, fructu atropurpureo. *Plum. ſpec.* 19. *ic.* 119?

Houttuyn Lin. Pfl. Syſt. 3. *p.* 626.

Sechseckige Fackeldiſtel. W.

Habitat Surinami. ♄

9. CACTUS *pentagonus.*

C. erectus ſubquinquangularis longus articulatus. *Hort. cliff.* 182. *. *Roy. lugdb.* 280.

Houttuyn Lin. Pfl. Syſt. 3 *p.* 627.

Funfeckige Fackeldiſtel. W.

Habitat in America. ♄

10. CACTUS *repandus.*

C. erectus longus octangularis: angulis compreſſis undatis: ſpinis lana longioribus. *Hort. cliff.* 182. *Roy. lugdb.* 279.

Cactus erectus cylindricus ſulcatus tenuior ſummitate attenuatus, aculeis confertis. *Brown. jam.* 238.

Cereus altiſſimus gracilior, fructu extus luteo, intus niveo ſeminibus nigris pleno. *Sloan. jam.* 197. *hiſt.* 2. *p.* 158. *Raj. dendr.* 22. *Trew. ehret. t.* 14.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 628.

Ausgeſchweifte Fackeldiſtel. W.

Habitat in America *calidiore.* ♄ (v. v. ſ. fl.)

11. CACTUS *lanuginoſus.*

C. erectus longus ſubnovemangularis: angulis obſoletis, ſpinis lana brevioribus. *Hort. cliff.* 182. *Roy. lugdb.* 279.

Cereus curaſſavicus erectus maximus, fructu rubro non ſpinoſo, lanugine flaveſcente. *Herm. par.* 115. *t.* 115.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 628.

Wollige Fackeldiſtel. W.

Habitat in Curaſſao. ♄

12. CACTUS *peruvianus.*

C. erectus longus ſuboctangularis: angulis obtuſis. *Hort. cliff.* 181. *Hort. upſ.* 120. *Roy. lugdb.* 279. *Swartz. obſ.* 199. *

C. cylindraceus erectus ſulcatus major ſummitate obtuſus, aculeis confertis. *Brown. jam.* 238.

Cereus erectus, fructu rubro non ſpinoſo. *Herm. par.* 114.

Cereus craſſiſſimus, fructu utrinque rubro. *Sloan. cat.* 196.

Cereus peruvianus ſpinoſus, fructu rubro nucis magnitudine. *Bauh. pin.* 458.

Euphorbii arbor cerei effigie. *Lob. ic.* 2. *p.* 25.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 629.

Peruvianiſche Fackeldiſtel. W.

Habitat in Jamaicae, Peru *apricis aridis maritimis.* ♄ Anguli *ſuperne inermes. Mill.*

13. CACTUS *Royeni.*

C. erectus articulatus novemangularis: articulis ſubovatis, ſpinis lanam aequantibus. *Roy. lugdb.* 279.

Cereus erectus gracilis ſpinoſiſſimus: ſpinis flavis, polygonus, lanugine albo-palleſcente. *Boerh. lugdb.* 1. *p.* 293.

Cereus erectus, fructu rubro non ſpinoſo, lanuginoſus. *Herm. parad.* 115.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 629.

Royenſche Fackeldiſtel. W.

Habitat in America. ♄

* * Cerei *repentes, radiculis lateralibus.*

14. CACTUS *grandiflorus.*

C. repens ſubquinquangularis. *Herm. par.* 120. *Knorr. del.* 1. *t. F.* 6.

C. ſcandens, angulis quinque pluribusque obtuſis. *Hort. cliff.* 182. *Hort. upſ.* 121.* *Roy. lugdb.* 280.

Cereus ſcandens minor polygonus articulatus. *Mill. icon. t.* 90.

Cereus gracilis ſcandens ramoſus plerumque ſexangularis, flore igenti atque fragranti. *Trew. ehr. t.* 31, 32. *E. N. C.* 1752. *v.* 9. *app.* 184. *t.* 11, 12, 13.

Cereus americanus major articulatus, flore maximo noctu ſe aperiente ſ. ſuaviſſimum odorem ſpirante. *Volk. heſp.* 1. *p.* 133. *t.* 134.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 630.
Groſsblumige Fackeldiſtel. W.
Habitat in Jamaica, Vera Cruce. ♄ (v. v.)

15. CACTUS *flagelliformis.*
C. repens decemangularis. *Mill. dict. n.* 12. *ſub* Cereo *Kniph. cent.* 1. *t.* 12. *Knorr. del.* 1. *t. F.* 8.
C. ſcandens angulis decem obſoletis. *Hort. upſ.* 121.
C. cylindraceus ſulcatus puſillus repens, aculeis ſetaceis confertis. *Brown. jam.* 238.
Cereus minor ſcandens polygonus ſpinoſiſſimus, flore purpureo. *Ehret. ſel.* 2. *f.* 2. *Trew. ehret. t.* 30.
Opuntia minima ſerpens americana. *Sloan. cat.* 197.
Ficoides americanum ſ. Cereus minima ſerpens americana. *Pluk. alm.* 148. *t.* 158. *f.* 6. *Sloan. jam.* 197. *hiſt.* 2. *p.* 158.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 631.
Peitſchenförmige Fackeldiſtel. W.
Habitat in America calidiore. ♄ (v. v.)

*16. CACTUS *pendulus.*
C. pendulus, ramis verticillatis teretibus glabris muticis. *Swartz prodr.* 77. *Fl. ind. occd.* 2. *p.* 876.* *Ait. Kew.* 2. *p.* 153.
C. paraſiticus inermis aphyllus ramoſus propendens, ramulis gracilibus teretibus ſtriatis. *Brown. jam.* 238.
Rhipſalis Caſſutha. *Gaert. ſem.* 1. *p.* 137. *t.* 28. *f.* 1.
Caſſyta baccifera. *J. Mill. illuſt.*
Hängende Fackeldiſtel. W.
Habitat in ramis arborum altiſſimarum Jamaicae *et* Hiſpaniolae. ♄ (v. v. ſ. fl.)

17. CACTUS *paraſiticus.*
C. repens teres ſtriatus muticus. *Sp. pl.* 668. *Swartz obſ.* 200.
Opuntia minima flagelliformis. *Plum. ſpec.* 6. *ic.* 197. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 932.
Schmarotzer Fackeldiſtel. W.
Habitat in America.

18. CACTUS *triangularis.*
C. repens triangularis. *Iacq. amer.* 152. *Riſler in Act. helv.* 5. *p.* 268. *t.* 2.*
C. triangularis ſcandens articulatus. *Hort. cliff.* 182. *Hort. upſ.* 121. *Roy. lugdb.* 280.

C. debilis brachiatus aequalis triquetrus ſcandens ſ. repens, ſpinis breviſſimis confertis. *Brown jam.* 468.
Cereus americanus triangularis radicoſus. *Brad. ſucc.* 1. *p.* 4. *t.* 3. *E. N. C.* 1752. *v.* 9. *app.* 199. *t.* 10. *f.* 14. *E. N. C.* 1754. *v.* 9. *app.* 349. *t.* 3.
Ficoides americanum, ſ. Cereus erectus criſtatus, foliis triangularibus profunde canaliculatis. *Pluk. alm.* 147. *t.* 29. *f.* 3.
Melocactus americanus repens trigonus, flore albo, fructu violaceo. *Plum. ic.* 199, 200?
β. Cactus (*triangularis folioſus*) fructu folioſo, inſipido. *Iacq. amer.* 152. *t.* 181. *f.* 65.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 632.
Dreyeckige Fackeldiſtel. W.
Habitat in Braſilia, Jamaica, Martinica. ♄ (v. v.)
Var. β. Fructus *multo major*, *nitide coccineus*, *foliis veſtitus fere integris.* Pulpa *candida*, *ſubdulcis inſipida*, *edulis.* Semina *atra*, *nitida.* Iacq.

* * * * Opuntiae *compreſſae articulis proliferis.*

19. CACTUS *moniliformis.*

C. articulato-prolifer, articulis globoſis ſpinoſis glomeratis. *Sp. pl.* 669.
Melocactus, ex plurimis globulis opuntiae modo naſcentibus, ſpinoſiſſimus. *Plum. ſpec.* 20. *ic.* 198.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 634.
Halsbandförmige Fackeldiſtel. W.
Habitat in America *calidiore.* ♃

20. CACTUS *Opuntia.*

C. articulato-prolifer laxus, articulis ovatis, ſpinis ſetaceis. *Kniph. cent.* 8. *n.* 19. *Knorr. del.* 1. *Tab. F. a.*
C. compreſſus articulatus ramoſiſſimus, articulis ovatis, ſpinis ſetaceis. *Hort. cliff.* 183. *Hort. upſ.* 120. *Gron. virg.* 54. *Roy lugdb.* 280.
C. brachiatus et articulatus, articulis ovatis compreſſis, aculeis longiſſimis confertis. *Brown. jam.* 237.
Opuntia foliis ovatis compreſſis, ſpinis ſetaceis. *Hall. helv. n.* 1099.
Opuntia vulgaris. *Mill. dict.* 1
Opuntia vulgo herbariorum. *Bauh. hiſt.* 1. *p.* 154. *Mill. ic.* 191.
Ficus indica, folio ſpinoſo, fructu majore, *Bauh. pin.* 458.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 634.
Gemeine Fackeldiſtel. W.
Habitat in America, Peru, Virginia, *nunc* in Hiſpania, nia, Italia, Luſitania, Helvetia, Minorca. ♄ (v. v.)

21. CACTUS *Ficus indica.*

C. articulato-prolifer, articulis ovato-oblongis, ſpinis ſetaceis. *Sp. pl.* 669.
C. compreſſus articulatus ramoſus, articulis ovato-oblongis, ſpinis ſetaceis. *Hort. cliff.* 283. *Hort. upſ.* 120. *Roy. lugdb.* 280.
Opuntia articulis ovato-oblongis, ſpinis ſetaceis. *Mill. dict. n.* 2.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 636.
Indiſche Fackeldiſtel. W.
Habitat in America *calidiore.* ♄ (v. v. ſ. fl.)

22. CACTUS *Tuna.*

C. articulato-prolifer, articulis ovato-oblongis, ſpinis ſubulatis. *Mill. dict. n.* 3. *ſub* Opuntia. *Knorr. del.* 2. *t.* O.
C. compreſſus articulatus ramoſus, articulis ovato-oblongis, ſpinis ſubulatis. *Hort. cliff.* 183. *Hort. upſ.* 122. *Roy. lugdb.* 280.
C. brachiatus articulatus, articulis oblongo-ovatis compreſſis, caudice tereti erecto ferociſſimo, aculeis brachiorum brevibus confertis. *Brown. jam.* 237.
Tuna major, ſpinis validis flavicantibus, flore gilvo. *Dill. elth.* 396. *t.* 295. *f.* 380.
Opuntia major, folio oblongo rotundo: ſpinis longis et validiſſimis, flore luteo. *Sloan. jam.* 103. *hiſt.* 2. *p.* 149. *t.* 244. *f.* 1.
β. C. (*elatior*) articulato-prolifer, articulis ovato-oblongis, ſpinis ſubulatis caule ſubtereti arboreo. *Euphraſ. itin. ed. germ.* 198.
Tuna elatior ſpinis validis nigricantibus. *Dill. elth.* 2. *p.* 395. *t.* 294. *f.* 379.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 636.
Breite Fackeldiſtel. W.
Habitat in Jamaica *et* America *calidiore.* ♄ (v. v.)
Munimenta *urbium in inſula Euſtachii. Iacq. amer.* 92.

23. CACTUS *coccinellifer.*

C. articulato-prolifer, articulis ovato-oblongis ſubiner-

mibus. *Mill. dict. n.* 6. *ſub* Opuntia. *Knorr. del.* 2. *t. O.* 1.

C. compreſſus articulatus ramoſus, articulis ovato-oblongis ſubinermibus. *Hort. upſ.* 121.

C. brachiatus et articulatus ſubinermis major, articulis oblongis et leviter compreſſis. *Brown. jam.* 237.

Tuna mitior, flore ſanguineo, cochenillifera. *Dill. elth.* 399. *t.* 297. *f.* 383.

Ficus indica major laevis ſpinoſa vermiculos proferens. *Pluk. alm.* 146. *t.* 281. *f.* 2.

Opuntia maxima, folio oblongo rotundo majore ſpinulis nonnullis et innocentibus obſito. *Sloan. jam.* 194. *hiſt.* 2. *p.* 152. *t.* 8. *f.* 1. 2.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 637.

Cochenillen Fackeldiſtel. W.

Habitat in Jamaica *et* America *calidiore.* ♄

24. CACTUS *curaſſavicus.*

C. articulato-prolifer, articulis cylindrico-ventricoſis compreſſis. *Mill. dict. n.* 7. *ſub* Opuntia. *Knorr. del.* 2. *t. O.* 2.

Cactus tereti-compreſſus articulatus ramoſus. *Hort. cliff.* 182. *Hort. upſ.* 120. *Roy. lugdb.* 280.

Opuntia minima americana ſpinoſiſſima. *Bradl. ſucc.* 1. *p.* 5. *t.* 4.

Ficus indica ſ. Opuntia curaſſavica minima. *Comm. hort.* 1. *p.* 107. *t.* 56.

Ficus indica ſ. Opuntia minor cauleſcens arbuſculae in modum, ramis cineritiis, ſpinoſiſſima. *Pluk. alm.* 147. *t.* 281. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 638.

Curaſſaviſche Fackeldiſtel. W.

Habitat in Curaſſao. ♄ (v. v. ſ. fl.)

*25. CACTUS *ſpinoſiſſimus.*

C. caule erecto compreſſo, ramis oppoſitis bifariis compreſſis, ſpinis ſetaceis. *Ait. Kew.* 2. *p.* 155. *Martyn. hort. cantabr.* 88.

C. (*ſpinoſiſſimus*) ſuberectus compreſſus, articulis cruciatim diſpoſitis, ſpinis creberrimis longis flaveſcentibus. *Lamarck. encycl.* 1. *p.* 557. *

Vielſtachliche Fackeldiſtel. W.

Habitat in Jamaica. ♄ (v. v. ſ. fl.)

*26. CACTUS *alatus.*

C. prolifer, enſiformi-compreſſus crenatus. *Swartz prodr.* 77. *Fl. ind. occid.* 2. *p.* 878. *

C. mitis minor, ſarmento flexili rotundo, frondibus longis compreſſis crenatis, ad crena floridis. *Brown. jam* 237.

Opuntia non ſpinoſa caulescens e crenis foliorum fructum et florem proferens. *Sloan. cat.* 216. *hiſt.* 2. *p.* 159.

Geflügelte Fackeldiſtel. W.

Habitat in truncis et ad radices arborum Jamaicae *frigidioris.* ♄

27. CACTUS *Phyllanthus.*

C. prolifer enſiformi-compreſſus ſerrato-repandus. *Mill. dict. n.* 9. *ſub* Opuntia.

C. foliis enſiformibus obtuſe ſerratis. *Hort. cliff.* 183. *Roy. lugdb.* 281.

C. mitis minor, ſarmento flexili rotundo, frondibus longis compreſſis crenatis, ad crenas floridis. *Brown. jam.* 237.

Cereus ſcolopendri folio brachiato. *Dill. elth.* 73. *t.* 64. *f.* 74.

Opuntia non ſpinoſa minima caulescens, foliis piloſis ſtrictiſſimis e crenis foliorum florem proferens. *Sloan. jam.* 216. *hiſt.* 2. *p.* 159.

Phyllanthos americana, ſinuoſis foliis longis. *Pluk. alm.* 296. *t.* 247. *f.* 5.

Ficus ſ. Opuntia non ſpinoſa, ſcolopendriae folio ſinuato. *Raj. dendr.* 21.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 339.

Dünngliedrige Fackeldiſtel. W.

Habitat in Braſilia, Surinamo, America *meridionali.* ♄ (v. v. ſ. fl.)

28. CACTUS *Pereskia.*

C. caule tereti arboreo: aculeis geminis recurvis, foliis lanceolato-ovatis. *Hort. upſ.* 122.

C. ſarmentoſus foliatus ſpinoſus, ſpinis geminis recurvis, foliis mollibus ovatis. *Brown. jam.* 237.

Pereskia. *Hort. cliff.* 122. *Roy. lugdb.* 281. aculeata. *Mill. dict.*

Pereskia aculeata, flore albo, fructu flaveſcente. *Plum. gen.* 37. *Dill. elth.* 305. *t.* 227. *f.* 294.

Malus americana ſpinoſa, portulacae folio, fructu foliosſo, ſemine reniformi ſplendenti. *Comm. hort.* 1. *p.* 145. *t.* 70.

Portulaca americana latifolia ad foliorum ortum lanu-

gine obducta, longioribus aculeis horrida. *Pluk. alm.* 135. *t.* 215. *f.* 6.

Groſſulariae fructu majore arbor ſpinoſa, fructu folioſo viridi-albicante. *Sloan. jam.* 165, *hiſt.* 2. *p.* 86. *Raj. dendr.* 27.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 639.

Groſsblättrige Fackeldiſtel. W.

Habitat in America *calidiore*, Jamaica, Margaretha. ♄ (v. v. ſ. fl.)

29. CACTUS *portulacifolius.*

C. caule tereti arboreo ſpinoſo, foliis cuneiformibus retuſis. *Sp. pl.* 671.

Opuntia arbor ſpinoſiſſima, foliis portulacae cordatis. *Plum. ſpec.* 6. *ic.* 197. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 640.

Portulacblättrige Fackeldiſtel. W.

Habitat in America *calidiore.* ♄

Fructus hujus minime foliis adſperſus, uti praecedentis.

967. PHILADELPHUS.

Gen. plant. ed. Schreb. n. 840.

Cal. 4-ſ. 5-partitus, ſuperus. *Petala* 4 ſ. 5. *Styl.* 4-fidus. *Capſ.* 4-ſ. 5-locularis, polyſperma.

1. PHILADELPHUS *coronarius.*

P. foliis ſubdentatis. *Mill. dict. n.* 1. *Hall. helv. n.* 1100. *Du H. m. arb.* 2. *t.* 83. *Kniph. cent* 5. *n.* 65. *Hoffm. germ.* 169. *Moench. haſſ. n.* 401. *Willd. arb.* 204. *Roth. germ. I.* 210. *II.* 538.

Philadelphus. *Hort. cliff.* 188. *Hort. upſ.* 122.

Syringa alba ſ. Philadelphus athenaei. *Bauh pin.* 399.

Frutex coronarius. *Cluſ. hiſt.* 1. *p.* 55.

β. Philadelphus nanus. *Mill. dict. n.* 2. *Du Roi. harbk.* 2. *p.* 6.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 641.

Wohlriechender Pfeifenſtrauch. W.

Habitat in Europa *auſtrali.* ♄ (v. v.)

Flos *primarius quinquefidus calyce, corolla, piſtillo, capſula. Phil. bot.* 178. *Petala aeſtivantia obvoluta cum ſole. Gemmis caret.*

Stylus *in calyce florente apice 4-fido; in calyce ſeminifero uſque ad baſin 4-fidus. Moench. l. c.* Var. *β. flore eſt pleno, ſeu potius flore du-vel triplicato.* Du Roi. *l. c.* R.

2. PHILADELPHUS *inodorus.*
P. foliis integerrimis. *Mill. dict.* 3.
Philadelphus flore albo majore inodoro. *Cateſb. car.* 2. *p.* 84. *t.* 84.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 643.
Geruchloſer Pfeifenſtrauch. W.
Habitat in Carolina. ♄.

*968. LEPTOSPERMUM.

Smith. Act. ſoc. Lin. Lond. 3. *p.* 260.

Cal. 5-fidus ſemiſuperus Petala 5-unguiculata ſtaminibus longiora. *Stigma* capitatum. *Capſ.* 4-vel 5-locularis. *Semina* anguloſa.

1. LEPTOSPERMUM *ſcoparium.*
L. foliis ovatis mucronatis obſolete trinerviis, calycibus glabris, dentibus membranaceis coloratis. *Smith. ſoc. Lin. Lond.* 3. *p.* 262.
Melaleuca (*ſcoparia*) foliis alternis mucronatis trinerviis, floribus terminalibus ſolitariis ſeſſilibus. *Wendl. et Schrad. ſert. hannov. p.* 25. *t.* 15. *
Philadelphus (*ſcoparius*) foliis lanceolatis integerrimis rigidis trinerviis, floribus omnibus quinquefidis, lacinis calycinis coloratis deciduis. *Ait. Kew.* 2. *p.* 156.
α. linifolia foliis lanceolatis.
Leptoſpermum ſcoparium. *Forſt. gen.* 36. *n.* 6. *Coock. itin.* 1. *p.* 100. *t.* 22.
Melaleuca (*ſcoparia*) foliis alternis ovatis acutis ſubtrinerviis, floribus terminalibus ſolitariis ſeſſilibus. *Forſt. prodr. n.* 210.
Melaleuca (*ſcoparia*) foliis alternis ovatis acutis trinerviis punctatis, floribus terminalibus ſolitariis. *Lin. ſuppl.* 343.
Melaleuca ſcoparia *α.* dioſmatifolia. *Wendl. et Schrad. l. c. t.* 15. *f.* 1.
β. myrtifolia, foliis ovato-ellipticis.
Melaleuca ſcoparia *β.* myrtifolia. *Wendl. et Schrad. l. c. t.* 15. *f.* 2.

Philadelphus floribundus. *Uſteri. et Roemer. Mag.* 7. *Stuck. p.* 177. *t.* 2.
Leptoſpermum ſquarroſum. *Gaert. ſem.* 1. *p.* 174. *t.* 35.
Beſenartige Südſeemyrte. W.
Habitat in Nova Zeelandia. ♄ (v. v.)

*2. LEPTOSPERMUM *Thea.*

L. foliis lineari-lanceolatis ſubmucronatis trinerviis calycibus glabris, dentibus membranaceis coloratis. W.
Melaleuca (*Thea*) ramis pendulis, foliis alternis obovàto-lanceolatis trinerviis, floribus lateralibus ſolitariis ſubſeſſilibus. *Wendl. et Schrad. ſert. hannov. p.* 24. *t.* 14.*
Theegebende Südſeemyrte. W.
Habitat in Nova Hollandia. ♄ (v. v. ſ. fl.)
Affine ſequenti, ſed foliis *mucrone parvo inſtructis trinerviis diſtinctum; a L. attenuato: calycibus glabris* pedunculis*que unifloris diverſum.* W.

*3. LEPTOSPERMUM *flaveſcens.*

L. foliis lineari-lanceolatis obtuſis enerviis, calycibus glabris, dentibus membranaceis nudis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 262.
Gelbliche Südſeemyrte. W.
Habitat in Nova Hollandia. ♄

*4. LEPTOSPERMUM *attenuatum.*

L. foliis lanceolato-linearibus acutis trinerviis, calycibus ſericeo-villoſis, dentibus membranaceis coloratis nudiuſculis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 262.
Verdünnte Südſeemyrte. W.
Habitat in Nova Hollandia. ♄
Nunquam vivum vidi. Flores *videntur albi, ſunt plerumque bini in pedunculo brevi ſericeo.* Smith. l. c.

*5. LEPTOSPERMUM *lanigerum.*

L. foliis ovato-lanceolatis trinerviis, calycibus ſericeo-villoſis, dentibus foliaceis perſiſtentibus. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 263.
Leptoſpermum trinerve. *White' itin. p.* 229. *c. icone.*
Philadelphus (*laniger*) foliis oblongis acutis integerrimis pubeſcentibus, calycibus lanatis. *Ait. Kew.* 2. *p.* 156.
Wollige Südſeemyrte. W.
Habitat in Nova Hollandia. ♄ (v. ſ. ſ. fl.)
Variat foliis *glabris et pubeſcentibus*, calyce *ſericeo et villoſo, villis longis et craſſis.* Smith. l. c.

*6. LEPTOSPERMUM *pubeſcens.* †
L. foliis lanceolato-oblongis piloſis ſubobliquis apice reflexis.
Philadelphus laniger β. piliger. *Ait. Kew.* 2. *p.* 156.
Weichbehaarte Südſeemyrte. W.
Habitat in Nova Hollandia. ♄

*7. LEPTOSPERMUM *parvifolium.*
L. foliis obovatis enerviis, ramulis calycibusque piloſis, dentibus membranaceis coloratis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 263.
Kleinblättrige Südſeemyrte. W.
Habitat in Nova Hollandia.

*8. LEPTOSPERMUM *arachnoideum.*
L. foliis ſubulatis pungentibus, ramulis hirtis, calycibus dentibusque villoſis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 263.
Leptoſpermum arachnoides. *Gaert. ſem.* 1. *p.* 175. *t.* 35.
Spinnenwebenartige Südſeemyrte. W.
Habitat in Nova Hollandia. ♄

*9. LEPTOSPERMUM *juniperinum.*
L. foliis lineari-lanceolatis pungentibus, ramulis ſericeis, calycibus glabris, dentibus membranaceis coloratis nudis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 263.
Melaleuca (*tenuifolia*) frutex erectus ramoſus, ramis pendulis piloſis, foliis alternis acutis tenuibus obliquis mucronatis piloſis quinquenervoſis, floribus lateralibus ſolitariis ſubſeſſilibus. *Wendl. obſ.* 50.
Wachholderblättrige Südſeemyrte. W.
Habitat in Nova Hollandia. ♄ (v. ſ.)

*10. LEPTOSPERMUM *baccatum.*
L. foliis lineari-lanceolatis pungentibus, ramulis hirtis, calycibus glabris, dentibus membranaceis coloratis pubeſcentibus, capſula baccata. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 264.
Beerentragende Südſeemyrte. W.
Habitat in Nova Hollandia. ♄

*11. LEPTOSPERMUM *ambiguum.*
L. foliis linearibus apice recurvis, calycibus glabriuſculis, dentibus foliaceis lanceolatis nudis ſtaminibus corolla longioribus. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 264.
Zweifelhafte Südſeemyrte. W.
Habitat in Nova Hollandia. ♄

Ab omnibus a me examinatis ſpeciebus ſtaminibus *corolla longioribus diverſa, qua nata cum Metroſideris convenit, ſed differt ab his* ſtigmate *capitato charactere Leptoſpermis proprio ut et habitu.* Smith. l. c.

12. LEPTOSPERMUM *virgatum.* †
L. foliis oppoſitis lineari-lanceolatis enerviis, umbellis terminalibus.
Melaleuca virgata. *Lin. ſuppl.* 343.
Melaleuca (*virgata*) foliis oppoſitis linearibus enerviis, pedunculis axillaribus umbelliferis. *Forſt. prodr. n.* 211.
Leptoſpermum virgatum. *Forſt. gen.* 36. 7.
Myrtus amboinenſis. *Rumph. amb.* 2. *p.* 77. *t.* 18.
Ruthenförmige Südſeemyrte. W.
Habitat in Nova Caledonia. ♄

Folia *pagina inferiori punctis atris reſinoſis tecta. In binis ſpeciminibus hujus plantae, in herbario Linnaeano aſſervatis, vidi* ſtamina *decem* fructumque *uni-vel bilocularem, videtur itaque ſpecies Imbricariae cum duplici ſtaminum numero eſſe. Si vero inpoſterum fructus* 3-*ſ.* 4-*locularis obſervetur, eſſet ſpecies decandra Beckiae, qua cum* foliis *oppoſitis, ut et* magnitudine *habituque floris convenit, quum reliqua Leptoſperma foliis gaudeant alternis; hinc plantae hujus genus futuris obſervationibus eſt determinandum.* Smith. Act. ſoc. Lin. Lond 3. p. 265.

*969. FABRICIA.

Smith. Act. ſoc. Lin. Lond. 3. *p.* 265.

Cal. 5-fidus ſemiſuperus. *Petala* 5 ſeſſilia. *Stigma* capitatum. *Capſ.* multilocularis. *Semina* alata.

*1. FABRICIA *myrtifolia.*
F. foliis lanceolato-obovatis oppoſitis, dentibus calycinis orbiculatis. W
Fabricia myrtifolia. *Gaert. ſem.* 1. *p.* 175. *t.* 35. *f.* 4.
Myrtenblättrige Fabricie. W.
Habitat in Nova Hollandia. ♄

*2. FABRICIA *laevigata.*
F. foliis obovatis alternis glaucis, dentibus calycinis triangularibus. W.

Fabricia laevigata. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 265.
Glatte Fabricie. W.
Habitat in Nova Hollandia. ♄
Folia *ſunt alterna obovata glabra obſcure* 3-*ſ.* 5-*nervia glauca.* Calycis *dentes triangulares, qui orbiculati ſunt in F. myrtifolia, quae et omnibus partibus duplo major eſt.* Smith. l. c.

*970. METROSIDEROS.

Smith. Act. ſoc. Lin. Lond. 3. *p.* 266.

Cal. 5-fidus ſemiſuperus. *Petala* 5. *Stamina* longiſſima exſerta. *Stigma* ſimplex. *Capſula* 3-vel 4-locularis.

* *Foliis oppoſitis.*

*1. METROSIDEROS *hiſpida.*
M. foliis oppoſitis baſi cordatis amplexicaulibus, ramulis pedunculis calycibusque hiſpidis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 267.
Rauhes Eiſenmaſs. W.
Habitat in Nova Hollandia. ♄
Planta *ſpecioſa, foliis latis ſeſſilibus oppoſitis, ramiſque hiſpidis a reliquis diſtincta.* Flores *flavi, ſtaminibus divaricatis, umbellati,* umbellis *in corymbum magnum terminalem collectis,* pedunculis *hiſpidis pilis rubro-fuſcis tectis ut in Robinia hiſpida.* Smith. l. c.

*2. METROSIDEROS *floribunda.*
M. foliis oppoſitis petiolatis ovato-lanceolatis, panicula brachiata, pedicellis umbellatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 267.
Vielblütiges Eiſenmaſs. W.
Habitat in Nova Hollandia. ♄
Flores *minores quem in praecedente, videntur albi.* Panicula *e ramis decuſſatis, umbella parva terminatis, compoſita.* Pedunculi *hiſpidi interdum glabri.* Smith. l. c.

*3. METROSIDEROS *coſtata.*
M. foliis oppoſitis petiolatis lineari-lanceolatis acuminatis obliquis, panicula brachiato-decompoſita, pedicellis ſubumbellatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 267.

Metroſideros coſtata. *Gaert. ſem.* 1. *p.* 171. *t.* 34. *f.* 2.
Ribbenfrüchtiges Eiſenmaſs. W.
Habitat in Nova Hollandia. ♄
Diſtincta a praecedente: foliis *anguſtioribus longioribus, magis rigidis et nitidis, obliquis vel falcatis*, panicula *irregulari, duplicato-ramoſa*, ramis *terminalibus ſubumbellatis*, floribus *majoribus flaveſcenti-albis*. Smith. l. c.

4. METROSIDEROS *diffuſa*.
M. foliis oppoſitis ovatis venoſis utrinque glabris, paniculis axillaribus terminalibusve, pedicellis oppoſitis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 268.
Melaleuca (*diffuſa*) foliis oppoſitis ovatis acuminatis, pedunculis rameis cymoſis. *Forſt. prodr. n.* 213.
Melaleuca (*lucida*) glabra, foliis oppoſitis ovatis venoſis, paniculis terminalibus, pedicellis oppoſitis bifloris. *Lin. ſuppl.* 342.
Ausgebreitetes Eiſenmaſs. W.
Habitat in Nova Zeelandia. ♄
Folia *pagina inferiori trinervia, nervis ante apicem evaneſcentibus*. Flores *lutei*

5. METROSIDEROS *villoſa*.
M. foliis oppoſitis ovatis venoſis ſubtus pubeſcentibus, thyrſis axillaribus terminalibusve oppoſitis villoſis, floribus ſeſſilibus confertis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 268.
Melaleuca (*villoſa*) tomentoſa, foliis oppoſitis ovatis venoſis, paniculis terminalibus, pedicellis bi-ſ. trifloris. *Lin. ſuppl.* 342.
Melaleuca (*aeſtuoſa*) foliis oppoſitis ovatis obovatisque venoſis (ſubtrinerviis), pedunculis cymoſis terminalibus tomentoſis. *Forſt. prodr. n.* 215.
Leptoſpermum collinum. *Forſt. gen.* 36. *n.* 2.
Metroſideros ſpectabilis. *Gaert. ſem.* 1. *p.* 172. *t.* 34. *f.* 9.?
Haariges Eiſenmaſs. W.
Habitat in Taheiti. ♄
Simillima M. diffuſae, at illa glaberrima, floribus *luteis; haec vero caule*, petiolis, panicula, pedunculis *et* calyce *tomento albis;* floribus *rubicundis*, ſtaminibus *multo majoribus*.

*6. METROSIDEROS *florida*.
M. foliis oppoſitis obovato-oblongis venoſis glabris, thyrſo terminali, calycibus turbinatis nudis, *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 269.

Melaleuca (*florida*) foliis oppoſitis oblongo-obovatis venoſis, pedunculis terminalibus cymoſis, calycibus turbinatis. *Forſt. prodr. n.* 114.
Leptoſpermum ſcandens. *Forſt. gen.* 36. *n.* 1.
Dichtblütiges Eiſenmaſs. W.
Habitat in Nova Zeelandia. ♄
Rami *longi thyrſo magno terminati.* Flores *flaveſcentes.* Calyx *valde elongatus forma Caryophyllorum.* Folia *glabra.* Pedunculi *et* Calyces *ſericei, pilis adpreſſis tecti.* Smith. l. c.

*7. METROSIDEROS *glomulifera.*

M. foliis oppoſitis ovatis reticulato-venoſis ſubtus pubeſcentibus, capitulis lateralibus pedunculatis bracteisque tomentoſis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 269.
Kugelblütiges Eiſenmaſs. W.
Habitat in Nova Hollandia. ♄
Arbor *ramis teretibus oppoſitis.* Folia *oppoſita brevi petiolo pubeſcenti ſuffulta, ovata, integerrima ſubundulata, venis numeroſis reticulata, ſubtus pubeſcentia.* Flores *vireſcenti-flavi, in capitulo parvo globoſo collecti, pedunculo unciali ſimplici pubeſcenti ſuffulti.* Pedunculi *laterales ſupra inſertionem petioli foliorum ſummorum oppoſiti.* Capitulum *florum binis oblongis pubeſcentibus bracteis ſuffultum.* Smith. l. c.

8. METROSIDEROS *anguſtifolia.*

M. foliis oppoſitis lineari-lanceolatis nudis, pedunculis axillaribus umbellatis, bracteis lanceolatis glabris. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 270.
Myrtus (*anguſtifolia*) pedunculis umbellatis, foliis lineari-lanceolatis ſubſeſſilibus. *Mant.* 74.
Phillyrea foliis longis ſubtus flavis, fructu deciduo. *Burm. afr.* 237. *t.* 83. *f.* 2.?
Houttuyn Lin. Pfl. Syſt. 3. *p.* 647. *t.* 25. *f.* 2.
Schmalblättriges Eiſenmaſs. W.
Habitat ad Cap. b. ſpei. ♄ (v. ſ.)
Arbuſcula *ramis teretibus, ramulis tetragonis glabris.* Folia (*Phillyreae anguſtifoliae*) *ſubſeſſilia longa lineari-lanceolata integerrima glabra, internodiis multoties longiora.* Pedunculi *laterales oppoſiti breves internodiis vix longiores apice* Umbellula *ſimplici, rarius compoſita parva multiflora.* Petala *ſubrotunda.* Stamina *multa erecta ſetacea petalis longiora.*

Eſt vera Metroſideri ſpecies, fructus eſt capſula nec bacca, ſimillimus capſulae Melaleucae ſuaveolentis. Gaertneri cui planta nullo modo ſimilis. Burmanni ſynonymum vix huc pertinere videtur, quum fructum, baccam unilocularem monoſpermam eſſe, dixerit. Smith. l. c.

** *Foliis alternis.*

*9. METROSIDEROS *ciliata.*

M. foliis ſparſis ſuboppoſitis ellipticis obtuſis coriaceis baſi ſubciliatis, corymbis terminalibus piloſis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 271.

Melaleuca (*ciliata*) foliis ſparſis, ſuperioribus ſuboppoſitis oblongis obtuſis ciliatis aveniis enerviis, cymis corymboſis terminalibus. *Forſt. prodr. n.* 217.

Leptoſpermum ciliatum. *Forſt. gen.* 36. *n.* 3.

Gefranztes Eiſenmaſs. W.

Habitat in Nova Caledonia. ♄

Folia *rigida craſſa concava, margine revoluta, iis Celaſtri buxifolii ſimilia, ſed minus nitida, inferne pallida, a venis transverſalibus reticulata, cum coſta media tenui. Clariſſ. Forſter* folia *avenia et enervia deſcripſit, venae forte in recenti planta ſunt minus viſibiles.* Foliorum *baſis*, rami *juniores*, pedunculi, calyx *et* petala *plana, pilis longis patentibus obſita.* Flores *pulchri magni, ſaturate rubri, pauci in corymbo vel umbella terminali.* Fructus *magnus depreſſus trilobus margine calycis altior.* Smith. l. c.

*10. METROSIDEROS *linearis.*

M. foliis ſparſis linearibus canaliculatis acutis rigentibus, floribus lateralibus confertis ſeſſilibus. *Smith. Act. ſoc. Lin. Lond.* 2. *p.* 273. *

Melaleuca (*linearis*) foliis linearibus carinatis, junioribus villoſis, floribus lateralibus ſeſſilibus confertis. *Wendl. et Schrad. ſert. hannov.* 19. *t.* 11. *

Linienblättriges Eiſenmaſs. W.

Habitat in Nova Hollandia. ♄ (v. ſ.)

*11. METROSIDEROS *lanceolata.*

M. foliis alternis lanceolatis mucronatis, floribus lateralibus confertis ſeſſilibus pubeſcentibus. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 272. *

Metroſideros citrina. *Curt. mag.* 260.

Lanzenblättriges Eiſenmaſs. W.

Habitat in Nova Hollandia. ♄ (v. ſ.)

*12 METROSIDEROS *ſaligna.*

M. foliis alternis lanceolatis utrinque attenuatis mucronatis, floribus lateralibus confertis ſeſſilibus glabris. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 272.

Weidenblättriges Eiſenmaſs. W.

Habitat in Nova Hollandia. ♄ (v. ſ. ſ. fl.)

A praecedente diſtinta: foliis *minus rigidis attenuatis*, floribus *minoribus flaveſcentibus*, calyce petalisque *glabris*, *neque pubeſcentibus*, *neque ciliatis*. Smith. l. c.

*13. METROSIDEROS *viminalis.*

M. foliis alternis lineari-lanceolatis, floribus confertis lateralibus pubeſcentibus.

Metroſideros viminalis. *Gaert. ſem.* 1. *p.* 171. *t.* 34. *f.* 4.

Schlankes Eiſenmaſs. W.

Habitat in Nova Hollandia. ♄

Differt a praecedente: foliis *lineari-lanceolatis non attenuatis* floribus*que pubeſcentibus*. Smith. Act. ſoc. Lin. Lond. 3. p. 273.

*14 METROSIDEROS *capitata.*

M. foliis ſparſis obovatis mucronulatis, capitulis terminalibus, calycibus ramulisque piloſis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 273.

Kopfförmiges Eiſenmaſs. W.

Habitat in Nova Hollandia. ♄

Ab omnibus ſpecies, valde diſtincta. Folia *vix quadrilinearia parce venoſa, obſolete crenata vel potius ſcabra e punctis minutis marginalibus.* Flores *capitati in pedunculo brevi.* Calyx *tubuloſus hirſutus, dentibus foliaceis perſiſtentibus.* Petala *parva purpurea.* Stamina *purpurea corolla triplo longiora.* Germen *minutum in fundo calycis.* Stylus *longitudine ſtaminum.* Stigma *parum incraſſatum nec capitatum.* Fructum *maturum non vidi, ſed non dubito fore Metroſideri ſpeciem, quum rudimenta trium valvularum detexerim.* Florum *forma eſt ut in Lythris.* Folia *ſunt punctata parum aromatica.* Smith. l. c.

Leptoſpermi et Metroſideri ſpecies a Clariſ. Salisbury (in prodromo ſtripium in horto ad chapel allerton vagentium) indicatae, ob nimis brevem descriptionem heic omiſi. W.

971. PSIDIUM.

Gen. plant. ed. Schreb. n. 841.

Cal. 5-fidus, ſuperus. *Petala* 5. *Bacca* 1-locularis, polyſperma.

1. PSIDIUM *pyriferum.*
P. foliis ellipticis ſubtus pubeſcentibus, pedunculis unifloris. W.
P. foliis lineatis obtuſiuſculis unifloris. *Mill. dict.* 1. *Jacq. obſ.* 2. *p.* 6.
P. caule quadrangulo. *Hort. cliff.* 148. *Fl. zeyl.* 192.
Guajava foliorum angulis quadrangulis, fructu oblongo. *Trew. ehr. t.* 43.
Guajava alba dulcis. *Comm. hort.* 1. *p.* 121. *t.* 63. *Mer. ſurin.* 19. *t.* 19.
Guajavos fructu pallido dulci. *Burm. zeyl.* 112.
Cujavus domeſtica. *Rumph. amb.* 1. *p.* 140. *t.* 47.
Pela. *Rheed. mal.* 3. *p.* 31. *t.* 34.
Pſidium fruticoſum, foliis ovatis venoſis, fructu majore. *Browm. jam.* 238.
Malo puniceo affinis pomifera, flore pentapetalo albo, fructu ex toto eſculento majore albo. *Sloan. hiſt.* 2. *p.* 163. Guajava *vulgo.*
Houttuyn Lin. Pfl. Syſt. 1. *p.* 699.
Birnentragender Gujavabaum. W.
Habitat in Indiis. ♄ (v. ſ.)

*2. PSIDIUM *pumilum.*
P. foliis lanceolatis acutis ſubtus tomentoſis, pedunculis unifloris. W.
P. (*pumilum*) foliis lanceolatis lineatis acutis, pedunculis unifloris, fructu globoſo. *Vahl. ſymb.* 2. *p.* 56. *
P. (*anguſtifolium*) foliis lanceolatis lineatis ſubtus tomentoſis, pedunculis unifloris. *Lamarck. encycl.* 3. *p.* 16. *
Cujavillus. *Rumph. amb.* 1. *p.* 145. *t.* 49.
Niedriger Gujavabaum. W.
Habitat in Zeylona. ♄

*3. PSIDIUM *aromaticum.*
P. foliis oblongis acuminatis glabris, pedunculis unifloris. W.
P. (*aromaticum*) foliis ovatis acuminatis, fructu dulci ceraſiformi. *Aubl. guj.* 1. *p.* 485. *t.* 191. *

Gewürzhafter Gujavabaum. W.
Habitat in ſylvis Cajennae, Gujanae. ♄

*4. PSIDIUM *grandiflorum.*

P. foliis ovatis acuminatis glabris, pedunculis unifloris bracteatis. W.

P. (*grandiflorum*) foliis ovatis acuminatis, fructu parvo quadriloculari. *Aubl. guj.* 1. *p.* 483. *t.* 190. *

Groſsblütiger Gujavabaum. W.

Habitat in ſylvis Cajennae. ♄

Pedunculi *longi uniflori.* Bracteae *binae lanceolatae oppoſitae ſub flore.* W.

5. PSIDIUM *Decaſpermum.*

P. foliis ovatis acuminatis pubeſcentibus, pedunculis unifloris bracteatis. *Forſt. prodr. n.* 219.

P. foliis ovatis acuminatis planis, pedunculis unifloris bracteatis. *Lin. ſuppl.* 252.

Decaſpermum fruticoſum. *Forſt. gen. n.* 37.

Ungenieſsbarer Gujavabaum. W.

Habitat in Societatis *inſulis.* ♄

Frutex *glaber.* Folia *petiolata parva integerrima venoſa non lineata.* Pedunculi *axillares foliis ſesqui breviores bracteati ſericei uniflori.* Bractea *una ſ. altera oblonga.* Flores *parvi albi.* Calyces *ſericei.* Capſula *ſ.* Bacca *parva globoſa toruloſa, calyce perſiſtente coronata.*

6. PSIDIUM *pomiferum.*

P. foliis oblongo-lanceolatis ſubtus pubeſcentibus, pedunculis trifloris. W.

P. foliis lineatis acuminatis, pedunculis trifloris, *Mill. dict. n.* 2. *Iacq. obſ.* 2. *p.* 7.

Guajava alba acida, fructu rotundiore. *Pluk. alm.* 181. *t.* 193. *f.* 4.

Guajaba pomifera indica, pomis rotundis. *Bauh. pin.* 437.

Cujavus agreſtis. *Rumph. amb.* 1. *p.* 142. *t.* 48. *Mer. ſurin.* 57. *t.* 57.

Malacca pela. *Rheed. mal.* 3. *p.* 33. *t.* 35. *Burm. ind. p.* 113.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 703.

Apfeltragender Gujavabaum. W.

Habitat in Indiis. ♄

*7. PSIDIUM *guineenſe.*

P. foliis ovatis integerrimis ſubtus tomentoſis, peduncu-

lis trifloris. *Swartz prodr.* 77. *Fl. ind. occid.* 2. *p.* 881. *

Africaniſcher Gujavabaum. W.

Habitat in inſula Principis *prope* Guineam. ♄

*8. PSIDIUM *montanum.*

P. foliis oblongis acuminatis crenulatis nitidis, pedunculis multifloris. *Swartz prodr.* 77. *Fl. ind. occid.* 2. *p.* 879. *

P. arboreum maximum, foliis ovatis nitidis, ligno fuſco, fibris undulatis. *Brown. jam.* 238.

Berg-Gujavabaum W.

Habitat in montibus Jamaicae. ♄

972. EUGENIA.

Gen. plant. edit. Schreb. n. 842.

Cal. 4-partitus, ſuperus. *Petala* 4. *Bacca* 1-locularis 1-ſperma.

1. EUGENIA *malaccenſis.*

E. foliis integerrimis oblongis, pedunculis ſubquadrifloris lateralibus. W.

E. foliis integerrimis, pedunculis ramoſis lateralibus. *Fl. zeyl.* 187. *Mill. dict. n.* 1.

Perſici oſſiculo fructus malaccenſis rubens. *Bauh. pin.* 441.

Jamboſa domeſtica. *Rumph. amb.* 1. *p.* 121. *t.* 37, 38.

Nati-ſchanibu. *Rheed. mal.* 1. *p.* 29. t. 18. *Raj. hiſt.* 1748. *Burm. ind.* 114.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 705.

Malacciſcher Jambuſenbaum. W.

Habitat in India. ♄

2. EUGENIA *Jambos.*

E. foliis integerrimis lanceolatis, pedunculis ſubquadrifloris terminalibus. W.

E. foliis integerrimis, pedunculis ramoſis terminalibus. *Fl. zeyl.* 188. *. *Mill. dict. n.* 2.

Perſici oſſiculo fructus malaccenſis ex candido rubeſcens. *Bauh. pin.* 441.

Jamboſa ſylveſtris alba. *Rumph. amb.* 1. *p.* 127. *t.* 39.

Malacca ſchambu. *Rheed. mal.* 1. *p.* 27. *t.* 17. *Raj. hiſt.* 1478. *Burm. ind.* 114.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 708.

Gemeiner Jambuſenbaum. W.
Habitat in India. ♄ (v. ſ.)

*3. EUGENIA *baruenſis.*

E. foliis integerrimis ovato-lanceolatis, pedunculis multifloris axillaribus petiolo brevioribus. W.

E. (*baruenſis*) foliis integerrimis ovato-lanceolatis, pedunculis axillaribus unifloris et aggregatis vel multifloris. *Iacq. ic. rar.* 3. *t.* 486. *Collect.* 3. *p.* 183.*

E. pedunculis partialibus elongatis. *Iacq. amer.* 153.

Baruſcher Jambuſenbaum. W.

Habitat in inſula Baru *prope* Carthagenam *in* America. ♄ (v. ſ.)

Pedunculi *ſunt uniflori ſolitarii vel aggregati, ſaepius vero multiflori fere racemoſi, ſemper vero petiolo breviores.* Calyx *ſic dictus inferus, vel potius bracteae, non adſunt.* W.

*4. EUGENIA *floribunda.*

E. foliis integerrimis margine revolutis oblongis acuminatis, pedunculis multifloris axillaribus petiolorum longitudine. W.

Eugenia floribunda. *Weſt. ſt. cruc. p.* 216.

Vielblütiger Jambuſenbaum. W.

Habitat in inſula St. Crucis. ♄ (v. ſ.)

Folia *parva uncialia vel ſesquiuncialia anguſta elliptica baſi anguſtiora, longiſſime acuminata integerrima, margine revoluta, tenuiſſime venoſa, inferne punctata.* Pedunculi *ſolitarii, aggregati vel multiflori, longitudine petiolorum, baſi bracteati.* Flores *parvi.* W.

*5. EUGENIA *buxifolia.*

E. foliis integerrimis obovato-lanceolatis obtuſis, pedunculis multifloris axillaribus lateralibusque petiolorum longitudine. W.

Myrtus (*buxifolia*) racemulis breviſſimis confertis axillaribus, foliis cuneatis oblongis obtuſis convexiuſculis. *Swartz. prodr.* 78. *Fl. ind. occid.* 2. *p.* 899.*

Buchsbaumblättriges Jambuſenbaum. W.

Habitat in Hiſpaniolae *ſteriliſſimis campis petroſis verſus mare.* ♄ (v. ſ.)

*6. EUGENIA *axillaris.*

E. foliis integerrimis oblongis acuminatis obtuſis planis, pedunculis axillaribus multifloris petiolorum longitudine. W.

Myrtus (*axillaris*) racemulis multifloris confertis axilla-

ribus breviſſimis, foliis oblongis acuminatis nitidis. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 901. *

Winkelblütiger Jambuſenbaum. W.

Habitat in montoſis Jamaicae. ♄

Differt a Myrto monticola, cui ſtatura proxima: magnitudine, ſoliis *majoribus magis acuminatis*, racemulis *confertis*, floribus *copioſioribus*, calyce *duplici*. Swartz l. c.

*7. EUGENIA *ſinemarienſis*. †

E. foliis integerrimis ovato-oblongis acuminatis reticulato-venoſis, pedunculis axillaribus terminalibusque multifloris petiolorum longitudine. W.

E. foliis ovatis acuminatis, floribus ſeſſilibus congeſtis terminalibus et axillaribus, fructu rubro triſpermo. *Aubl. guj.* 1. *p.* 501. *t.* 198.

Sinemariſcher Jambuſenbaum. W.

Habitat in Gujana *ad ripas fluvii* Sinemarienſis. ♄

An Myrtis adnumeranda? W.

*8. EUGENIA *lateriflora*.

E. foliis integerrimis margine revolutis ſubrotundo-ellipticis reticulato-venoſis, floribus ſparſis lateralibus ſeſſilibus. W.

E. floribus lateralibus ſeſſilibus, foliis oblongis integerrimis nitidis ſubtus punctatis. *Vahl. ſymb.* 3. *q.* 64. *

Seitenblütiger Jambuſenbaum. W.

Habitat in inſula St. Crucis. ♄ (v. ſ.)

*9. EUGENIA *crenulata*.

E. foliis repandis ſubrotundis, pedunculis axillaribus ſubtrifloris petiolorum longitudine. W.

Myrtus (*crenulata*) pedunculis ſolitariis axillaribus ſubtrifloris, foliis ſubrotundis crenatis. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 889 *

Gekerbter Jambuſenbaum. W.

Habitat in Hiſpaniolae *fruticetis montoſis*. ♄ (v. ſ.)

*10. EUGENIA *alpina*.

E. foliis integerrimis ellipticis aveniis, pedunculis ſolitariis ſubterminalibus unifloris breviſſimis. W.

Myrtus (*alpina*) pedunculis ſolitariis terminalibus unifloris breviſſimis, foliis ovatis, ramulis quaternis faſtigiatis, caule arboreo. *Swartz prodr.* 77. *Fl. ind. occid.* 2. *p.* 883 *

Gebirgs Jambuſenbaum. W.

Habitat in Jamaicae *cacumine montium coerulcorum*. ♄ (v. ſ.)

Folia *parva margine revoluta, parum majora quam in Vaccinio oxycocco*. W.

*11. EUGENIA *liguſtrina*.

E. foliis integerrimis lanceolatis baſi anguſtatis obtuſis aveniis, pedunculis unifloris ſolitariis terminalibus. W.

Myrtus (*liguſtrina*) pedunculis ſolitariis ſubſimplicibus terminalibus unifloris, foliis lanceolatis obtuſis convexis lucidis. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 885.*

Liguſterartiger Jambuſenbaum. W.

Habitat in fruticetis depreſſis montium Hiſpaniolae. ♄ (v. ſ.)

12. EUGENIA *Pſeudo-pſidium*.

E. foliis integerrimis ovatis acuminatis, pedunculis unifloris terminalibus lateralibusque pluribus. W.

E. foliis integerrimis, pedunculis unifloris pluribus lateralibus terminalibusque. *Syſt. veg.* 384.

E. foliis integerrimis, pedunculis unifloris axillaribus et terminalibus. *Jacq. amer.* 152. *t.* 93.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 709.

Gujavaartiger Jambuſenbaum. W.

Habitat in Martinica. ♄ (v. ſ.)

Fructus *laevis*.

13. EUGENIA *uniflora*.

E. foliis integerrimis ovato-lanceolatis, pedunculis unifloris ſolitariis lateralibus. W.

E. (*uniflora*) foliis integerrimis cordato-lanceolatis, pedunculis unifloris lateralibus. *Sp. pl.* 673.

Myrtus (*braſiliana*) floribus ſolitariis pedunculis nudis, petalis ſubciliatis. *Sp. pl.* 674.

Plinia (*rubra*) floribus tetrapetalis. *Mant.* 402.

Plinia (*pedunculata*) floribus pedunculatis polyandris. *Lin. ſuppl.* 253.

Eugenia. *Roy. lugdb.* 265.

E. indica, myrti folio deciduo, flore albo, fructu ſuave rubenti molli leviter ſulcato et odoro. *Mich. gen.* 226. *t.* 108.

Myrtus indica, foliis rigeſcentibus latis ac recurvis parum odoratis. *Till. piſ.* 117. *t.* 44.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 710. 714. 715.

Einblütiger Jambuſenbaum. W.

Habitat in Braſilia. ♄ (v. ſ.)

Pedunculi *ſolitarii*.

*14. EUGENIA *Patriſii.*

E. foliis integerrimis ovatis acuminatis acutis, pedunculis unifloris geminatis axillaribus, bracteis ſetaceis. W.
E. (*Patriſii*) foliis ellipticis glabris acuminatis acutis, pedunculis unifloris capillaribus axillaribus geminis, terminalibus ſubquaternis. *Vahl. eclog.* 2. *p.* 35. *
Patriſiſcher Jambuſenbaum. W.
Habitat in America *meridionali.* ♄
Affinis E. uniflorae, diverſa habitu *et* bracteis *ſetaceis nec ovatis.* Vahl. l. c.

*15. EUGENIA *zeylanica.* W.

E. foliis integerrimis oblongis acuminatis coriaceis impunctatis, pedunculis unifloris filiformibus ſubgeminatis axillaribus lateralibusque. W.
Zeyloniſcher Jambuſenbaum. W.
Habitat in Zeylona. ♄ (v. ſ.)
Folia *oppoſita breve petiolata oblonga baſi anguſtiora apice acuminata obtuſa, coriacea nitida venoſa non punctata.* Pedunculi *uniflori filiformes geminati vel ſolitarii in foliorum axillis vel in ramulis aphyllis.* Bracteae *binae breviſſimae ſubulatae ſub germine. Differt a binis praecedentibus* foliis *coriaceis nec membranaceis*, defectu *punctorum*, bracteis *breviſſimis.* W.

16. EUGENIA *cotinifolia.*

E. foliis integerrimis ſubrotundo-obovatis obtuſis, pedunculis unifloris ſubgeminatis lateralibus. W.
E. foliis ovatis obtuſis integerrimis, pedunculis unifloris. *Mant.* 243. *Iacq obſ* 3. *p.* 3. *t.* 53. †
Houttuyn Lin. Pfl. Syſt. 1. *p.* 710.
Parückenbaumblättriger Jambuſenbaum. W.
Habitat in Cajenna. ♄
Rami *juniores angulati.* Folia *oppoſita, ſubpetiolata, nitida, coriacea.* Pedunculi *axillares*, 1, 2, *ſ.* 3.

*17. EUGENIA *latifolia.*

E. foliis integerrimis ovato-oblongis acuminatis reticulato-venoſis, pedunculis unifloris ſubternis axillaribus, fructiferis nutantibus. W.
E. foliis ampliſſimis ovatis acutis, fructu nigro olivaeformi. *Aubl. guj.* 1. *p.* 502. *t.* 199. *
Streifblättriger Jambuſenbaum. W.
Habitat ad agrorum margines Gujanae. ♄

*18. EUGENIA *undulata.*

E. foliis integerrimis ovato-lanceolatis acuminatis margine ſubundulatis, pedunculis unifloris aggregatis ſupraxillaribus. W.

E. foliis ovato-lanceolatis acuminatis glabris ſubundulatis, pedunculis lateralibus unifloris aggregatis. *Vahl. eclog.* 2. *p.* 36. *

C. foliis ovato-oblongis acuminatis, fructu parvo rubro olivaeformi. *Aubl. guj.* 1. *p.* 508. *t.* 202. *

Wellenblättriger Jambuſenbaum. W.

Habitat in Cajenna, Gujana. ♄

*19. EUGENIA *Mini.*

E. foliis integerrimis oblongo-lanceolatis acuminatis, pedunculis axillaribus multifloris ſubracemoſis folio brevioribus. W.

E. (*Mini*) foliis lanceolatis, floribus racemoſis, fructu parvo albo ex rubro variegato. *Aubl. guj.* 1. *p.* 498. *t.* 197. *

Weidenblättriger Jambuſenbaum. W.

Habitat in ſylvis Cajennae *et* Gujanae. ♄

*20. EUGENIA *fragrans.*

E. foliis integerrimis ſubrotundo-ovatis obtuſiuſculis, pedunculis axillaribus multifloris trichotomis longitudine foliorum. W.

Myrtus *fragrans*) pedunculis axillaribus trichotomis ſimplicibusque, foliis ovatis convexiuſculis ſubcoriaceis punctatis glabris, caule arboreo. *Swartz prodr.* 79. *Fl. ind. occid.* 1. *p.* 914. * *Vahl. ſymb.* 2. *p.* 59. *

E. (*paniculata*) foliis integerrimis ovatis obtuſis, pedunculis paniculatis axillaribus et terminalibus. *Iacq. Collect.* 2. *p.* 108. *t. f.* 1.?

E. (*montana*) foliis ovatis acutis, floribus terminalibus, fructu parvo albo, punctis rubris notato. *Aubl. guj.* 1. *p.* 495. *t.* 195.?

Wohlriechender Jambuſenbaum. W.

Habitat in montibus excelſis Jamaicae *auſtralis.* ♄

*21. EUGENIA *punctata.*

E. foliis integerrimis oblongis obtuſis glabris, tenellis ſericeis, pedunculis trifloris axillaribus longitudine foliorum. W.

E. (*punctata*) foliis oblongis utrinque punctatis, pedunculis oppoſitis trifloris longitudine foliorum. *Vahl. ſymb.* 3. *p.* 65. *

Punctirter Jambuſenbaum. W.
Habitat in inſula St. Crucis. ♄ (v. ſ.)

*22. EUGENIA *Cumete.*
E. foliis integerrimis ovatis acuminatis reticulato-venoſis, pedunculis trichotomis axillaribus folio longioribus, floribus pentapetalis. W.
E. (*Cumete*) foliis latis ovatis acuminatis, floribus racemoſis. fructu albo. *Aubl. guj.* 1. *p.* 479. *t.* 196.*
Netzblättriger Jambuſenbaum. W.
Habitat in ſylvis remotioribus Gujanae. ♄ (v. ſ. ſ. fl.)

*23. EUGENIA *gujanenſis.*
E. foliis integerrimis oblongis acutiuſculis, pedunculis trifloris folio brevioribus. W.
E. foliis ovatis obtuſiuſculis ſubpetiolatis, pedunculis ſolitariis bi- et trifloris, fructu monoſpermo. *Aubl. guj.* 1. *p.* 506. *t.* 201.*
Rothfrüchtiger Jambuſenbaum. W.
Habitat in Gujanae *ſylvis prope montem* Serpent. ♄
Similis E. punctatae, diſtincta tamen: foliis *minoribus acutioribus,* venis *foliorum crebrioribus,* pedunculis *brevioribus,* floribus *minoribus.* W.

24. EUGENIA *caryophyllata.*
E. foliis integerrimis oblongis acutiuſculis, pedunculis trichotomis, paniculis axillaribus et terminalibus, calycibus repandis, fructu elliptico. W.
E. (*caryophyllata*) foliis integris, panicula trichotoma decompoſita. *Thunb. diſſ. de. Caryoph. aromat. p.* 1.*
Caryophyllus aromaticus. *Sp. pl.* 735. *Hort. cliff.* 207. *Mat. med. p.* 136. *Blackw. t.* 338. *Gaert. ſem. p.* 167. *t.* 33. *f.* 2.
Caryophyllus foliis ovato-lanceolatis oppoſitis, floribus terminalibus, ſtaminibus corolla longioribus. *Mill. dict. n.* 1.
Caryophyllus aromaticus fructu oblongo. *Bauh. pin.* 410.
Caryophyllus aromaticus indiae orientalis, fructu clavato monopyreno. *Pluk. alm.* 88. *t.* 155. *f.* 1.
Caryophyllus. *Cluſ. exot.* 16. *Rumph. amb.* 2. *p.* 3. *t.* 1. 2. 3.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 50. *t.* 12. *f.* 1.
Gewürznäglein-Jambuſenbaum. W.
Habitat in Moluccis, *ſolo aridiſſimo.* ♄ (v. ſ.)
Fructus *eſt* Drupa *ſecundum Illuſt. Thunbergii deſcriptionem, Gaertnerus vero* Baccam *monoſpermam ſe-*

mine accoco eſſe contendit. Fructus *nonnulli a me examinati erant*, baccae *monoſpermae*, *nec* drupae. W.

*25. EUGENIA *elliptica.*

E. foliis integerrimis ellipticis acuminatis, pedunculis paniculatis axillaribus et terminalibus, calyce repando, fructibus globoſis. W.

E. (*elliptica*) foliis ellipticis acuminatis, floribus paniculatis, calyce repando, bacca globoſa. *Smith. Act. ſoc. Lin. Lond.* 3 *p.* 281. *

Elliptiſcher Jambuſenbaum. W.

Habitat in Nova Hollandia. ♄

Calyx *clavatus margine repando, nec dentato.* Petala 4 *alba parva decidua.* Stylus *brevis ſtigmate ſimplici.* Bacca *magnitudine Piſi majoris, alba, calyce coronata monſperma.* Smith. l. c.

*26. EUGENIA *tomentoſa.*

E. foliis oblongis integerrimis ramisque villoſis, racemis axillaribus compoſitis. *Vahl. eclog.* 2. *p.* 37. *

E. foliis ovatis ſubſeſſilibus, floribus racemoſis. *Aubl. guj.* 1. *p.* 504. *t.* 200. *

Filziger Jambuſenbaum. W.

Habitat in Gujana. ♄

*27. EUGENIA *trinervia.*

E. foliis integerrimis ovatis trinerviis glabris, racemis axillaribus oppoſitis ſolitariis, infrafoliaceis geminatis. *Vahl. eclog.* 2 *p.* 36. *

Dreynerviger Jambuſenbaum. W.

Habitat in Cajenna. ♄

28. EUGENIA *acutangula.*

E. foliis crenatis, pedunculis terminalibus, pomis oblongis acutangulis. *Sp. pl.* 673. *Fl. zeyl.* 190. *

Butonica terreſtris rubra. *Rumph. amb.* 3. *p.* 181. *t.* 115.

Tſieria-ſamſtravadi. *Rheed. mal.* 4. *p.* 15. *t.* 7. *Raj. hiſt.* 140. *Burm. ind.* 114.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 711.

Eckenfrüchtiger Jambuſenbaum. W.

Habitat in India. ♄ (v. ſ.)

29. EUGENIA *racemoſa.*

E. foliis crenatis, racemis longiſſimis, pomis ovatis quadrangulis. *Sp. pl.* 673. *Fl. zeyl.* 191. *

Butonica ſylveſtris alba. *Rumph. amb.* 3. *p.* 181. *t.* 116.

Samſtravadi ſ. Caipatſiambu. *Rheed. mal.* 4. *p.* 11. *t.* 6. *Raj. hiſt.* 1479. *Burm. ind.* 115.

Houttuyn Lin. Pſl. Syſt. 1. *p.* 712.
Langtraubiger Jambuſenbaum. W.
Habitat in India. ♄ (v. ſ.)

30. EUGENIA *laurina.* W. †
E. foliis ſerrulatis oblongis glabris, racemis axillaribus conjugatis. W.
Myrtus (*laurinus*) pedunculis racemoſis bracteatis, foliis ovato-lanceolatis ſerrulatis glabris. *Retz. obſ.* 4. *p.* 26.
Laurus ſerrata floribus ſpicatis, ex foliorum alis provenientibus. *Burm. zeyl. p.* 139. *t.* 62. *
Lorbeerartiger Jambuſenbaum. W.
Habitat in Zeylona. ♄

975. MYRTUS.

Gen. plant. edit. Schreb. n. 844.

Cal. 5-fidus. ſuperus. *Petala* 5. *Bacca* 2-ſ. 3-locularis polyſperma.

1. MYRTUS *communis.*
M. floribus ſolitariis: involucro diphyllo. *Hort. cliff.* 42. *Hort. upſ.* 123. *Mat. med.* 122. *Roy. lugdb.* 269. *Scop. carn. n.* 584. *Regn. bot.*
α. Myrtus communis, *romana.*
M. foliis ovatis, pedunculis longioribus. *Mill. dict. n.* 1. *et ic. t.* 184. *f.* 1. *Kniph. cent.* 4. *n.* 52.
M. latifolia romana. *Bauh. pin.* 468.
β. Myrtus communis, *tarentina.*
M. foliis ovatis, baccis rotundioribus. *Mill. dict. n.* 6.
M. minor vulgaris. *Bauh. pin.* 469.
γ. Myrtus communis, *italica.*
M. foliis ovato-lanceolatis acutis, ramis rectioribus. *Mill. dict. n.* 5.
M. communis italica. *Bauh. pin.* 468.
δ. Myrtus communis, *boetica.*
M. foliis ovato-lanceolatis confertis. *Mill. dict. n.* 4.
M. latifolia boetica 2. *Bauh. pin.* 469. *Cluſ. hiſt.* 1. *p.* 65. *Blackw. t.* 114.
ε. Myrtus communis, *luſitanica.*
M. foliis lanceolato-ovatis acutis. *Mill. dict. n.* 3.
M. ſylveſtris, foliis acutiſſimis. *Bauh. pin.* 469. *Cluſ. hiſt.* 1. *p.* 66. *f.* 1.
ζ. Myrtus communis, *belgica.*

M. foliis lanceolatis acuminatis. *Mill. dict. n.* 2.
M. latifolia belgica. *Bauh. pin.* 469.
ι. Myrtus communis, *mucronata.*
M. foliis lineari-acuminatis. *Mill. dict.*
M. foliis minimis et mucronatis. *Bauh. pin.* 469. *Cluſ. hiſt.* 1. *p* 67.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 643.
Gemeine Myrte. W.
Habitat in Europa *auſtrali*, Aſia, Africa. ♄ (v.v.)

*2. MYRTUS *tenuifolia.*
M. pedunculis axillaribus ſolitariis unifloris, foliis linearibus mucronulatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p* 280.*
Schmalblättrige Myrte. W.
Habitat in Nova Hollandia. ♄
Folia *oppoſita uncialia et ultra, lineam lata, margine revoluta, ſubtus pubeſcentia.* Pedunculi *ſericei foliis breviores.* Flores *albi ſaepius rubicundi duplo minores quam in M. communi.* Germen *ſericeum.* Calyx *glaber.* Petala *pubeſcentia.* Smith. l. c.

*3. MYRTUS *tomentoſa.*
M. pedunculis unifloris, foliis triplinerviis ſubtus tomentoſis. *Ait. Kew.* 2. *p.* 159. *Vahl. ſymb.* 2. *p.* 56.* *et* 3. *p.* 65.
M. (*caneſcens*) floribus ſolitariis ſparſis, foliis ovatis trinerviis tomentoſis, petiolis cauli connatis. *Loureiro. cochin. p.* 381.*
Arbor ſinenſis Canellae folio minore trinervi, prona parte villoſo, fructu Caryophylli aromatici majoris, villis ſimiliter obducto. *Pluk. amalth.* 21. *t.* 372. *f.* 1.
Filzige Myrte. W.
Habitat in China, Cochinchina. ♄

*4. MYRTUS *ceraſina.*
M. pedunculis lateralibus terminalibusque unifloris, foliis oblongis nitidis ſubtus punctatis. *Vahl. ſymb.* 2. *p.* 57.*
Arbor baccifera myrtifolia latiore fructu nigro ceraſino dipyreno. *Sloan. jam. hiſt.* 2. *p.* 107.
Kirſchartige Myrte. W.
Habitat in inſulis Caribaeis. ♄

*5. MYRTUS *procera.*
M. pedunculis confertis axillaribus unifloris, foliis ova-

tis acuminatis planis glabris, ramis virgatis, caule arboreo. *Swartz prodr.* 77. *Fl. ind. occid.* 2. *p.* 887.*
Hohe Myrte. W.
Habitat in ſylvis Hiſpaniolae *interioribus.* ♄
Foliis *cum M. monticola convenit*, pedunculis *vero florum diſtincta.* Folia *inodora.* Flores *fragrantes,* Swartz l. c.

*6. MYRTUS *bracteata.* W.
M. pedunculis unifloris axillaribus bracteatis, foliis ellipticis glabris punctatis, junioribus ſericeo-pubeſcentibus. W.
Beblätterte Myrte. W.
Habitat in India *orientali.* ♄ (v. ſ.)
Folia *oppoſita petiolata integerrima elliptica, utrinque parum attenuata obtuſa leviſſime venoſa glabra punctata imprimis in pagina inferiore.* Folia *juniora utrinque pilis ſericeis adpreſſis flaveſcentibus tecta.* Pedunculi *uniflori villoſi ſolitarii axillares in ramulis junioribus.* Bracteae *binae lanceolatae piloſae ad baſin germinis.* Germen *pubeſcens.* Calycis *laciniae* 4, *ſubrotundae pubeſcentes ciliatae pellucido-punctatae.* Fructus *mihi ignotus.* W.

7. MYRTUS *biflora.*
M. pedunculis bifloris, foliis lanceolatis. *Amoen. acad.* 5. *p.* 398.*
Caryophyllus fruticoſus, foliis lanceolatis oppoſitis, floribus geminatis alaribus. *Brown. jam.* 248. *t.* 25. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 716.
Zweyblütige Myrte. W.
Habitat in Jamaica. ♄

*8. MYRTUS *trinervia.*
M. pedunculis axillaribus trifloris, foliis ovatis acuminatis trinerviis ſubtus tomentoſis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 280.*
Habitat in Nova Hollandia. ♄
Folia *lata pulchra oppoſita ovata acuminata ſubtus tomentoſa trinervia ut in Blakea.* Flores *parvi plerumque* 3 *in pedunculo brevi piloſo furcato. Licet* 4 *ſint calycis laciniae et petala tamen eſt Myrti nec Eugeniae ſpecies.* Bacca *enim continet ſemina multa giboſo-curvata et eſt unilocularis; germen vero videtur bi-ſ. triloculare.* Smith. l. c.

9. MYRTUS *lucida.*

M. pedunculis ſubtrifloris, foliis ſubſeſſilibus lanceolatis attenuatis. *Sp. pl.* 674.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 717.

Glänzende Myrte. W.

Habitat Surinami. *Rolander.* ♄

Folia *ſingulari ſtructura, ex ovato admodum attenuata in apicem lanceolatum.* Flores *pentapetali.*

*10. MYRTUS *laevis.*

M. pedunculis umbellatis, foliis ovatis acuminatis. *Thunb. jap.* 198. *

Glatte Myrte. W.

Habitat in Japonia. ♄

Rami *teretes glabri.* Ramuli *pubeſcentes.* Folia *e geminis alternis ſolitaria, bina vel terna, petiolata ovata acuminata, acute ſerrata venoſa utrinque glabra, ſubtus pallida patentia inaequalia pollicaria et ultra.* Flores *in ramulis terminales umbellati.* Bacca 2-*ſ.* 3-*locularis.* Semina *ſolitaria in ſingulo loculamento. Valde affinis Crataego laevi et villoſae, differt vero a C. villoſa: quod tota glabra,* piſtillo *unico,* foliis *aggregatis; a C. laevi:* piſtillo *unico,* foliis *ovatis tenuibus longe acuminatis.* Thunb. l. c.

*11. MYRTUS *ruſcifolia.* W.

M. pedunculis ſubmultifloris axillaribus, foliis glaberrimis elliptico-ſubrotundis ſupra punctatis. W.

Mäuſedornblättrige Myrte. W.

Habitat in India *orientali.* ♄ (v. ſ.)

Folia *quodammodo iis Ruſci aculeati ſimilia, ſed parum majora, oppoſita breviſſime petiolata conferta elliptico-ſubrotunda obtuſiuſcula integerrima venoſa, ſupra impreſſo-punctata, inferne punctis elevatis obſita.* Folia *juniora pilis rufeſcentibus adpreſſis tecta.* Pedunculi *breves axillares conferti* 3-4-*ſ. multiflori pubeſcentes.* Floris *ſtructuram in ſpecimine male ſiccato haud axaminare potui.* Fructus *mihi ignotus.* W.

12. MYRTUS *dioica.*

M. pedunculis trichotomo-paniculatis, foliis oblongis floribus dioicis. *Sp. pl.* 675.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 645.

Getrennte Myrte. W.

Habitat in America. *Miller.* ♄

Folia *oppoſita, lanceolato-ovata craſſa.* Pedunculi

axillares et terminales, brachriato-paniculati, longitudine foliorum. Petala *pauca.*

*13. MYRTUS *lineata.*

M. floribus axillaribus ſubſeſſilibus, foliis ovatis acuminatis rigidis lineatis ſubtus incanis. *Swartz. prodr.* 78. *Fl. ind. occid.* 2. *p.* 891. *

Geſtreifte Myrte. W.

Habitat in campeſtribus montium Domingenſium. ♄

*14. MYRTUS *cordata.*

M. floribus axillaribus lateralibusque ſubſeſſilibus, foliis ſeſſilibus cordatis ovatis nitidis. *Swartz. prodr.* 78. *Fl. ind. occid.* 2. *p.* 893. * *Vahl. ſymb.* 2. *p.* 57. *

Herzblättrige Myrte. W.

Habitat in India *occidentali.* (v. ſ.)

*15. MYRTUS *pallens.*

M. racemis terminalibus pubeſcentibus, pedicellis unifloris, foliis lato-lanceolatis attenuatis ſupra punctatis. *Vahl. ſymb.* 2. *p.* 57. *

Bleiche Myrte. W.

Habitat in America *calidiore.* ♄

*16. MYRTUS *dumoſa.*

M. racemis axillaribus breviſſimis, foliis petiolatis lato-lanceolatis acuminatis acutis. *Vahl. ſymb.* 2. *p.* 57. *

Buſchige Myrte. W.

Habitat in America *meridionali.* ♄

*17. MYRTUS *glabrata.*

M. racemulis breviſſimis axillaribus multifloris, foliis ellipticis acuminatis convexis coriaceis glaberrimis. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 903. *

Spitzblättrige Myrte. W.

Habitat in fruticetis montium Hiſpaniolae. ♄ (v. ſ.)

*18. MYRTUS *diſticha.*

M. pedunculis axillaribus multifloris foliis brevioribus, foliis diſtichis deflexis ovato-lanceolatis ramis patentibus. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 894. *

Zweyzeilige Myrte. W.

Habitat in montibus Jamaicae *ſeptentrionalioris.* ♄

*19. MYRTUS *monticola.*

M. racemis multifloris breviſſimis axillaribus ſolitariis, foliis ovatis obtuſatis planis glaberrimis. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 898. *

Berg-Myrte. W.

Habitat in montibus excelſis Jamaicae. ♄. (v. ſ.)

*20. MYRTUS *Gregii.*

M. pedunculis multifloris axillaribus, foliis ellipticis acutis integerrimis ſubtus pubeſcentibus. *Swartz prodr.* 78. *Fl. ind. occid.* 2. *p.* 898. *

Greggia aromatica. *Gaert. ſem.* 1. *p.* 168. *t.* 33. *f.* 3.

Gregiſche Myrte.

Habitat in India *occidentali.* ♄ (v. ſ.)

*21. MYRTUS *virgultoſa.*

M. racemis lateralibus terminalibusque, foliis lato-lanceolatis attenuatis. *Vahl. ſymb.* 2. *p.* 58. *

M. racemis axillaribus terminalibusque paniculatis, pedicellis remotis decuſſatis unifloris, foliis ovato-lanceolatis attenuatis, ramis virgatis, caule arboreo. *Swartz prodr.* 79. *Fl. ind. occid* 2. *p.* 905. *

M. foliis ovato-lanceolatis. *Plum. amer.* 202. *ic.* 208. *f.* 1.

Vieläſtige Myrte. W.

Habitat in montibus altis Jamaicae. ♄

22. MYRTUS *zeylanica.*

M. racemis axillaribus terminalibusque, pedicellis ſubquadrifloris ovatis attenuatis ſupra punctatis. *Vahl. ſymb.* 2. *p.* 58.

M. pedunculis multifloris, foliis ovatis ſubpetiolatis. *Mill. dict. n.* 8.

M. foliis ovatis acuminatis obtuſiuſculis. *Fl. zeyl.* 182.

M. zeylanica odoratiſſima, baccis niveis monococcis. *Herm. lugdb.* 434. *t.* 435.

Myrtoides foliis ovatis. *Hort. cliff.* 489. *Roy. lugdb.* 535.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 718.

Zeylaniſche Myrte. W.

Habitat in Zeylona. ♄

*23. MYRTUS *ſplendens.*

M. pedunculis axillaribus ſolitariis trichotomis, foliis ellipticis attenuatis planis venoſis ſcarioſis nitidis, ramis virgatis. *Swartz prodr.* 79. *Fl. ind. occid.* 2. *p.* 907. *

Eugenia (*periplocifolia*) foliis integerrimis acuminato-lanceolatis, pedunculis paniculatis axillaribus et terminalibus *Iacq. collect.* 2. *p.* 108. *t.* 4. *

Langſpitzige Myrte. W.

Habitat in fruticetis montoſis Hiſpaniolae. ♄ (v. ſ.)

24. MYRTUS *adroſaemoides.*

M. racemis axillaribus terminalibusque ſubcompoſitis,

pedicellis trifloris, foliis ovato-oblongis ſeſſilibus. *Vahl. ſymb. 2. p. 58.* *

M. pedunculis trifido-multifloris, foliis ſubovalibus ſeſſilibus. *Fl. zeyl.* 184. *

Houttuyn Lin. Pfl. Syſt. 1. *p.* 719.

Apocynumblättige Myrte. W.

Habitat in Zeylona. ♄

25. MYRTUS *caryophyllata.*

M. pedunculis trichotomis terminalibus, calycibus indiviſis, foliis obovatis impunctatis. *Swartz obſ.* 202. *

M. pedunculis trifido-multifloris, foliis obovatis. *Fl. zeyl.* 183. *Mat. med.* 122.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 720.

Nägleinartige Myrte. W.

Habitat in Zeylona. ♄

*26. MYRTUS *acris.*

M. pedunculis axillaribus terminalibus corymboſisque trichotomis foliis longioribus, foliis ellipticis convexis coriaceis venoſis punctatis, caule arboreo. *Swartz prodr.* 79. *Fl. ind. occid.* 2. *p.* 909.

Myrtus caryophyllata. *Iacq. obſ.* 2. *p.* 1. *

Caryophyllus folio et fructu rotundo. *Breyn. prodr.* 2.

Caryophyllus aromaticus indiae occidentalis foliis et fructu rotundo. *Pluk. alm.* 188. *t.* 155. *f.* 3.

Caryophyllus foliis oblongo-ovatis oppoſitis, racemis lateralibus et terminalibus. *Brown. jam.* 247.

Caryophyllus racemoſus. *Mill. dict.*

Scharfe Myrte. W.

Habitat in montibus Jamaicae, Antiguae, Barbados. ♄ (v. ſ.)

*27. MYRTUS *coriacea.*

M. pedunculis trichotomis terminalibus, foliis ſubrotundo-ellipticis convexis coriaceis aveniis punctatis utrinque nitidis. *Vahl. ſymb.* 2. *p.* 59. * *Swartz Fl. ind. occid.* 2. *p.* 912. *

Myrtus acris. *β. Swartz prodr.* 79.

M. cotini folio. *Plum. ic.* 208. *f.* 2.

Lederartige Myrte. W.

Habitat in inſulis Hiſpaniola, St. Lucia. ♄

28. MYRTUS *Pimenta.*

M. foliis alternis. *Fl. zeyl.* 180. *Mat. med.* 122. *Swartz. obſ.* 202. *

M. calycibus absque appendiculis. *Hort. cliff.* 501.

M. foliis oblongo-ovatis glabris alternis, racemis terminalibus et lateralibus. *Brown. jam.* 247.
M. arborea aromatica, foliis laurinis. *Sloan. jam.* 161. *hiſt.* 2. *p.* 76. *t.* 191. *f.* 1.
Caryophyllus foliis oblongo-ovatis alternis, racemis terminalibus et lateralibus. *Brown. jam.* 247.
Caryophyllus aromaticus americanus, lauri acuminatis foliis, fructu orbiculari. *Pluk. alm.* 88. *t.* 155. *f.* 4.
Piper jamaicenſe. *Raj. hiſt.* 1507. *Blackw. t.* 355. R.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 722.
Pfefferartige Myrte. W.
Habitat in India *occidentali.* ♄ (v. ſ. ſ. fl.)
Calyx *fructus infe:us quadridentatus; floris vero ſuperus quadripartitus.* Bacca 2-*ſperma.*

*974. CALYPTRANTHES.
Gen. pl. ed. Schreb. n. 845.

Cal. ſuperus truncatus ante antheſin tectus *operculo* integerrimo deciduo. *Cor.* 0. *Bacca* unilocularis 1-4 ſperma.

1. CAYPTRANTHES *Suzygium.*
C. arboreſcens, pedunculis axillaribus trichotomis patulis, foliis ovatis obtuſis, ramis dichotomis. *Swartz prodr.* 79. *Fl. ind. occid.* 2. *p.* 919.*
Myrtus (*Zuzygium*) pedunculis multifloris, ramis dichotomis, foliis geminis ſubovatis terminalibus. *Amoen acad.* 5. *p.* 398.* *Swartz obſ.* 202.
Suzygium fruticoſum, foliis ovatis nitidis et ramulis ubique jugatis. *Brown. jam.* 240. *t.* 7. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 646.
Gabeläſtige Deckelmyrte. W.
Habitat in maritimis aridis calcareis Jamaicae. ♄

*2. CALYPTRANTHES *guineenſis.* W.
C. arboreſcens, pedunculis axillaribus et terminalibus corymboſis, foliis oblongis utrinque attenuatis reticulatis. W.
Guineaſche Deckelmyrte. W.
Habitat in Guinea. ♄ (v. ſ.)
Folia *oppoſita oblonga ſeu elliptica integerrima, utrinque parum attenuata, ſuperne nitida, reticulato-venoſa.* Corymbi *divaricati in apice caulis et axillis ſupremis foliorum.* Pedunculi *laterales biflori.* W.

3. CALYPTRANTHES *caryophyllifolia.*

C. arborea, paniculis lateralibus, foliis elliptico-ovatis integerrimis. *Willd. in Uſter. neue Annalen* 11. *Stück. p.* 22. *

Eugenia (*caryophyllifolia*) foliis ovato-lanceolatis acuminatis, racemis paniculatis, calyce truncato. *Lamarck. encycl.* 3. *p.* 189. *

Eugenia (*corticoſa*) foliis ovatis acuminatis, racemis corymboſis, filamentis breviſſimis. *Loureiro. cochin. p.* 376. ?

Myrtus (*Cumini*) pedunculis multifloris, foliis lanceolato-ovatis. *Sp. pl.* 674. *Fl. zeyl.* 185. *

Jamboſa ceramica. *Rumph. amb.* 1. *p.* 130. *t.* 41. *Burm. ind.* 115.

Arbor zeylanica cuminum redolens. *Burm. zeyl.* 27.

Caryophyllus langueſcente vi aromaticus, Malabarenſis folio, fructu maximo. *Pluk. alm.* 88. *t.* 274. *f.* 2.

Peria Njara. *Rheed. mal.* 5. *p.* 57. *t.* 29.

Gewürznägleinblättrige Deckelmyrte. W.

Habitat in India *orientali.* ♄ (v. ſ.)

Panicula *in ramis aphyllis praeteriti anni.* W.

*4. CALYPTRANTHES *Jambolana.* W.

C. arborea, paniculis ſubterminalibus, foliis ovatis emarginatis. *Willd. in Uſteri neue Annalen* 11. *Stück. p.* 23. *

Eugenia (*Jambolana*) foliis ovalibus venoſis et tenuiſſime punctatis, racemis paniculatis lateralibus, calyce truncato. *Lamarck. encycl.* 3. *p.* 150.

Jambolana. *Rumph. amb.* 1. *p.* 131. *t.* 42.

Jambolifera pedunculata. *Houttuyn Lin. Pfl. Syſt.* 1. *p.* 462. *tab.* 7. *f.* 2.

Ausgerandete Deckelmyrte. W.

Habitat in India *orientali.* ♄

5. CALYPTRANTHES *Chytraculia.*

C. arborea, pednuculis terminalibus paniculatis trichotomis tomentoſis, foliis ovatis apice attenuatis. *Swartz prodr.* 79. *Fl. ind. occid.* 2. *p.* 921. *

Myrtus (*Chytraculia*) pedunculis dichotomis paniculatis tomentoſis, foliis geminis ſubovatis terminalibus. *Amoen. acad.* 5. *p.* 398. * *Swartz obſ.* 202.

Chytraculia arborea, foliis ovatis glabris oppoſitis, racemis terminalibus. *Browu. jam.* 239. *t.* 37. *f.* 2. *Houttuyn Lin. Pfl. Syſt.* 3 *p.* 646.

Behaartſtielige Deckelmyrte. W.
Habitat in montoſis ſiccis Jamaicae. ♄

*6. CALYPTRANTHES *rigida.*
C. arboreſcens, pedunculis ſolitariis ſubtrifloris, foliis ovatis acutis convexis aveniis rigidis. *Swartz prodr.* 80. *Fl. ind. occid.* 2. *p.* 923.*
Steifblättrige Deckelmyrte. W.
Habitat in ſummis montibus Jamaicae, *ut in excelſo monte* Catharin Hill *paroecia* St. Andreae. ♄

*975. EUCALYPTUS.
l'Herit. ſert. angl. 18.

Cal. ſuperus perſiſtens truncatus ante antheſin tectus *operculo* integerrimo deciduo. *Corolla* nulla. *Capſula* quadrilocularis apice dehiſcens polyſperma.

* *operculo conico.*

*1. EUCALYPTUS *robuſta.*
E. operculo conico medio conſtricto calyce latiori, umbellis lateralibus terminalibusque, foliis ovatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 283.
E. (*robuſta*) operculo conico medio conſtricto, umbellis lateralibus terminalibusque, pedunculis pedicellisque compreſſis. *Smith. Nov. Holl. p.* 39. *t.* 13.*
Hohe Schönmütze. W.
Habitat in Nova Hollandia. ♄ (v. ſ.)

*2. EUCALYPTUS *pilularis.*
E. operculo conico medio conſtricto longitudine calycis umbellis lateralibus, fructu globoſo, foliis lineari-lanceolatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 284.
Pillentragende Schönmütze. W.
Habitat in Nova Hollandia. ♄
Folia *multo anguſtiora quam in praecedente et* flores *dupla minores; operculum vero calyce duplo latior,* fructus *globoſus.* Fructus *E. robuſtae videtur turbinatus et margine reflexus, ſed immaturum tantum vidi.* Smith. l. c.

*3. EUCALYPTUS *tereticornis.*
E. operculo conico tereti laeviſſimo membranaceo ca-

lyce latiore triploque longiore, umbellis lateralibus ſolitariis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 284.

E. (*tereticornis*) operculo conico tereti laeviſſimo calyce triplo longiori, umbellis lateralibus ſolitariis. *Smith. Nov. Holl. p.* 41.

Rundhörnige Schönmütze. W.

Habitat in Nova Hollandia. ♄

Diſtincta operculo *membranaceo laeviſſimo baſi dehiſcente et calyci annuli ad inſtar adhaerente*, foliisque *lanceolatis obliquis.* Smith. l. c.

*4. EUCALYPTUS *reſinifera*

E. operculo conico tereti coriaceo calyce duplo longiori, umbellis lateralibus ſolitariis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 284.

Eucalyptus reſinifera. *White. itin.* 231. *cum. icon.*

Metroſideros gummifera. *Gaert. ſem.* 1. *p.* 170. *t.* 34. *f.* 1.

Harzbringende Schönmütze. W.

Habitat in Nova Hollandia. ♄

Primo intuitu ſimillima praecedenti, *ſed attente examinata*, *differt*: operculo *calyce duplo longiore latitudine calycis, nec latiore*, *craſſo coriaceo, nec membranaceo laeviſſimo*, *nec baſi calyci adhaerente.* Smith. l. c.

*5. EUCALYPTUS *capitellata.*

E. operculo conico obtuſiuſculo calyceque anguloſo ſubancipiti, capitulis lateralibus ſolitariis, fructu globoſo, foliis ovato-lanceolatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 285.

E. (*capitellata*) operculo conico calyceque anguloſo ſubancipiti, capitulis lateralibus pedunculatis ſolitariis. *Smith. Nov. Holl.* 42.* *White. itin.* 226. *c. icone. fig. a.*

Kopfförmige Schönmütze. W.

Habitat in Nova Hollandia. ♄ (v. ſ.)

Differt ab omnibus praecedentibus; floribus *capitatis*, *id eſt absque pedunculis partialibus*, *umbellatis.* Operculum *eſt longitudine calycis angulatum apice compreſſum.* Folia *ovato-lanceolata rigida obliqua.* Smith l. c.

*6. EUCALYPTUS *ſaligna.*

E. operculo conico acuto calyceque anguloſo ſubancipiti, capitulis lateralibus ſolitariis fructu turbinato, foliis

lineari - lanceolatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 285.
Weidenblättrige Schönmütze. W.
Habitat in Nova Hollandia. ♄

Folia *anguſtiora et minus coriacea quam in omnibus reliquis.* Capitula *parva florum brevi pedunculo ſunt ſuffulta et ſolitaria ad foliorum baſin.* Flores *reliquis ſpeciebus minores.* Operculum *acutum longitudine calycis.* Fructus *turbinatus parum margine reflexus et baſi ſtyli pyramidali coronatus.* Smith. l. c.

* * *Operculo hemiſphaerico.*

*7. EUCALYPTUS *botryoides.*

E. operculo hemiſphaerico ſubmutico, capitulis lateralibus ſolitariis, pedunculis cuneatis compreſſis, fructu turbinato. *Smith. Act. ſoc. Lin. Lond* 3. *p.* 286.
Traubige Schönmütze. W.
Habitat in Nova Hollandia. ♄

Similis binis praecedentibus capitulis *ſolitariis, ſed diſtincta:* operculo *latiore hemiſphaerico apice ſubmucronato.* Flores *coarcti et faſciculo baccarum haud abſimiles.* Pedunculus *communis compreſſus et latus, praecipue apicem verſus.* Folia *lanceolata obliqua.* Smith. l. c.

*8. EUCALYPTUS *haemaſtoma.*

E. operculo hemiſphaerico depreſſo mucronulato, umbellis lateralibus terminalibusque, pedunculis compreſſis, ramulis angulatis, fructu ſubgloboſo. *Smith. Act. ſoc Lin. Lond.* 3. *p.* 286.
Rothrandige Schönmütze. W.
Habitat in Nova Hollandia. ♄

Folia *coriacea lanceolata in acumen lineare longum producta.* Flores *umbellati non capitati,* operculum *depreſſum mucronatum.* Fructus *globoſus apice dehiſcens, orificio margine lato rubro cincto.* Smith. l. c.

*9. EUCALYPTUS *piperita.*

E. operculo hemiſphaerico mucronulato, umbellis lateralibus ſubpaniculatis ſolitariisve, pedunculis compreſſis, ramulis angulatis. *Smith. Act. ſoc. Lin. Lond.* 3. *p.* 286. *Smith. Nov. Holl.* 42.* *White. itin.* 226. *c. icone.*
Pfefferartige Schönmütze. W.
Habitat in Nova Hollandia. ♄ (v. ſ.)

Diſtincta a praecedente cui ſimillima: foliis *ovatis licet obliqua ſint*, floribus *minoribus in ſuprema ramorum parte copioſis, non in apice ramorum. Paucae* umbellae *ſunt ſolitariae, reliquae in paniculis vel corymbis aggregatae.* Smith. l. c.

*10. EUCALYPTUS *obliqua.*

E. operculo hemiſphaerico mucronulato, umbellis lateralibus ſolitariis, pedunculis ramulisque teretibus. *Smith. Act. ſoc. Lin. Lond. 3. p. 287. Nov. Holl. 43.* * *Ait. Kew. 2. p. 157. l'Herit. ſert. angl. 18.*

Schiefblättrige Schönmütze. W.

Habitat in Nova Hollandia. ♄ (v. ſ. ſ. fl.)

*11. EUCALYPTUS *corymboſa.*

E. operculo hemiſphaerico mucronulato, calyce tereti, umbellis corymboſo-paniculatis terminalibus. *Smith. Act. ſoc. Lin. Lond. 3 p. 287. Nov. Holl.* 43.

Doldentraubige Schönmütze. W.

Habitat in Nova Hollandia. ♄ (v. ſ.)

Flores *magni ſpecioſi formant paniculam magnam terminalem ex umbellis compoſitam, qua nota a reliquis diſtincta.* Folia *lanceolata coriacea.* Fructus *turbinatus* calyce *urceolato perſiſtenti. et* ſtylo *in hac cavitate perſiſtenti coronatus.* Smith. l. c.

*12. EUCALYPTUS *paniculata.*

E. operculo hemiſphaerico ſubmutico, calyce anguloſo, umbellis ſubpaniculatis terminalibus. *Smith. Act. ſoc. Lin. Lond. 3. p. 287.*

Riſpenblütige Schönmütze. W.

Habitat in Nova Hollandia. ♄

Differt a praecedente: calyce *angulato* operculo *ſubmucronato, et quod omnibus partibus minor ſit.* Umbellae *formant nunquam paniculam amplam ſed parvam; rami inferiores paniculae ſunt axillares.* Smith. l. c.

*976. B A N A R A.

Gen. plant. edit. Schreb. n. 809.

Cal. inferus 4-fidus. *Corolla* tetrapetala. *Bacca* 1-locularis polyſperma.

*1. BANARA *fagifolia.*

B. fagifolia. *Vahl. ſymb. 3. p.* 65. *

B. gujanenſis. *Aubl. guj. 1. p. 547. t.* 217. *

Buchenblättrige Banare. W.
Habitat in Cajenna. ♄
Frutex *ramis teretibus cineraſcentibus.* Folia *alterna oblonga acuminata venoſa glabra ſerrata, ſerraturis ſubtus glanduloſis.* Racemus *terminalis ſimplex pedunculatus palmaris villoſus.* Vahl. l. c.

*977. ANTHERYLIUM.

Rohr. Act. ſoc. Hiſt. Nat. Hafn. 2. *p.* 211.

Cal. inferus 4-partitus. *Cor.* tetrapetala. *Capſ.* 1-locularis trivalvis polyſperma.

*1. ANTHERYLIUM *Rohrii.*
A. Rohrii *Vahl. Act. ſoc. Hiſt. Nat. Hafn.* 2. 1. *p.* 212. *t.* 8 * *Vahl. ſymb.* 3 *p.* 66.
Rohriſches Antherylium. W.
Habitat in inſula St. Thomae Indiae *occidetalis.* ♄ (v. ſ.)
Arbor. Rami *teretes glabri a caſu foliorum tuberculati.* Ramuli *ſuboppoſiti, inferne teretiuſculi, superne e lineis decurrentibus ſubtetragoni, lateribus duobus oppoſitis convexis, duobus ſulco exaratis.* Folia *petiolata oppoſita et alterna ovata, interdum obovata integerrima ſubemarginata.* Aculeus *brevis ad ſingugulum latus petioli.* Pedunculi 5-8 *axillares uniflori.* Stamina 12-16 *calyci inſerta. Affinis Legnoti.* Vahl. l. c.

*978. FOETIDIA.

Juſſ. gen. ed. Uſteri p. 361.

Cal. ſuperus 4-fidus. *Cor.* o. *Capſ.* lignoſa 4-locularis, loculis 1-ſ. 2-ſpermis.

*1. FOETIDIA *mauritiana.*
F. mauritiana. *Lamarck. encycl.* 2. *p.* 453. * *Lamarck. illuſt. gen. t.* 419.
Einblütiger Stinckbaum. W.
Habitat in inſula Mauritii. ♄
Arbor *ramis confertis fere verticillatis glabris.* Folia *ſubcoriacea elliptica integerrima ſeſſilia avenia glabra.* Pedunculi *uniflori terminales.* W.

*979. SCOLOPIA.

Gen. plant. edit. Schreb. n. 846.

Cal. inferus 5-ſ. 4-partitus. *Corolla* 5-ſ. 4-petala. *Bacca* ſtylo coronata 1-locularis 6-ſperma. *Sem.* arillata.

*1. SCOLOPIA *puſilla.* †

Limonia puſilla. *Gaert. ſem.* 1. *p.* 279. *t. t.* 58. *f.* 4.
Malus limonia lentiſci folio zeylanica, fructu minimo, uvariae magnitudine. *Pluk. alm.* 239. ?
Khatu-Karundu *Zeylon.*
Niedrige Scolopie. W.
Habitat in Zeylona. ♄

980. PUNICA.

Gen. plant. ed. Schreb. n. 847.

Cal. 5-fidus, ſuperus. *Petala* 5. *Pomum* multiloculare, polyſpermum.

1. PUNICA *Granatum.*

P. foliis lanceolatis, caule arboreo. *Hort. cliff.* 134. *Hort. upſ.* 122. *Mat. med.* 183. *Roy. lugdb.* 267. *Gron. orient.* 144. *Mill. dict. n.* 1. *Scop. carn. ed.* 2. *n.* 585. *Gmel. it.* 3. *p.* 33. *Ludw. ect. t.* 13. *Blackw. t.* 97 *et* 145. *Kniph. cent.* 2. *n.* 64. *Mill. illuſtr. Regn. bot.*
P. ſpinoſa, foliis nitentibus, ellipticis, integerrimis, floribus ſeſſilibus. *Hall. helv. n.* 1098.
Malus punica ſylveſtris. *Bauh. pin.* 438.
Malus punica ſativa. *Bauh. pin.* 438.
Malus punica. *Cam. epit.* 130. 131.
β. Balauſtia flore pleno majore et minore. *Bauh. pin.* 438. *Knorr. del.* 1. *t.* 5.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 648.
Gemeine Granate. W.
Habitat in Hiſpania, Italia, Mauritania, Perſia, Helvetia, Carniolia *ſolo cretaceo.* ♄ (v. v.)

2. PUNICA *nana.*

P. foliis linearibus, caule fruticoſo. *Mill. dict. n* 2. *Du Roi. harbk.* 2. *p.* 204.
P. fruticoſa humilior, ramulis gracilibus patentibus. *Brown. jam.* 239.

P. americana nana ſ. humillima. *Tournef. inſt.* 656. *Du Ham. arb.* 9.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 650.
Zwerg-Granate. W.
Habitat in Antillis. ♄ (v. ſ.)
Altitudo *pedum* 5. Folia *breviora, anguſtiora.* Floret *tota aeſtate.*

981. AMYGDALUS.

Gen. plant. ed. Schreb. n. 848.

Cal. 5-fidus, inferus. *Petala* 5. *Drupa* nuce poris perforata.

1. AMYGDALUS *Perſica.*

A. foliorum ſerraturis omnibus acutis, floribus ſeſſilibus ſolitariis. *Hort. cliff.* 189. *Hort. upſ.* 123. *Mat. med.* 123. *Roy. lugdb.* 267. *Du Roi. harbk.* 1 *p.* 56. *Ludw. ect. t.* 194. *Knorr. del* 1. *t. P.* 1. *Regn. bot. Willd. arb.* 18.
Perſica vulgaris, *Mill. dict. n.* 1.
Perſica molli carne et vulgaris. *Bauh. pin.* 440.
Perſica rubra. *Cam. epit.* 145. *Blackw. t.* 101.
β. Nuciperſica, quae nucium juglandium faciem repraeſentat. *Bauh. pin.* 440. *Raj. hiſt.* 1516.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 724.
Perſiſche Mandel. W.
Habitat in Perſia. ♄ (v. v.)
Flores *ſeſſiles, gemmales, ſolitarii.* Folia *e gemma terminali, nec infra flores.*
Variat *floribus plenis.* R.

2. AMYGDALUS *communis.*

A. foliorum ſerraturis infimis glanduloſis, floribus ſeſſilibus geminis. *Hort. cliff.* 186. *Hort. upſ.* 123. *Mat. med.* 123. *Roy. lugdb.* 267. *Gron. orient.* 146. *Du Roi. harbk.* 1. *p.* 48. *Lud. ect. t* 177. *Reg. bot.*
A. foliis glabris, ovato-lanceolatis, ſerratis, imis dentibus et petiolis glanduloſis. *Hall. helv. n.* 1078.
Amygdalus ſylveſtris. *Bauh. pin.* 442.
β. Amygdalus ſativa. *Bauh. pin.* 441. *Mill. ic. t.* 28. *f.* 1. *Du Ham. arb. fruit.* 5. *Kniph. cent.* 6. *n.* 5. *Knorr. del.* 1. *t. M.* 1.
Amygdalus. *Dod. pempt.* 798.

γ. Amygdalus amara. *Tournef. inſt.* 627. *Blackw. t.* 105. *Du Ham. arb. fruit.* 7.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 727.
Gemeine Mandel. W.
Habitat in Mauritaniae *ſepibus*, Helvetia. ♄ (v. v.)
Flores *gemmales*, *ſeſſilis ſ. ſubſeſſiles*, *ſaepius bini.* Folia *e gemmis infra et ſupra flores locatis.*
Italis *ignota Catonis aevo: unde Nuces graecae dictae.*

3. AMYGDALUS *pumila.*
A. foliis lanceolatis duplicato-ſerratis. *Willd. arb.* 19.
A. foliis venoſo-rugoſis. *Syſt. Veg.* 385. *Herm. lugdb.* 487. *t.* 489. *Medic. in. obſ. Soc. Lutr.* 1774. *p.* 143. *Du Roi. harbk.* 1. *p.* 53.
A. perſica nana flore carneo pleno. *Pluk. phyt. t.* 11. *f.* 4.
Perſica africana nana, flore incarnato pleno. *Tournef. inſt.* 625. *Mill.* 3.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 729.
Niedrige Mandel. W.
Habitat in Africa. ♄ (v. v.)
Rami *laevis, bi-ſ. tripedales, atropurpurei.* Folia *lanceolata, petiolata, venoſa, rugoſa. ſerrulata.* Flores *ad gemmas ſaepius bini, ſeſſiles.* Calyx *rufeſcens.* Petala *emarginata, incarnata, tubo calycis longiora.* Filamenta *magis pallida.* Germen *et* Stylus *inferne pubeſcens.* Stipulae *lineares, profundius ſerratae.* H V.
Variat *floribus plenis.* Petala *non emarginata, ſed acuminata.* R.

4. AMYGDALUS *nana.*
A. foliis ovatis baſi attenuatis ſimpliciter argute ſerratis. *Willd. arb.* 19.
A. foliis baſi attenuatis. *Hort. cliff.* 186. *Hort. upſ.* 124. *Roy. lugdb.* 267. *Mill. dict. n* 5. *Du Roi. harbk.* 1. *p.* 51. *Du Ham. arb. fruit.* 1. *t.* 3. *Pall. it.* 1. *p.* 81. *Kniph. cent.* 6. *n.* 4. *Pall. roſſ.* 1. *p.* 12. *t.* 6.
Prunus inermis, foliis ex lineari-lanceolatis, calycum laciniis oblongis. *Gmel. ſib.* 3. *p.* 171. *n.* 2.
A. indica nana. *Pluk. alm.* 28. *t.* 11. *f.* 3. *Mill. dict. t.* 28. *f.* 2.
Armeniaca perſicae foliis, fructu exſucco. *Amm. ruth.* 273. *t.* 30. *Burm. ind.* 117.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 730.

Zwerg-Mandel. W.
Habitat in Calmuckia *copioſiſſima.* ♄ (v. v.)

Caules *pedales.* Radix *repens.* Stipulae *lineares.* Folia *lanceolata, ſerrata, laevia, in petiolos deſinentia.* Flores *ſeſſiles, 2. ſ. 3. ſimul, rubri.*

*5. AMYGDALUS *incana.*

A. foliis oblongo-lanceolatis ſerratis ſubtus tomentoſis. W.
A. foliis lanceolatis ſerratis rugoſis ſubſeſſilibus, ſubtus albo-tomentoſis. *Pall.* 1. *p.* 13. *t.* 7. *
Weiſsblättrige Mandel. W.
Habitat in herbidis Caucaſi. ♄

Facies praecedentis, ſed folia anguſtiora ſupra viridia glabra, infra niveo-tomentoſa.

*6. AMYGDALUS *orientalis.*

A. foliis lanceolatis integerrimis argenteis perennantibus petiolo breviore. *Mill. dict. Ait. Kew.* 1. *p.* 162.
Amygdalus argentea. *Lamarck. encycl.* 1. *p.* 102. *
A. orientalis, foliis argenteis ſplendentibus. *Duhamel. arb.* 48.
Silberblättrige Mandel. W.
Habitat in Oriente. ♄ (v. v.)

Folia *vix uncialia oblonga baſi et apice acuta integerrima, utrinque ſericeo-tomentoſa.* W.

982. PRUNUS.

Gen. plant. edit. Schreb. n. 849.

Cal. 5-fidus, inferus. *Petala* 5. *Drupae* nux ſuturis prominulis.

1. PRUNUS *Padus.*

P. floribus racemoſis, racemis pendulis, foliis deciduis duplicato-ſerratis ſubrugoſis, petiolis biglanduloſis. *Willd. arb.* 237.
P. floribus racemoſis, foliis deciduis baſi ſubtus biglanduloſis. *Gmel. ſib.* 3. *p.* 171. *Du Roi. harbk.* 2. *p.* 188. *Crantz. auſtr.* 1. *p.* 94. *Scop. carn.* 2. *n.* 589. *Pollich. pal. n.* 465. *Flor. dan. t.* 205. *Doerr. naſſ. p.* 264. *Hoffm. germ.* 169. *Roth. germ.* 1. 211. *II.* 538.
Padus glandulis duabus baſi foliorum ſubjectis. *Hort. cliff.* 185. *Fl. ſuec.* 396, 431. *Roy. lugdb.* 269.
Padus foliis annuis. *Fl. lapp.* 198.

Padus avium. *Mill. dict. n.* 1.
Padus foliis ovato-lanceolatis ſerratis *Hall. helv. n.* 1086.
Ceraſus racemoſa ſylveſtris, fructu non eduli. *Bauh. pin.* 451. *Tabern.* 988. *Du Ham. arb.* 3.
β. Prunus (*rubra*) floribus racemoſis, racemis erectis, foliis deciduis tenuiſſime duplico-ſerratis laevibus, petiolis biglanduloſis. *Willd. arb.* 237. *t.* 4. *f.* 2. *excluſo ſynonymo.*
Houttuyn Lin. Pfl. Syſt. 1. *p.* 731.
Vogel Pflaume. W.
Habitat in Europa. ♄ (v. v.)

Racemi *foliacei infra folia.* Petala *ſerrata ut in ſequenti, nec emarginata ut in Ceraſo.*

Prunus rubra a me in arboret. berol. l. c. deſcripta, eſt ſola hujus varietas, foliorum ſerraturis minoribus et racemis erectoribus. W.

2. PRUNUS *viginiana.*

P. floribus racemoſis, racemis rectis, foliis deciduis duplicato-dentatis laevibus, petiolis ſubquadriglanduloſis. *Willd. arb.* 238. *t.* 5. *f.* 1.*
P. floribus racemoſis, foliis deciduis baſi antice glanduloſis. *Sp. pl.* 677.
Prunus (*rubra*) floribus racemoſis, racemis erectis, foliis deciduis laevibus baſi biglanduloſis. *Ait. Kew.* 2. *p.* 162.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 734. *nomen nec deſcriptio.*
Virginiſche Pflaume. W.
Habitat in Virginia. ♄ (v. v.)

Padi progenies, *ſed verrucae in ramis duplo majores.* Petioli *antice ad apicem binis paribus glandularum.* Foliis *minus ſubtus reticulata et fere glabra.* Petala *orbiculata, nec obovata.* Fructus *quadruplo majores rubri.*

Diſtincta a ſequente, qua cum, ob ſynonyma auctorum falſa, confuſa: foliis *oblongis acuminatis argute inaequaliter duplicato-dentatis utrinque glabris,* petiolis *in foliis adultis binis paribus glandularum inſtructis,* racemis *longioribus, junioribus rectis, ſenioribus inferne jam defloratis laxiuſculis, a praecedente vero diverſa,* foliorum *forma, glandularum petioli diſpoſitione,* floribus *minoribus,* petalis *ſubrotundis. Huc ſine dubio pertinere videtur Prunus rubra Aitoni.* W.

*3. PRUNUS *ſerotina.*

P. floribus racemoſis, racemis laxis, foliis deciduis ſimpliciter ſerratis, ſerraturis infimis ſubglanduloſis. *Willd. arb.* 239. *t.* 5. *f.* 2. *

P. (*ſerotina*) foliis lato-lanceolatis acuminatis ſimpliciter ſerratis, glaberrimis ſubglanduloſis deciduis, ſerraturis adpreſſo-inflexis, infimis ſaepe petiolo inſidentibus, coſta ſubtus bifariam pubeſcente, racemis ſimplicibus, petalis integerrimis. *Ehrh. Beitr.* 3. *p.* 20.

Prunus virginiana. *Mill. dict. n.* 3. *Du Roi. harbk.* 2. *p.* 191. *Wangenh. am.* 34 *t.* 14. *f.* 33

Ceraſus ſylveſtris, fructu nigricante in racemis longis pendulis phytolaccae inſtar congeſtis. *Gron. virg.* 54. *Roy. lugdb.* 537. *Du Ham. arb.* 5.

Spätblühende Pflaume. W.

Habitat in America *boreali.* ♄ (v. v.)

Dignoſcitur facillime, licet foliorum, forma valde ſit varians, coſta media, ſubtus baſin verſus, barbata. W.

4. PRUNUS *canadenſis.* †

P. floribus racemoſis, foliis deciduis eglanduloſis lato-lanceolatis rugoſis utrinque pubeſcentibus. *Sp. pl.* 678.

Ceraſus racemoſa, foliis amygdalinis, americana. *Pluk. alm.* 97. *t.* 58. *f.* 5.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 735.

Canadiſche Pflaume. W.

Habitat in America *ſeptentrionali.* ♄

Rami *laeves.* Folia *lato-lanceolato absque glandulis, venoſo-rugoſa minusque rigida quam in reliquis, argute ſerrata, utrinque viridia, ad tactum manifeſte villoſa, et baſi in petiolum deſinentia.*

*5. PRUNUS *elliptica.*

P. floribus racemoſis, foliis ellipticis ſerratis glabris. *Thunb. jap.* 199.

Elliptiſchblättrige Pflaume. W.

Habitat culta in Japonia. ♄

Caulis *arboreus.* Rami *et* Ramuli *alterni nodoſi albo-punctati rugoſi glabri.* Folia *in ramulis aggregata ſparſa breviter petiolata elliptica obtuſiuſcula venoſa glabra patentia digitalia.* Flores *racemoſi pauci.* Drupa *oblonga magnitudine Paſſulae.* Flores *non vidi.* Thunb. l. c.

*6. PRUNUS *occidentalis.*

P. floribus racemoſis, racemis lateralibus, foliis perennantibus eglanduloſis oblongis acuminatis integerrimis utrinque glabris. *Swartz prodr.* 80. *Fl. ind. occid.* 2. *p.* 925.* *Ait. Kew.* 2. *p.* 163.

Ceraſus laticre folio, fructu racemoſo purpureo majore. *Cateſb. carol.* 2. *p.* 94. *t.* 94.? W.

Amygdalis foliis magnis. *Nicolſ. doming.* 154.

Weſtindiſche Pflaume. W.

Habitat in Antillarum *montibus interioribus.* ♄

*7. PRUNUS *ſphaerocarpa.*

P. floribus racemoſis, racemis axillaribus, foliis perennantibus eglanduloſis integerrimis nitidis, drupis ſubrotundis. *Swartz prodr.* 81. *Fl. ind. occid.* 2. *p.* 927.*

Myrtifolia arbor, foliis latis ſubrotundis, flore albo. *Sloan. jam.* 162. *hiſt.* 2. *p.* 79. *t.* 193. *f.* 1. *mala. Roy. dendr.* 36.

Amygdalus foliis parvis. *Nicolſ. doming.* 145.

Kugelfrüchtige Pflaume. W.

Habitat in montium ſylvis Jamaicae, Hiſpaniolae. ♄ (v. ſ.)

8. PRUNUS *luſitanica.*

P. floribus racemoſis, foliis ſempervirentibus ovato-lanceolatis ſerratis eglanduloſis. *Ait. Kew.* 2. *p.* 163. *Willd. arb.* 241.*

P. floribus racemoſis, foliis ſempervirentibus eglanduloſis. *Mill. ic.* 131. *t.* 196. *f.* 1.

Padus luſitanica. *Mill. dict. n.* 5.

Padus foliis glandula deſtitutis. *Vir. cliff.* 42. *Hort. upſ.* 126. *Roy. lugdb.* 269.

Padus foliis ſempervirentibus ovatis. *Hort. cliff.* 185.

Lauro Ceraſus luſitanica minor. *Dill. elth.* 193. *t.* 159. *f.* 193. *Tournef. inſt.* 628.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 736.

Portugiſiſche Pflaume. W.

Habitat in Luſitania, Penſylvania. ♄ (v. v.)

*9. PRUNUS *caroliniana.*

P. foliis racemoſis, foliis ſempervirentibus oblongo-lanceolatis ſerratis eglanduloſis. *Ait. Kew.* 2. *p.* 163.

Padus (*carolina*) foliis lanceolatis acute denticulatis ſempervirentibus. *Mill. dict. n.* 6. *Du Roi. harbk.* 2. *p.* 198.

Caroliniſche Pflaume. W.
Habitat in Carolina *auſtrali.* ♄

10. PRUNUS *Lauro-Ceraſus.*

P. floribus racemoſis, foliis ſempervirentibus dorſo biglanduloſis. *Mat. med.* 124. *Du Roi. harbk.* 1. *p.* 196. *Blackw. t.* 512. *Willd. arb.* 242.

Padus lauroceraſus. *Mill. dict. n.* 4.

Padus glandulis duabus dorſo foliorum innatis. *Vir. cliff.* 42. *Hort. upſ.* 126. *Roy. lugdb.* 269.

Padus foliis ſempervirentibus lanceolato-ovatis. *Hort. cliff.* 185.

Ceraſus folio laurino. *Bauh. pin.* 450.

Lauroceraſus. *Cluſ. hiſt.* 1. *p.* 4. *Cam. hort. t.* 23. *Tournef. inſt.* 627. *Raj. hiſt.* 1549. *Du Ham. arb.* 1. *t.* 133.

Houttuyn Lin. Pfl. Syſt. 1 *p.* 737.

Lorbeerblättrige Pflaume. W.

Habitat in Trapezunte, *unde in* Europam *venit* 1576. ♄ (v. v.)

*11. PRUNUS *paniculata.*

P. floribus paniculatis patulis, foliis ovatis. *Thunb. jap.* 200.

Riſpenblütige Pflaume. W.

Habitat in Japonia. ♄

Caulis *arboreus totus glaber.* Rami *et* Ramuli *erecti.* Folia *ſparſa petiolata ovata acute ſerrata venoſa glabra inaequalia bipollicaria et ultra.* Petioli *erecti lineam longi.* Flores *paniculati albi.* Panicula *ampla patentiſſima.* *Differt a P. Mahaleb, cui quodammodo ſimilis:* panicula *florum longe majori et patentiſſima,* floribus *minoribus,* foliis *magis oblongis, inferne attenuatis acuteque ſerratis.* Thunb. l. c.

12. PRUNUS *Mahaleb.*

P. floribus corymboſis terminalibus, foliis ovatis. *Iacq. auſtr. t.* 227. *Grantz. auſtr. p.* 95. *Scop. carn. ed.* 2. *n.* 588. *Pollich pal. n.* 466. *Hoffm. germ.* 169. *Roth. germ. I.* 211. *II.* 539.

P. floribus corymboſis terminalibus, foliis ſubcordatis petiolis glanduloſis. *Du Roi. harbk.* 2. *p* 182.

Ceraſus foliis ſubrotundis ſerratis, pedunculis multifloris. *Hall. helv. n.* 1084.

Ceraſus Mahaleb. *Mill. dict. n.* 4.

Ceraſus foliis ovatis. *Vir. cliff.* 43. *Hort. upſ.* 125. *Roy. lugdb.* 268. crenatis. *Hort. cliff* 487.
Ceraſus ſylveſtris amara, Mahaleb putata. *Bauh. hiſt. Du Ham. arb.* 1. *t.* 55.
Ceraſo affinis. *Bauh. pin.* 451.
Mahaleb. *Cam. epit.* 91.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 740.
Mahaleb-Pflaume. W.
Habitat in Helvetia, Germania. ♄ (v. v.)
Folia conduplicata, ovata, obtuſa, crenata. Stipulae *crenatae dentibus fuſcis.* Corymbus *convexus.* Bracteae *breviſſimae, obtuſiſſimae, ſerratae.*

13. PRUNUS *Armeniaca.*

P. floribus ſeſſilibus, foliis ſubcordatis. *Sp. pl.* 679. *Du Roi. harbk.* 2. *p.* 169. *Willd. arb.* 243.
P. foliis ovato-cordatis. *Hort. cliff.* 186. *Hort. upſ.* 124. *Roy. lugdb.* 268.
Armeniaca. *Mill. dict. Blackw. t* 281.
Armeniaca mala ſ. Mala praecocia priſcorum. *Bauh. hiſt.* 1. *p.* 176. *Camer. epit.* 146.
Mala armeniaca majora. *Bauh. pin.* 442.
Mala armeniaca majora. nucleo dulci. *Bauh. pin.* 442. *Knorr. del.* 1. *t. A.* 2.
β. Malus armeniaca minor. *Bauh. pin.* 442.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 742.
Apricoſen-Pflaume. W.
Habitat in Oriente. ♄ (v. v.)
Folia vernantia convoluta.

14. PRUNUS *ſibirica.*

P. floribus ſeſſilibus, foliis ovatis acuminatis ſimplicibus ſerratis, petiolis eglanduloſis. W.
P. floribus ſeſſilibus, foliis ovato-oblongis. *Sp. pl.* 679. *Pall roſſ* 1. *p.* 15. *t.* 8. *
P. inermis foliis ex ovato-cordatis, longe mucronatis, fructu exſucco. *Gmel. ſib.* 3. *p.* 172.
P. foliis ſeſſilibus, foliis ovato-oblongis, petiolatis glanduloſis. *Du Roi. harbk.* 2. *p.* 170.
Armeniaca betulae folio et facie, fructu exſucco. *Amm. ruth.* 272. *t.* 29.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 744.
Sibiriſche Pflaume. W.
Habitat in Sibiriae *ulterioris montoſis.* ♄

*15. PRUNUS *daſycarpa.*

P. floribus ſeſſilibus, foliis ovatis acuminatis duplicato-ſerratis, petiolis glanduloſis. *Willd. arb.* 243. *

P. (*daſycarpa*) foliis ovatis acuminatis duplicato-ſerratis, petiolatis glanduloſis, pedunculis ſubſolitariis breviſſimis, drupa globoſa pubeſcente. *Ehrh. Beitr.* 6. *p.* 90.

Rauhe Pflaume W.

Habitat in Oriente? ♄ (v. v.)

Simili praecedenti, at diverſa foliis *duplicato-ſerratis,* petiolis *glanduloſis. Vulgo nota it hortis ſub titulo gallico: Abricot nois.* W.

16. PRUNUS *pumila.*

P. floribus ſubumbellatis, foliis anguſto-lanceolatis. *Mant.* 75. *Du Roi. harbk.* 2. *p.* 186. *Willd. arb.* 244.

P. ſylveſtris humilior, fructu rubro praecoci minore, radice reptatrice. *Gron. virg.* 2. *p.* 76.

Ceraſus (*canadenſis*) foliis lanceolatis glabris integerrimis, ſubtus caeſiis, ramis patulis. *Mill. ic. t.* 89. *f.* 2.

Ceraſus canadenſis pumila, oblonga anguſtoque folio, fructu parvo. *Du Hamel. arb.* 149. *n.* 17.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 745.

Niedrige Pflaume. W.

Habitat in Canada. ♄ (v. v.)

Frutex *ſtatura Amygdali nanae.* Radix *repens.* Rami *virgati, pupuraſcentes.* Folia *ſubtus glauca ſ. pallida.* Pedunculi *proprii, uniflori, terni e gemmis lateralibus.* H. U.

*17. PRUNUS *Chamaeceraſus.*

P. umbellis ſeſſilibus, foliis obovatis obtuſiuſculis glabris glanduloſo-ſerratis. W.

P. (*Chamaeceraſus*) pedunculis ab unico ad quinque, umbellis ſeſſilibus, foliis lanceolatis utrinque glaberrimis. *Jacq. ic. rar.* 1. *t.* 90. *Collect.* 1. *p.* 133. * *Hoffm. germ.* 170.

Prunus Ceraſus ϑ. pumila. *Sp. pl.* 679.

Ceraſus pumila. *Bauh. pin.* 450.

Chamaeceraſus. *Cluſ. hiſt. p.* 64.

β. Prunus (*fruticoſa*) umbellis ſeſſilibus, foliis obovatis glabris ſerratis, ſerraturis infimis glanduloſis. *Pall. roſſ.* 1. *p.* 19. *t.* 8. *B. Gmel. ſib.* 3. *p.* 171.

Stranchartige Pflaume. W.

Habitat in Auſtriae *alpibus* β *in* Sibiria *ad* Volgam. ♄ (v. ſ.)

Sibirica varietas differt magnitudine ſola foliorum, florum, fructumque. W.

18. PRUNUS *Ceraſus.*

P. umbellis ſubpedunculis, foliis ovato-lanceolatis glabris conduplicatis. *Mat. med.* 124. *Blackw. t.* 449. *Du Roi. harbk.* 2. *p.* 175. *Pollich p. n.* 467. *Moench. haſſ. n.* 403. *Crantz. auſtr.* 94. *Hoff. germ.* 170. *Roth. germ. I.* 211. *II.* 539.

Ceraſus vulgaris. *Mill. dict. n.* 1.

Ceraſus foliis glabris, ſerratis, ovato-lanceolatis, mucrone producto. *Hall. helv. n.* 1083.

Ceraſus foliis ovato-lanceolatis. *Vir. cliff.* 49. *Hort. upſ.* 125. *Roy. lugdb.* 267. *Du Ham. arb.* 2. *t.* 56.

α Ceraſus *caproniana,* ſativa rotunda rubra et acida. *Bauh. pin.* 449.

β. Ceraſus *roſea,* hortenſis, flore roſeo. *Bauh. pin.* 450.

γ. Ceraſus *plena* hortenſis, flore pleno. *Bauh. pin.* 450. *Mill. ic.* 89. *f.* 1.

δ. Ceraſa *dulcia,* alba dulcia. *Bauh. pin.* 450.

ε. Ceraſa *juliana,* carne tenera et aquoſa. *Bauh. pin.* 450.

ζ. Ceraſus *auſtera,* acidiſſima ſanguineo ſucco. *Bauh. pin.* 445.

ϑ. Ceraſus *avium,* racemoſa hortenſis. *Bauh. pin.* 450.

C. hortenſis. *Mill. dict. n.* 2.

ι. Ceraſus hortenſis, flore pleno. *Bauh. pin.* 450. *Mill. dict. n.* 3.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 746.

Saure Pflaume. W.

Habitat in Europa. ♄ (v. v.)

Folia *conduplicata.* Stipulae *dentatae.* Bracteae *trifidae, ſerratae: intermedia foliacea.* Gemmae *terminales foliosae, laterales floriferae.*

19. PRUNUS *avium.*

P. umbellis ſeſſilibus, foliis ovato-lanceolatis ſubtus pubeſcentibus conduplicatis. *Fl. ſuec.* 2. *p.* 474. *Hudſ. angl.* 187. *Mat. med.* 125. *Du Roi. harbk.* 2. *p.* 172. *Pollich. pal. n.* 468. *Ludw. ect. t.* 108. *Blackw. t.* 425. *Kniph. cent.* 5. *n.* 73. *Hoffm. germ.* 170. *Roth. germ. I.* 211. *II.* 540.

Ceraſus foliis ovato-lanceolatis, ſerratis, inferne ſubhirſutis, mucrone producto. *Hall. helv. n.* 1082.

Prunus Ceraſus. *Scop. carn.* 2. *n.* 587.

Ceraſus nigra. *Mill. dict. n.* 2.

Ceraſus ſylveſtris, fructu nigro, et rubro. *Bauh. hiſt.* 1. *p.* 220.

Ceraſus major ſ. ſylveſtris, fructu ſubdulci nigro colore inficiente. *Bauh. pin.* 450. *Du Ham. arb. fruit.* 1.

Ceraſus ſylveſtris, ſeptentrionalis anglica, fructu rubro parvo ſerotino. *Raj. hiſt.* 1539.

β. Prunus *duracina.*

Ceraſus craſſa carne dura. *Bauh. pin.* 450.

γ. Prunus *Bigarella.*

Ceraſus ſativa majora. *Bauh. pin.* 450.

Ceraſus major, fructu cordato magno. *Raj.* 1538.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 749.

Süſse Kirſche. W.

Habitat in Europa *borealiore.* ♄ (v. v.)

Folia *conduplicata, glandula* 1. *ſ.* 2 *in petioli apice.* Umbella *ſeſſilis, ſubtriflora:* Floribus *petiolatis.* Involucrum 4-*phyllum.*

Haec floret e ramis antepenultimi, Ceraſus *vero ex ultimi anni ramis.*

Ex urbe Ponti Ceraſa in Italiam, Romam transportata per Lucullum, ex victoria Attica Mithridato victo, ab urbe condita 680.

*29. PRUNUS *ſemperflorens.*

P. floribus racemoſis, calycibus ſerratis, foliis ovatis ſerratis baſi glanduloſis. *Willd. arb.* 247. *

Prunus (*ſemperflorens*) caule arboreſcente, foliis ovato-lanceolatis ſerratis conduplicatis glabris, petiolis eglanduloſis, racemis folioſis pendulis. *Ehrh. Beit.* 7. *p.* 132.

P. (*ſerotina*) racemis elongatis folioſis pendulis, foliis ovato-lanceolatis ſerratis glabris, petiolis eglanduloſis. *Roth. catalect.* 1. *p.* 58.

Ceraſus ſativa aeſtate continue florens et frugeſcens. *Du Ham. arb.* 1. *p.* 178. *Du Roi. harbk.* 2. *p.* 178.

Ceraſus fructu ſerotino, cum pediculo longiori foliato. *Tournef. inſt.* 626.

Ceraſus racemoſa. *Bauh. hiſt.* 1. *p.* 223.

Immerblühende Pflaume. W.

Habitat - - - - ♄ (v. v.)

Hybrida forte progenies nunc in hortis noſtris faciem ſuam ſervans. W.

21. PRUNUS *penſylvanica.*

P. umbellis ſubſeſſilibus aggregatis multifloris tandem

paniculaeformibus, foliis oblongo-lanceolatis acuminatis glabris baſi biglanduloſis. *Ait. Kew.* 2. *p.* 165. *Willd. arb.* 248. *

P. umbellis ſeſſilibus, foliis ovato-lanceolatis, ramulis pallide punctatis. *Lin. ſuppl.* 252.

Prunus lanceolata. *Willd. arb.* 240. *t.* 3. *f.* 3.

Penſylvaniſche Pflaume. W.

Habitat in America *ſeptentrionali.* ♄ (v. v.)

Facies *P. Ceraſi, ſed ramuli punctis albidis adſperſi ut in P. virginiana.* Folia *P. Ceraſi lanceolato-ovata ſerrulata laevia, baſi glandulis duabus ſaepe rubris.* Umbellae *ſeſſiles.* Flores *P. Ceraſi minores.*

P. lanceolata a me in arb. berol. l. c. deſcripta, eſt nil aliud, quam haec Pruni ſpecies; vidi hanc ſero autumno cum pedunculorum tantum rudimentis, et flores P. Padi forma et facie eſſe e narratione hortulani habui. In ſequenti vero anno florentem arborem vidi et mox errorem agnovi. Infloreſcentia vero hujus ſpeciei ſaepius elongata eſt, et formam fere racemi habet. W.

*22. PRUNUS *pygmaea.* W.

P. umbellis ſeſſilibus paucifloris, foliis ellipticis acutis baſi biglanduloſis glabris. *Willd. arb.* 248. *

Zwerg-Pflaume. W.

Habitat in America *boreali.* ♄ (v. v.)

Frutex *tri-ſ. quadripedalis inermis..* Folia *ovato-elliptica acutiuſcula utrinque glabra, baſi attenuata argute ſerrata, dente ultimo utrinque glandula inſtructo.* Umbellae *ſeſſiles quadri-ſ. quinqueflorae confertae.* Flores *magnitudine P. ſpinoſae.* Fructus *magnitudine piſi majoris, nigri parum ſucculenti.* W.

*23. PRUNUS *nigra.*

P. umbellis ſeſſilibus ſolitariis paucifloris, foliis deciduis ovatis acuminatis, petiolis biglanduloſis. *Ait. Kew* 2. *p.* 165.

Schwarze Pflaume. W.

Habitat in America *boreali.* ♄ (v. v.)

Diverſiſſima a praecedente. Folia *ovata valde acuminata inaequaliter argute ſerrata utrinque glabra* Umbellae *tri-ſ. quadriflorae.* W.

*24. PRUNUS *aſpera.*

P. floribus ſolitariis terminalibus, foliis ovatis ſerratis aſperis. *Thunb. jap.* 201.

Muk no ki. *Kaempf. amoen.* 799.
Scharfblättrige Pflaume. W.
Habitat in Japonia. ♄.

Rami *et* ramuli *teretes punctati:* Folia *alterna petiolata ovata acuminata ſerrata venoſa utrinque ſcabra, ſubtus pallida patentia bipollicaria.* Petiolus *vix unguicularis ſulcatus.* Flores *in ultimis ramulis terminales ſolitarii.* Drupa *magnitudine piperis globoſo-ovata ſtylo perſiſtente coerulea glabra.* Nux *triquetra dura ovata acutiuſcula.* Nucleus *albidus.* Fructus *edulis.* Folium *pagina ſuprema aſperum ad poliendum adhibetur.* Thunb. l. c.

*25. PRUNUS *japonica.*

P. pedunculis ſolitariis, foliis ovatis acuminatis glabris, ramis inermibus. *Thunb. jap.* 201.
Japaniſche Pflaume. W.
Habitat in Japonia. ♄

Caulis *fruticoſus totus glaber.* Rami *et* ramuli *alterni teretes cinerei rugoſi erecti.* Folia *alterna petiolata ovata acuminata duplicato-ſerrata utrinquè glabra, venoſa patentia pollicaria.* Stipulae *binae ſupra foliaceae ſetaceae longitudine petioli.* Flores *ſolitarii pedunculati.* Pedunculus *infra ramulos uniflorus unguicularis.* Drupa *magnitudine piſi.* Thunb. l. c.

*26. PRUNUS *glanduloſa.*

P. pedunculis ſolitariis, foliis oblongis glanduloſo-ſerratis, ramis inermibus. *Thunb. jap.* 202.
Drüſige Pflaume. W.
Habitat in Japonia. ♄

Caulis *fruticoſus ramuloſus, omnibus partibus glaber.* Rami *teretes ſtriati cinerei, ramulis ſenſim filiformibus.* Folia *ſparſa petiolata oblonga ſerrata, ſerraturis gladuloſis, patentia, ſubtus venoſo-reticulata unguicularia.* Petioli *foliis multoties breviores.* Flores *ſparſi ſolitarii pedunculati.* Pedunculi *penduli longitudine foliorum.* Calyx *ſubcampanulatus.* Corolla *incarnata. Differt a P. pumila,* foliis *acutis tenuiſſime ſerratis ſubtus concoloribus nec obtuſis ſubintegris et glaucis: a P. inciſa:* foliis *glanduloſo-ſerratis glabris.* Thunb. l. c.

*27. PRUNUS *inciſa.*

P. pedunculis ſolitariis, foliis ovatis inciſo-ſerratis villoſis, ramis inermibus. *Thunb. jap.* 202.

Tiefzähnige Pflaume. W.
Habitat in Japonia. ♄

Caulis *arboreus parum craſſus debilis orgyalis.* Rami et ramuli *alterni flexuoſi glabri cinerei inermes patentes.* Folia *ex apicibus ramulorum plura breviſſime petiolata ovata acute inciſo-ſerrata villoſa, vix unguicularia ſub floreſcentia dein excreſcentia.* Petioli *foliis duplo breviores.* Flores *ex apicibus ramulorum ſolitarii pedunculati incarnati.* Pedunculi *capillares glabri foliis duplo longiores.* Perianthium *cylindricum apice dilatatum glabrum ferrugineum corolla duplo brevius.* Thunb. l. c.

*28. PRUNUS *tomentoſa.*

P. pedunculis ſolitariis, foliis ovatis ſubtus tomentoſis. *Thunb. jap.* 203.
Filzblättrige Pflaume. W.
Habitat in Japonia. ♄

Caulis *arboreus ramoſus.* Rami *glabri erecti ramuloſi.* Ramuli *alterni patuli tomentoſi* Folia *alterna petiolata ovata acuta ſerrata venoſa ſupra villoſa ſubtus tomentoſa patentia ſubpollicaria.* Petioli *tomentoſi ſemilineam longi.* Flores *ſolitarii pedunculati.* Drupa *magnitudine piſi.* Thunb. l. c.

29. PRUNUS *domeſtica.*

P. pedunculis ſubſolitariis, foliis lanceolato-ovatis convolutis, ramis muticis. *Mat. med.* 125. *Du Roi. harbk.* 2. *p.* 160. *Hoffm. germ.* 170. *Roth. germ. I.* 212. *II.* 541.
P. inermis, foliis lanceolato-ovatis. *Hort. cliff.* 186. *Hort. upſ.* 124. *Roy. lugdb.* 268.
Prunus pruna offic. *Crantz. auſt. p.* 94.
P. foliis ſerratis, hirſutis, ovato-lanceolatis, floribus longe petiolatis. *Hall. helv. n.* 1079.
P. ſylveſtris fructu majore. *Vaill. pariſ.* 140.
Prunus. *Bauh. pin.* 443. *Raj. hiſt.* 1526. *Blackw. t.* 309.
β. Prunus *Damaſcena.*
Pruna majora dulcia et parva atro-coerulea. *Bauh. pin.* 443 *n.* 23. *Blackw. t.* 305. *Knorr. del* 1. *t. P.* 2.
γ. Prunus *hungarica.*
Pruna magna craſſa ſubacida. *Bauh. pin.* 443.

δ. Prunus *juliana.*
Pruna oblonga caerulea. *Bauh. pin.* 443.
ε. Prunus *pertigona.*
Pruna nigra, carne dura. *Bauh. pin.* 443.
ζ. Prunus *cerea.*
Pruna coloris cerae ex candido in luteum palleſcente. *Bauh. pin.* 443.
η. Prunus *acinaria.*
Pruna magna rubra rotunda. *Bauh. pin.* 443.
θ. Prunus *maliformis.*
Pruna rotunda flava dulcia mali amplitudine. *Bauh. pin.* 443.
ι. Prunus *auguſtana.*
Pruna auguſto matureſcentia minora et auſteriora. *Bauh. pin.* 443.
κ. Prunus *praecox.*
Pruna parva praecocia. *Bauh. pin.* 443.
λ. Prunus *cereola.*
Pruna parva ex viridi flaveſcentia. *Bauh. pin.* 443.
μ. Prunus *amygdalina.*
Pruna amygdalina. *Bauh. pin.* 443.
ν. Prunus *galatenſis.*
Pruneoli albi oblongiuſculi acidi. *Bauh. pin.* 443.
ξ. Prunus *Brignola.*
Pruna ex flavo rufeſcentia mixti ſaporis gratiſſima. *Bauh. pin.* 443.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 750.
Gemeine Pflaume. W.
Habitat in Europae *auſtralioris locis elevatis.* ♄ (v. v.)
Folia vernantia convoluta. Gemmae *florifera aphyllae.*

50. PRUNUS *inſiticia.*

P. pedunculis geminis, foliis ovatis ſubtus villoſis convolutis, ramis ſpineſcentibus. *Amoen. acad.* 4. *p.* 273.* *Hudſ. angl.* 186. *Blackw. t.* 305. *Hall. helv. n.* 1081. *Du Roi. harbk* 2. *p.* 165. *Hoffm. germ.* 170. *Roth. germ. I* 212. *II* 541.
P. ſylveſtris major. *Raj. hiſt.* 1528. *Du Ham. arb.* 2. *t.* 41.
Pruna ſylveſtria praecocia. *Bauh. pin.* 444.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 752.
Kriechen-Pflaume. W.
Habitat in Germania, Helvetia, Anglia. ♄ (v. v.).

*31. PRUNUS *ceraſifera.*

P. pedunculis ſolitariis, foliis ellipticis glabris, fructibus pendulis ramis ſubinermibus. *Willd arb.* 250. *

P. (*ceraſifera*) ramis ſubſpineſcentibus glaberrimis, foliis ellipticis glabris convolutis, pedunculis ſolitariis tenuiſſimis glaberrimis, calyce reflexo, drupa ſubgloboſa pendula. *Ehrh. Beitr.* 4. *p.* 17.

Prunus domeſtica ξ. myrobalan. *Sp. pl.* 680.

Prunus domeſtica c. *Du Roi. harbk.* 2. *p* 164.

P. fructu rotundo nigro - purpureo dulci. *Bauh. pin.* 444.

P. fructu majori, rotundo rubri. *Tournef. inſt.* 622.

Runde Pflaume, W.

Habitat in America *boreali.* ♄ (v. v.)

32. PRUNUS *ſpinoſa.*

P. pedunculis ſolitariis, foliis elliptico - lanceolatis ſubtus pubeſcentibus, fructibus rectis, ramis ſpinoſis. W.

P. pedunculis ſolitariis, foliis lanceolatis glabris, ramis ſpinoſis. *Hort. cliff.* 186. *Fl. ſuec.* 397, 432. *Mat. med.* 125. *Roy. lugdb.* 268. *Du Roi. harbk.* 2. *p.* 166. *Scop. carn. n.* 586. *Pollich. pal. n.* 469. *Du Ham. arb.* 2. *p.* 186. *Blackw. t.* 494. *Hoffm. germ.* 170. *Roth. germ. I.* 212. *II* 541.

P. ramis ſpinoſis pubeſcentibus, foliis ellipticis convolutis ſubtus villoſis, pedunculis ſolitariis ſubpubeſcentibus, calyce patente, drupa globoſa. *Ehrh. Beitr.* 4. *p.* 16.

P. Acacia. *Crantz. auſtr. p.* 193.

P. ſpinoſa, foliis glabris, ſerratis, ovato - lanceolatis, floribus breviter petiolatis. *Hall helv. n.* 1080.

P. ſylveſtris. *Bauh. pin* 444. *Tabern. ic.* 992.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 755.

Schlehen - Pflaume. W.

Habitat in Europae *collibus apricis* ♄ (v. v.)

*33. PRUNUS *proſtrata.*

P. pedunculis geminis, foliis ovatis inciſo - ſerratis eglandulosis ſubtus tomentoſo - albicantibus, caule proſtrato. *Billard. pl. ſyriac. p.* 15. *t.* 6. *

P. cretica montana minima humifuſa, flore ſuave rubente. *Tournef. cor.* 43.

Liegende Pflaume. W.

Habitat in Cretae *montibus, inque cacumine montis* Libani. ♄

Amygdalis habitu accedit, ſed nux laevis. W.

683. PLINIA.

Gen. plant. edit. Schreb. n. 851.

Cal. 5-ſ. 4-partitus. Petala 5 ſ. 4. *Drupa* ſupera, ſulcata.

1. PLINIA *crocea.*

P. floribus pentapetalis. *Mant.* 402.
P. pinnata. *Sp. pl.* 2. *p.* 735.
P. fructu coroceo odorato. *Plum. gen.* 9. *ic.* 225.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 713.
Safranfarbige Plinie. W.
Habitat in America. ♄

Folia *pinnata, abrupta, foliolis circiter* 12, *ovato-lanceolatis, integerrimis.* Flores *ſeſſiles, ſparſi per ramos nudos ſeniores.* Fructus *edulis.*

Planta adhuc obſcura a recentioribus Botanicis nondum obſervata; quae ſecundum characterem incompletum, a Plumiero datum, ſitu germinis ab Eugenia diverſa videtur. W.

984. CHRYSOBALANUS.

Gen. pl. ed. Schreb. n. 850.

Cal. 5-fidus. *Petala* 5. *Stylus* lateralis *Drupae* nux 5-ſulcata, 5-valvis.

1. CHRYSOBALANUS *Icaco.*

Chryſobalanus. *Iacq. amer.* 154. *t.* 94. *Mill. dict. n.* 1.
Frutex folio cotini craſſo in ſummitate deliquium patiente, fructu ovali coeruleo. *Catesb. car.* 1. *p.* 25. *t.* 25.?
Guaiera. *Marcgr. braſ. t.* 2. *c.* 4.
α. Icaco fructu ex albo rufeſcente. *Plum. gen.* 44. *ic.* 158.
β. Icaco fructu nigro. *Plum. gen.* 44.
γ. Icaco fructu purpureo. *Plum. gen.* 44.
Chryſobalanus (*purpurea*) foliis decompoſitis, foliolis ovatis integerrimis. *Mill. dict. n.* 2. R.
δ. Chryſobalanus fruticoſus, foliis orbiculatis alternis, floribus laxe racemoſis. *Brown. jam.* 250. *t.* 17. *f.* 5.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 756. *t.* 11. *f.* 2.
Weſtindiſche Icacopflaume. W.
Habitat in America *meridionali.* ♄ (v. ſ.)

Species *an una vel plures vix dixero, quam tantum unica mihi viſa δ, quae caule ſcabro non eſt, ſed* Folia

alterna, ſubrotunda, integerrima, ſubemarginata. Flores *paniculati*: Pedunculis *exterius rotundatis, interius planis.*

*985. ROBINSONIA.

Gen. pl. ed. Schreb. n. 852.

Cal. 5-dentatus. *Petala* 5. *Bacca* ſtriata 2-locularis, loculis 1-ſpermis. *Sem.* villoſa.

*1. ROBINSONIA *melianthifolia.* W.

Touroulia gujanenſis. *Aubl. guj.* 1. *p.* 492. *t.* 194. *
Honigblumenblättrige Robinſonie. W.
Habitat in ſylvis Gujanae. ♄

Arbor *magna.* Folia *alterna fere Melianthi majoris, impari pinnata, foliolis lanceolato acuminatis argute ſerratis venoſis, ſuperioribus baſi in petiolum communem decreſcentibus.* Stipulae *oblongae acutae parvae.* Flores *flavi minuti in panicula terminali.* Bacca *Ceraſi magnitudine edulis acidula et grati ſaporis.* W.

986. SONNERATIA.

Gen. pl. ed. Schreb. n. 853.

Cor. 6-fidus. *Petala* 6-lanceolata. *Bacca* multilocularis, loculis polyſpermis.

1. SONNERATIA *acida.*

S. acida. *Lin. ſuppl.* 252.
Rhizophora (*caſeolaris*) foliis ovatis obtuſis, floribus ſolitariis, fructibus orbiculatis depreſſis mucronatis. *Sp. pl.* 635.
Aubletia caſeolaris. *Gaert. ſem.* 1. *p.* 379. *t.* 78. *f.* 2.
Pagapate. *Sonnerat. it. nov. guin. p.* 16. *t.* 10. 11.
Manginm caſeolare. *Rumph. amb.* 1. *p.* 111. *t.* 73. 74.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 183.
Sänerliche Sonneratie. W.
Habitat in Moluccis *inque uliginoſis* Nova Guineae. ♄

Arbor. Folia *oppoſita ſubſeſſilia oblonga integerrima Hyperici.* Flores *terminales ſolitariis magni.* Petala *rubra.* Fructus *ſimilis Meſembryanthemi, ſucco acido veſiculoſo.*

D I G Y N I A.

987. CRATAEGUS.

Gen. plant. ed. Schreb. n. 854.

Cal. 5-fidus. *Petala* 5. *Bacca* infera diſperma.

1. CRATAEGUS *coccinea.*

C. ſpinoſa, foliis cordato-ovatis inciſo-angulatis glabris, petiolis calycibusque glandulosis floribus pentagynis. *Ait. Kew.* 2. *p.* 167. *Willd. arb.* 81.

C. foliis cordatis repando-angulatis ſerratis glabris. *Mill. dict. n.* 4. *Du Roi.* 1. *p.* 139.

C. foliis ovatis repando angulatis ſerratis glabris. *Hort. cliff.* 187. *Hort. upſ.* 126. *Gron. virg.* 54. *Roy. lugdb.* 272. *Du Hamel. arb.* 12.

Meſpilus foliis cordatis ovatis acuminatis, marginibus acute ſerratis, ramis ſpinoſis. *Mill. dict. t.* 170.

Meſpilus, apii folio, virginiana ſpinis horrida, fructu amplo coccineo *Pluk. alm.* 249. *t.* 46. *f.* 4.

Meſpilus ſpinoſa ſ. Oxyacantha virginiana maxima *Angl. hort.* 49. *t.* 13. *f.* 1.

Oxyacantha ſpina ſancta dicta. *Raj. hiſt.* 1799.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 763.

Rother Weiſsdorn. W.

Habitat in Virginia, Canada. ♄ (v. v.)

Rami *adſperſi punctis vagis.* Spinae *axillares: validae, patentiſſimae, ramorum e rudimentis.* Folia *cuneato-ovata, ſublobata, ſerrata.* Pedunculi *pubeſcentes, corymboſi.*

2. CRATAEGUS *cordata.*

C. ſpinoſa, foliis cordato-ovatis inciſo-angulatis glabris petiolis calycibusque eglanduloſis, floribus pentagynis. *Ait. Kew.* 2. *p.* 168. *Willd. arb.* 82. *

C. (*populifolia*) foliis ſubcordatis trilobis, lobis inciſo-ſerratis, utrinque laevibus, petiolis praelongis laevibus. *Walt. carol. p.* 147.

Meſpilus (*Phaenopyrum*) ſpinoſa, foliis cordatis tri-quinque-ſ. ſeptemfidis ſerratis glabris, floribus corymboſis, ſegmentis calycinis deciduis, fructibus depreſſo-globoſis late umbilicatis, nucibus apice denudatis. *Lin. ſuppl.* 254. *Ehrh. Beitr.* 2. *p.* 67.

Meſpilus (*cordata*) foliis cordato-ovatis acuminatis acute

ſerratis, ramis ſpinoſis. *Mill. dict. ic.* 119. *tab.* 179. *Du Roi. harbk.* 1. *p.* 615.

Meſpilus acerifolia. *Burgsdorf. Anleit. n.* 232.

Ahornblättriger Weiſsdorn. W.

Habitat in America *boreali* ♄ (v. v.)

Arbor *mediocris.* Rami *ſpinoſi albo-maculati.* Folia *cordata trifida, quinquefida ſ. ſeptemfida acuminata, acute ſerrata glabra magnitudine Betulae albae.* Petiolus *tenuiſſimus folio brevior.* Corymbus *compoſitus.* Brateae *ad baſin pedunculorum ſolitariae ſubulatae minimae deciduae.* Flos *Crataegi Oxyacanthae paulo minor.* Dentes *calycini breviſſimi obtuſi, fructu maturo decidui.* Styli *quinque.* Fructus *depreſſo-globoſus ſ. verticilliformis coccineus magnitudine Ribis rubri, umbilico laxo nudo.* Nuces 5 *oſſeae, apicibus umbilicum replentibus nudis.* Floreſcentia *ſerotina et ultima hujus generis.*

3. CRATAEGUS *viridis.*

C. foliis lanceolato-ovatis ſubtrilobis ſerratis glabris caule inermi. *Sp. pl.* 683.

Meſpilus inermis foliis oblongis integris acuminatis ſerratis parvis utrinque viridibus. *Gron. virg.* 163.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 764.

Grüner Weiſsdorn. W.

Habitat in Virginia. ♄

Stipulae ſemicordatae.

Varietas *C. coccineae. Gron. virg.* 76. Variant *certe hae. Americae ſeptentrionalis ſpecies plurimum ſpinis ſ. ſine ſpinis, Stylorum numero, foliorum figura et videntur in natali mixtae. Sed inermis ſunt C. Aria, torminalis et indica.*

Incerta ſpecies, denuo obſervationibus determinanda. W.

*4. CRATAEGUS *pyrifolia.*

C. ſpinoſa inermisve, foliis ovato-ellipticis inciſo-ſerratis ſubplicatis ſubhirtis, calycibus villoſiuſculis, foliolis lineari-lanceolatis ſerratis, floribus trigynis. *Ait. Kew.* 2. *p.* 168. *Willd. arb.* 83. *

C. (*Leucophleos*) inermis, foliis ovatis inaequaliter ſerratis, baſin verſus integerrimis, ſtylo ſemibifido. *Moench. weiſſenſt. p.* 31. *t.* 2.

Crataegus tomentoſa. *Du Roi. harbk.* 1. *p.* 183.

Crataegus edulis. *Loddiges catal. p.* 8.

Meſpilus laciniata. *Walt. carol. p.* 147 ?
Meſpidus (*Calpodendron*) ſpinoſa, foliis ovalibus acuminatis ſubangulatis ſerratis pubeſcentibus baſi cuneatis, pedunculis multifloris, laciniis calycinis pinnatifidis. fructibus urceolatis. *Ehrh. Beitr.* 2. *p.* 67.
Birnenblättriger Weiſsdorn. W.
Habitat in America *boreali.* ♄ (v. v.)

*5. CRATAEGUS *elliptica.*
D. ſpinoſa, foliis ellipticis inaequaliter ſerratis glabris, petiolis calycibusque glandulosis, baccis globoſis pentaſpermis. *Ait. Kew.* 2. *p.* 168.
Ovalblättriger Weiſsdorn. W.
Habitat in America *boreali.* ♄

*6. CRATAEGUS *glanduloſa.*
C. ſpinoſa, foliis obovato-cuneiformibus angulatis glabris nitidis, petiolis ſtipulis calycibusque glanduloſis, baccis ovalibus pentaſpermis. *Ait. Kew.* 2. *p.* 168. *Willd. arb.* 84. *. *Moench. weiſſenſt. p.* 31.
C. (*ſanguinea*) ſpinoſa, foliis ſeptangulis ſerratis, baſi productis, petiolis ſubmarginatis. *Pall. roſſ.* 1. *p.* 25. *t.* 11.
Meſpilus (*rotundifolia*) ſpinoſa, foliis ſubrotundis ſubangulatis ſerratis glabris ſplendentibus, corymbis multifloris, laciniis calycinis ſerrato-glanduloſis, drupis globoſis. *Ehrh. Beitr.* 3. *p.* 20.
Drüſiger Weiſsdorn. W.
Habitat in America *boreali*, *inque* Sibiria. ♄ (v. v.)
In hac ſpecie ſpinae *validae, in ſequentibus duabus tenues, apice fere incurvo, et in flava ſaepe foliolis parvis inſtructae.* Ait. l. c.

*7. CRATAEGUS *flava.*
C. ſpinoſa, foliis obovato-cuneiformibus angulatis glabris nitidis, petiolis ſtipulis calycibusque glanduloſis, baccis turbinatis tetraſpermis. *Ait. Kew.* 2. *p.* 169.
Meſpilus (*flexiſpina*) foliis ovatis crenatis, ſtipulis ſemicordatis ſerratis, ſpinis foliaceis, floribus ſolitariis, *Moench. weiſſenſt. p.* 62. *t.* 4.
Gelbfrüchtiger Weiſsdorn. W.
Habitat in America *boreali.* ♄

8. CRATAEGUS *parvifolia.*
C. ſpinoſa, foliis cuneiformi-ovatis inciſis ſerratis, foliolis calycinis lanceolatis inciſis longitudine fructus,

floribus pentagynis. *Ait. Kew.* 2. *p.* 169. *Willd. arb.* 85. *

C. (*tomentoſa*) foliis cuneiformi-ovatis ſerratis ſubangulatis ſubtus villoſis, ramis ſpinoſis. *Sp. pl.* 682.

C. (*uniflora*) foliis lanceolato-ovatis ſerratis ſubtus villoſis, ſtipulis ſemicordatis, floribus ſolitariis, calycibus foliaceis, ſpinis longiſſimis. *Du Roi. harbk.* 1. *p.* 184.

Meſpilus (*Xanthocarpus*) ſpinoſa, foliis ſubcuneiformibus crenatis, floribus ſolitariis, laciniis calycinis ſubfoliaceis inciſo-ſerratis longis reflexis fructibus ſubturbinatis punctato-verrucoſis. *Lin. ſuppl.* 254.

Meſpilus inermis foliis ovato-oblongis ſerratis ſubtus tomentoſis. *Gron. virg.* 55.

Meſpilus caroliniana apii folio, vulgari ſimilis major, fructu luteo. *Trew. ehret. t.* 17.

Meſpilus virginiana groſſulariae foliis. *Pluk. phyt. t.* 100. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 766.

Kleinblättriger Weiſsdorn. W.

Habitat in Virginia. ♄ (v. v.)

Fruticulus *rigidus ſpinoſus.* Folia *ſubcuneiformia crenata craſſiuſcula ſubtomentoſa ſesquipollicaria.* Petiolus *breviſſimus.* Pedunculus *ex apicibus ramulorum ſolitarius tomentoſus, longitudine germinis.* Bracteae 2-ſ. 3 *lineari-lanceolatae ſerratae diciduae.* Germen *tomentoſum.* Laciniae *calyciniae ſubfoliaceae lanceolatae inciſo-ſerratae reflexae germine longiores.* Styli *quinque.* Fructus *ſubturbinatus pallide luteus, verrucis nigrispunctatus, profunde umbilicatus, magnitudine drupae Pruni inſiticiae.* Nuces 5 *oſſeae.*

*9. CRATAEGUS *villoſa.*

C. foliis oblongis acuminatis villoſis ſerratis, florum umbella compoſita. *Thunb. jap.* 204. *

Haariger Weiſsdorn. W.

Habitat in Japonia. ♄

Caulis *fruticoſus.* Folia *alterna petiolata ovato-oblonga acuminata ſerrata venoſa villoſa bipollicaria.* Flores *terminales umbellati.* Umbella *compoſita.* Styli *duo ſtigmatibus capitatis.* Fructus *villoſus.* Thunb. l. c.

*10. CATRAEGUS *laevis.*

C. foliis ovatis acuminatis glabris ſerratis, florum umbella ſubſimplici. *Thunb. jap.* 204. *

Doldenblütiger Weiſsdorn. W.
Habitat in Japonia. ♄

Caulis *fruticoſus totus glaber.* Folia *e gemmis alternis* 3 *vel* 4 *petiolata ovata acuminata tenuiſſime ſerrata plana glabra, tenuia, patentiſſima inaequalia pollicaria.* Flores *terminales umbellati patentiſſimi.* Umbella *ſubſimplex, rarius radio unico trifido vel racemoſo.* Styli. 2 *ſtigmatibus capitatis.* Thunb. l. c.

*11. CRATAEGUS *glabra.*

C. foliis oblongis acutis glabris serratis, florum panicula compoſita. *Thunb. jap.* 205.*

Glatter Weiſsdorn. W.
Habitat in Japonia. ♄

Differt a binis praecedentibus: foliis *oblongis nec ovatis, ſed craſſioribus et ſempervirentibus,* petalis *baſi villoſis,* panicula *magna compoſita.* Thunb. l. c.

*12. CRATAEGUS *punctata.*

C. ſpinoſa inermisve, foliis obovato-cuneiformibus glabris ſerratis, calycibus ſubvilloſis, foliolis ſubulatis integris. *Ait. Kew.* 2. *p.* 169. *Willd. arb.* 86.* *Iacq. hort.* 1. *p.* 10. *t.* 28.

Crataegus Crus galli. *Du Roi. harbk.* 1. *p.* 195. *Wangenh. amer.* 52.

Meſpilus (*cuneifolia*) ſpinoſa, foliis ovatis cuneatis in petiolos decurrentibus obſolete angulatis ſerratis vix pubeſcentibus, ſupra glabris, ſubtus venoſo-rugoſis, corymbis multifloris, pedunculis apice incraſſatis, ſegmentis calycinis integerrimis reflexis, drupis ſubgloboſis punctato-verrucoſis florum magnitudine, umbilico depreſſo. *Ehrh. Beitr.* 3. *p.* 21.

Punctirter Weiſsdorn. W.
Habitat in America *boreali.* ♄ (v. v.)

Variat fructibus rubris et flavis. W.

13. CRATAEGUS *Crus galli.*

C. ſpinoſa, foliis obovato-cuneiformibus ſubſeſſilibus nitidis coriaceis, foliolis calycinis lanceolatis ſubſerratis, floribus digynis. *Ait. Kew.* 2. *p.* 170. *Willd. arb.* 87.*

C. foliis lanceolato-ovatis ſerratis glabris, ramis ſpinoſis. *Kalm. it.* 1. *p.* 244.

Crataegus lucida. *Wangenh. amer. p.* 53. *t.* 17. *f.* 42.
Meſpilus lucida. *Ehrh. Beitr.* 4. *p.* 17.

Meſpilus foliis lanceolatis ſerratis, ſpinis robuſtioribus floribus corymboſis. *Mill. dict. t.* 178. *f.* 2.

a. ſplendens foliiis obovato-cuneiformibus. *Ait. l. c.*
Meſpilus aculeata pyrifolia denticulata ſplendens fructu inſigni rutilo, virginienſis. *Plum. alm.* 249. *t.* 46. *f.* 1. *Du Ham. arb.* 17. *Hort. angl. t.* 13.
Meſpilus pruni foliis, ſpinis longiſſimis fortibus, fructu rubro magno. *Clayt. virg.* 55.

β. pyracanthifolia foliis oblongo-lanceolatis ſubcuneiformibus. *Ait. l. c.*

γ. ſalicifolia foliis lanceolatis. *Ait. l. c.*
Houttuyn Lin. Pfl. Syſt. 1. *p.* 765.
Glänzender Weiſsdorn. W.
Habitat in Virginia. ♄ (v. v.)
Folia *ovalia, baſi cuneata, nitida.*

14. CRATAEGUS *indica.*
C. foliis lanceolatis ſerratis, caule inermi corymbis ſquamoſis. *Sp. pl.* 683.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 767.
Indiſcher Weiſsdorn. W.
Habitat in India. ♄
Arbor *magna ramis inermibus.* Folia *lato-lanceolata, obtuſe ſerrata, craſſiuſcula, petiolata.* Corymbi *terminales, pedunculis ſquamoſis, bracteis ſubulatis.*

15. CRATAEGUS *maura.*
C. foliis oblongis apice dentatis. *Lin. ſuppl* 253.
Mauritaniſcher Weiſsdorn. W.
Habitat in Mauritania. *S. M. Wänſtröm.* ♄
Statura et Facies C. Oxyacanthae. Frutex *cortice cinereo* Stipuleae *in meo nullae.* Folia *oblonga S. ſublanceolata obtuſa laevia apice tridentata S. quinquedentata, dentibus acutis.* Flores *in corymbis ramoſis ſimiles C. Oxyacanthae.*

16. CRATAEGUS *Oxyacantha.*
C. foliis obtuſis ſubtrifidis ſerratis glabris, floribus digynis, pedunculis calycibusque ſubglabriuſculis calycinis ſegmentis lanceolatis acutis. W.
C. foliis obtuſis ſubtrifidis ſerratis *Hort. cliff.* 188. *Fl. ſuec.* 399, 434. *Roy. lugdb.* 272. *Gmel ſib.* 3. *p.* 176. *Du Roi. harbk.* 1. *p.* 180. *Iacq. auſtr.* 3. *t.* 292. *f.* 2. *Pollich. pal. n.* 472. *Flor. dan. t.* 634. *Blackw. t* 149. *Kniph. cent.* 7. *n.* 17. *Knorr. del.* 1. *t. S. Medic. in*

obſ. Soc. Lutr. 1774. *p.* 294. *Hoffm. germ.* 171. *Roth. germ. I.* 213. *II.* 543.
Meſpilus ſpinoſa, foliis glabris, ſerratis, retuſis, trifidis. *Hall. helv. n.* 1087.
Meſpilus, apii folio, ſylveſtris ſpinoſa ſ. Oxyacantha. *Bauh. pin.* 454.
Oxyacantha ſ. Spina acuta. *Dod. pempt.* 751.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 768.
Gemeiner Weiſsdorn. W.
Habitat in Europae *pratis apricis duris.* ♄ (v. v.)

*17. CRATAEGUS *monogyna.*

C. foliis acutis ſubtrifidis, ſerratis glabris, floribus monogynis, pedunculis calycibusque ſubpubeſcentibus calycinis ſegmentis lanceolatis acuminatis. W.
C. (*monogyna*) foliis ſubtrifidis acutis floribus monogynis. *Murr. ſyſt. ed.* 14. *p.* 465. *Iacq. auſt.* 3. *t.* 292. *f.* 1.
C. (*monogyna*) rubra ſpinoſa, foliis quinquefidis ſerratis, lobis infimis divaricatis, ſtipulis ſemicordatis. *Pall. roſſ.* 1. *p.* 26.
β. C. (*monogyna nigra*) ſpinoſa, foliis quinquefidis inciſis cuneiformibus, ſtipulis lanceolatis. *Pall. roſſ.* 1. *p.* 26. *t.* 12.
γ. floribus ſimplicibus rubris.
δ. floribus plenis.
Einſamiger Weiſsdorn. W.
Habitat in Europae *dumetis, inque* Sibiria. ♄ (v. v.)

Folia *acuta profunde diviſa, et flores conſtanter monogyni,* pedunculi *calyeesque plerumque pubeſcentes denique* laciniae *calycinae acuminatae.* W.

*18. CRATAEGUS *pentagyna.*

C. foliis ovatis ſubtrifidis ſerratis, axillis venarum ſubtus villoſis, floribus pentagynis, pedunculis calycibusque pubeſcentibus, calycinis ſegmentis ſubrotundis. W.
Crataegus pentagyna. *Waldſt. et Kitaibel pl. hung.*
Fünfſamiger Weiſsdorn. W.
Habitat in comitatu Syrmienſi Hungariae *inque* Banatu. ♄ (v. ſ.)

Quoad foliorum formam ſimillima praecedenti, ſed diverſa: foliis *ſubtus ad venarum axillas villoſis,* calycinis *dentibus breviſſimis vix conſpicuis ſubrotundis obtuſis, denique* flore *conſtanter pentagyno.* W.

19. CRATAEGUS *Azarolus.*

C. foliis obtuſis ſubtrifidis ſubdentatis pubeſcentibus, ſegmentis calycinis ovatis. W.

C. foliis obtuſis ſubtrifidis ſubdentatis. *Sp. pl.* 683. *Mill. dict. n.* 7. *Willd. arb.* 90.*

Pyrus Azarolus. *Scop. carn.* 2. *n.* 597.

Meſpilus apii folio laciniato. *Bauh. pin.* 453.

Meſpilus Aronia veterum. *Bauh. hiſt.* 1. *p.* 67.

β. Crataegus *Aronia.*

Meſpilus orientalis, apii folio, ſubtus hirſuto. *Pock. orient.* 189. *t.* 85.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 770.

Azarol-Weiſsdorn. W.

Habitat Florentiae, Monſpelii, *in* Carniolia. ♄ (v. v.)

Differre videtur ab Oxyacantha *ſola magnitudine, ut Sorbus domeſtica a ſylveſtri; in hac quadruplo majora omnia; ſpinae in culta pauciores et ſaepe nullae.* Locula *quina, ut in Pyro.* Scop. l. c. R.

*988. WALDSTEINIA. W.

Willd. Nov. Act. ſoc. Berol. 2. *p.* 105.

Cal. 10-fidus, laciniis alternis minoribus. *Petala* 5. *Semina* 2 obovata.

*1. WALDSTEINIA *geoides.* W.

W. geoides. *Willd. Nov. Act. ſoc. Nat. ſcrut. Berol.* 2. *p.* 106. *t.* 4. *f.* 1.*

Geumartige Waldſteinie. W.

Habitat in ſylvis umbroſis Hungarie. ♃ (v. f.)

Folia *radicalia petiolata quinqueloba nervoſo-venoſa ſubpiloſa, lobis obtuſis obſolete trilobis dentatis.* Caulis *adſcendens foliorum longitudine teres ſtriatus piloſiuſculus.* Folia *caulina triloba inciſo-dentata.* Stipulae *oblongae acutae integerrimae.* Pedunculi *bini vel terni terminales filiformes longiſſimi.* Flores *flavi.* Planta *Geo affinis, ſed* piſtillorum *numero,* ſtylisque *deciduis clavatis diſtincta, a Potentillis vero habitu* piſtillorum *numero, et* ſtylorum *forma diverſiſſima.* W.

TRIGYNIA.

989. SORBUS.

Gen. plant. edit. Schreb. n. 855.

Cal. 5-fidus. *Petala* 5. *Bacca* infera, 3-ſperma.

1. SORBUS *aucuparia*.

S. foliis pinnatis utrinque glabris. *Mat. med.* 126. *Hall. helv. n.* 1091. *ſub* Meſpilo. *Mill. dict. n.* 1. *Gmel. ſib.* 3. *p.* 178.. *Gunn. norv. n.* 99. *Pollich. pal. n.* 473. *Du Roi. harbk.* 2. *n.* 420. *Blackw. t.* 73. *Willd. arb.* 365. *Hoffm. germ.* 171. *Roth. germ. I.* 213. *II.* 544.

S. foliis pinnatis. *Hort. cliff.* 188. *Fl. ſuec.* 400, 435. *Roy. lugdb.* 274. *Gron. orient.* 151.

S. foliis pinnatis, fructu pomiformi 5-loculari. *Crantz. auſt. p.* 88. *t.* 1. *f.* 4.

Meſpilus floribus trigynis: foliis pinnatis, glabris. *Scop. carn.* 1. *p.* 584. *n.* 3. *ed.* 2. *n.* 593.

S. aucuparia. *Tournef. inſt.* 634. *Bauh. hiſt.* 1. *p.* 62. *Du Ham. arb.* 2. *t.* 73.

S. ſylveſtris, foliis domeſticae ſimilis. *Bauh. pin.* 415. *Raj. hiſt.* 1457.

S. ſylveſtris. *Cam. epit.* 161.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 772.

Gemeine Ebereſche. W.

Habitat in Europae *frigidioribus*, Libano. ♄ (v. v.)

Styli 3, 4 *totque Semina.* Póll. *Fructus* 5-*locularis pentaſpermus, licet aliqua ſemina abortiant.* Crantz. l. c. R.

Folia *ſunt ſubtus pubeſcentia, in varietate vero ex America boreali, quae forte ſpecie differt, utrinque glaberrima nec juniora pubeſcentia.* W.

2. SORBUS *hybrida*.

S. foliis ſemipinnatis ſubtus tomentoſis. *Linn. dec.* 6. *Oed. dan. t.* 30. *Willd. arb.* 366. *Hoffm. germ.* 171. *Roth germ. I.* 213. *II.* 545.

Crataegus fennica. *Fl. ſuec.* 2. *n.* 433. *γ.*

Houttuyn Lin. Pfl. Syſt. 1. *p.* 775.

Baſtard Ebereſche. W.

Habitat in Gotlandia, Thuringia. ♄ (v. v.)

Statura *et foliatura aliquatenus Pyri Ariae.* Flores, Corymbus *et* Piſtilla 3. *S. aucupariae.*

3. SORBUS *domeſtica.*

S. foliis pinnatis ſubtus villoſis. *Hall. helv. n.* 1092. *Mill. dict. n.* 2. *Du Roi. harbk.* 2. *p.* 423. *Hoffm. germ.* 171. *Roth. germ. I.* 214. *II.* 546.

S. foliis pinnatis, fructu pyriformi 5-loculari. *Crantz auſtr. p.* 87. *t.* 2. *f.* 3.

S. ſativa. *Bauh. pin.* 415. *Edw. av. t.* 211. *Raj. hiſt.* 1456. *Tournef. inſt.* 633. *Blackw. t.* 174.

S. legitima. *Cluſ. hiſt.* 1. *p.* 10. *Du Ham. arb.* 1.

S. domeſtica. *Matthiol. dioſc. L.* 1. *c.* 136. *Cam. epit.* 160. *Lob. ic.* 2. *p.* 106.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 775.

Zahme Ebereſche. W.

Habitat in Europae *calidioribus.* ♄ (v. ſ.)

Fructus magnitudine minoris pomi Mali. Grandaeva *primum frutificat.*

Loculi *quini, raro ſemine completi omnes.* Crantz. l. c. R.

990. SESUVIUM.

Gen. plant. edit. Schreb. n. 856.

Cal. 5-partitus, coloratus, *Petala* 0. *Capſ.* ovata, trilocularis, circumſciſſa, polyſperma.

1. SESUVIUM *Portulacaſtrum.*

Seſuvium. *Iacq. amer.* 155. *t.* 95. *Swartz obſ.* 204. *

Portulaca foliis lanceolatis convexis, pedunculis unifloris. *Sp. pl.* 1. *p.* 446. *Loefl. ic.* 191. *

Aizoon repens, foliis oblongis turgidis foliis ſeſſilibus ſingularibus ad alas. *Brown. jam.* 241.

Portulaca curaſſavica, anguſto longo lucidoque folio procumbens, floribus rubris. *Pluk. alm.* 303. *t.* 216. *f.* 1. *Herm. par.* 212. *t.* 212.

Portulaca marina latifolia, flore ſuaverubente. *Plum. ſpec.* 6. *ic.* 223. *f.* 2.

Portulaca aizoides maritima procumbens, flore purpureo. *Sloan. jam.* 88. *hiſt* 1. *p.* 204.

Halimus indicus. *Rumph. amb.* 5. *p.* 165. *t.* 72. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 96.
Portulacartiges Seſuvium. W.
Habitat in Indiae *maritimis.* ⊙ (v. ſ.)

P E N T A G Y N I A.

991. M E S P I L U S.

Gen. plant. edit. Schreb. n. 857.

Cal. 5-fidus. *Petala* 5. *Bacca* infera, 5-ſperma.

1. MESPILUS *germanica.*

M. inermis, foliis lanceolatis ſubtus tomentoſis, floribus ſeſſilibus ſolitariis. *Hort. cliff.* 189. *Hort. upſ.* 129. *Mat. med.* 127. *Roy. lugdb.* 270. *Mill. dict. n.* 2. *Du Roi. harbk.* 1. *p.* 409. *Pollich pal n.* 474. *Blackw. t.* 154. *Kniph. cent.* 2. *n.* 45. *Ludw. ect. t.* 38. *Hoffm. germ.* 172. *Roth. germ. I.* 214. *II.* 546. *Willd. arb.* 192. *Pall. roſſ.* 1. *p.* 29. *t.* 13. *f.* 1.

α. ſtricta foliis duplicato ſerratis. *Ait. Kew* 2 *p.* 172.

M. foliis elliptico-lanceolatis ſerratis, calycibus longiſſimis perſiſtentibus. *Hall. helv. n.* 1094.

M. foliis oblongis arguto ſerratis, bacca pyriformi truncata. *Crantz. auſtr.* 80.

M. germanica, folio laurino non ſerrato. *Bauh. pin.* 453.

Meſpilus. *Dod. pempt.* 801.

β. diffuſa foliis ſubintegris. *Ait. l. c.*

Meſpilus folio laurino, major. *Bauh. pin.* 453. *Du Ham. arb. frut.* 1. *t.* 3.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 778.

Gemeine Miſpel. W.

Habitat in Europa *auſtrali.* ♄ (v. v.)

Rami *villoſi.* Folia *ovali-lanceolata; ad apicem ſerrata, ſubvilloſa, Petiolis breviſſimis canaliculatis.* Calyces *terminales, hirſuti, bractea longitudine corollae.*

*2. MESPILUS *japonica.*

M. inermis, foliis obovatis apice ſerratis, ſubtus tomentoſis, racemis paniculatis terminalibus. W.

M. inermis, foliis oblongis apice ſerratis, ſubtus tomentoſis. *Thunb. jap.* 206.* *Banks.* 5. *Ic. Kaempf. t.* 18.

Bywà. *Kaempf. amoen. p.* 800.

Japaniſche Miſpel. W.
Habitat in Japonia. ♄

3. MESPILUS *Pyracantha.*

M. ſpinoſa, foliis lanceolato-ovatis crenatis, calycibus fructus obtuſis. *Hort. cliff.* 189. *Vir. cliff.* 44. *Roy. lugdb.* 271. *Mill. dict. n.* 3. *Scop. carn. n.* 596. *Du Roi. harbk.* 1. *p.* 413. *Med. in obſ. Soc. oecon. Lutr.* 1774. *p.* 227. *Willd. arb.* 192. *Pall. roſſ. p.* 29. *t.* 13. *f.* 2.
M. aculeata amygdali folio. *Tournef. inſt.* 642. *Du Ham. arb.* 7.
Oxyacantha Dioſcoridis ſ. Spina acuta, pyri folio. *Bauh. pin.* 454. *Raj. hiſt.* 1459.
Uva urſi. *Dalech. hiſt.* 164.
Rhamnus 3. Dioſcoridis. *Lob. ic.* 2. *p.* 182.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 182.
Immergrüne Miſpel. W.
Habitat in Galloprovinciae, Italiae *ſepibus.* ♄ (v. v.)

4. MESPILUS *Chamae-Meſpilus.*

M. inermis, foliis ovalibus acute ſerratis glabris, floribus corymboſo-capitatis. *Sp. pl.* 685. *Scop. carn. n.* 592. *Mill. dict. n.* 8. *Willd. arb.* 192. *
M. foliis ovatis ſerratis glabris. *Hall. helv. n.* 1090.
Crataegus foliis ovalibus acute ſerratis, glabris, caule inermi. *Iacq. vind.* 243. *auſtr. t.* 231.
Crataegus follis ovalibus utrinque glabris plicatis indiviſis ſerratis. *Hort. cliff.* 497. *Vir. cliff.* 43. *Roy. lugdb.* 272.
Sorbus fruticoſa, foliis oblongo-ovalibus ſerratis, fructu ovato biloculari. *Crantz. auſtr. p.* 83. *t.* 1. *f.* 3.
Cotoneaſter folio oblongo ſerrato. *Bauh. pin.* 452.
Cotoneaſter forte Geſneri. *Cluſ. hiſt.* 1. *p.* 63.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 652.
Niedrige Miſpel. W.
Habitat in alpibus Auſtriacis, Pyrenaicis. ♄ (v. v.)

Dum tenera lanata, adultior nuda. Folia *obovata.* Dentes *calycis tubo breviores, in Amelanchier tubo longiores.* Flos *digynus viſus Hallero, Iacquinio, Scopolio.*

Folia *juniora glaberrima absque lanugine, quam tamen ſenſim induunt et in ſenectute iterum abiiciunt. Crantz. l. c.* R.

5. MESPILUS *Cotoneaſter.*

M. inermis, foliis ovatis integerrimis acutiuſculis ſubtus tomentoſis, germinibus glabris, baccis di-vel triſpermis. *Ait. Kew.* 2. *p.* 174. *Willd. arb.* 194. *

M. inermis, foliis ovatis integerrimis, ſubtus tomentoſis. *Oed. dan t.* 112. *Mill. dict. n.* 7. *Hall. helv. n.* 1093. *Hort. cliff.* 189. *Fl. ſuec.* 403, 438. * *Roy. lugdb.* 271. *Pollich. pal. n.* 476. *Leers. herb. n.* 375. *Scop. carn. n.* 594. *Kniph. cent.* 1. *n.* 58. *Hoffm. germ.* 172. *Roth. germ.* 1. 214. *II.* 548. *Pall. roſſ.* 1. *p.* 30 *t.* 14.

M. foliis ovato-acuminatis integerrimis ſubtus lanatis, bacca globoſa. *Crantz. auſtr. p.* 81.

Cotoneaſter folio rotundo non ſerrato. *Bauh. pin.* 452.

Chamae Meſpilus Geſneri. *Cluſ. hiſt.* 1. *p.* 60.

Chamae Meſpilus Cordi. *Bauh. pin.* 452.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 653.

Quitten-Miſpel. W.

Habitat in Europae *frigidioris colibus apricis, inque* Pyrenaeis, Sibiria *et monte* Ararat. ♄ (v. v.)

Styli *tres;* Bacca 3-*rarius* 4-*ſperma. Pollich l. c. Crantz.* Styli 3-4. Bacca 1-*locularis,* 3-*ſperma. Grimm. flor. iſen. in Nov. Act. A. N. C. III. app. p.* 321. Styli 3-5 *totque ſemina.* Scop. l. c. R.

*6. MESPILUS *tomentoſa.*

M. inermis, foliis ovatis integerrimis obtuſis ſubtus tomentoſis, germinibus lanatis, baccis pentaſpermis. *Ait. Kew.* 2. *p.* 174.

Meſpilus orientalis. *Mill. dict.*

Wollige Miſpel. W.

Habitat - - - - ♄

992. PYRUS.

Gen. plant. edit. Schreb. n. 858.

Cal. 5-fidus. *Petala* 5. *Pomum* inferum, 5-loculare, polyſpermum.

1. PYRUS *arbutifolia*

P. foliis obovatis acuminatis ſerratis, ſubtus tomentoſis, rachi ſupra glanduloſa, floribus corymboſis. *Willd. arb.* 257. *

P. (*arbutifolia*) inermis, foliis ovato-lanceolatis tenuiſ

ſime ſerratis, rachi ſupra glandulifere, corymbo compoſito. *Lin. ſuppl.* 256. *Wangenh. amer.* 89. *t.* 28. *f.* 64.

Meſpilus (*arbutifolia*) inermis, foliis lanceolatis crenatis ſubtus tomentoſis. *Sp. pl.* 685. *Hort. cliff.* 189. *Roy. lugdb.* 271. *Du Roi. harbk.* 1. *p.* 418.

Crataegus virginiana, foliis arbuti. *Tournef. inſt.* 633. *Mill. dict. t.* 109.

Sorbus virginiana, folio arbuti. *Herm. lugdb.* 578. *t.* 669.

Sorbus aucuparia virginiana foliis arbuti. *Breyn prodr.* 1. *p.* 15.

α. *rubra* fructu rubro.

β. *nigra* fructu nigro.

γ. *alba* fructu albo.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 782.

Sandbeerblättrige Birne. W.

Habitat in Virginia. ♄ (v. v.)

Fruticulus *inermis.* Folia *ovato-lanceolata interdum obovata ſ. cuneiformia, tenuiſſime ſerrata, ſubtus villoſa, ſeſqui-aut bipollicaria, adulta pagina ſuperiori rubra, inferiori lutea.* Rachis *ſupra tota glandulifera.* Glandulae *confertae ſubulato-cylindricae luteo-ruſeſcentes.* Serraturae *apice glanduloſo-cartilagineae.* Stipulae *laterales petiolo adnatae ſubulatae.* Corymbus *compoſitus tomentoſus* 6-12 *florus.* Bracteae *ſubulatae caducae.* Flores *vix magnitudine Crataegi Oxyacanthae.* Petala *obovata alba, extus rubella.* Antherae *rubrae.* Styli *quinque.* Frurtus *ſubgloboſus* 5-*locularis.* Semina *in quovis loculamento* 2, *oblonga fuſca.*

In varietate α *ſunt.* Folia *magis tomentoſa minora.* Fructus *rubri, magnitudine Ribis rubri, acerbi; in* β. Folia *minus tomentoſa.* Fructus *nigri magnitudine fructus Pyri Ariae.*

2. PYRUS *Botryapium.*

P. foliis oblongo-ellipticis cuſpidatis glabris, floribus racemoſis, petalis lineari-lanceolatis, germinibus pubeſcentibus, calycinis ſegmentis glabris. *Willd. arb.* 258.

P. (*Botryapium*) inermis, foliis, ovato-oblongis ſerratis acutis, racemis ſimplicibus elongatis. *Linn. ſuppl.* 255. *Wangenh. amer.* 90. *t.* 28. *f.* 65.

Meſpilus (*canadenſis*) inermis, foliis ovato-oblongis

glabris ſerratis acutiuſculis. *Sp. pl.* 685. *Mill. dict. n.* 6. *Du Roi. harbk.* 1. *p.* 416.
Meſpilus inermis, foliis ſubtus glabris obverſe ovatis. *Gron. virg.* 54. *Du Ham. arb.* 9.
Crataegus racemoſa. *Lamarck. encycl.* 1. *p.* 84.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 653.
Trauben-Birne. W.
Habitat in Virginia, Canada. ♄ (v. v.)

Arbor 10-*ſ.* 12-*pedalis inermis.* Folia *ovato-oblonga ſerrata acuta tripollicaria, tomentoſa, aldulta nuda glabra.* Racemi *ſimplices elongati ex apicibus ramulorum.* Pedunculi *tomentoſi.* Bracteae *ſubulato-filiformes villoſae caducae.* Petala *lineari-lanceolata obtuſa.* Styli *quinque.* Fructus *coeruleo-nigricans, magnitndine baccae* Ribis *nigri* 5-*locularis ſucculentus dulcis gratiſſimus.* Semina 10 *interdum pauciora brunea.* Gemmae *P. communis ſ. Populi tremulae, ſquamis interne villoſis.*

*3. PYRUS *ovalis.*

P. foliis ſubrotundo-ellipticis acutis glabris, floribus racemoſis, petalis obovatis, germinibus calycinisque ſegmentis pubeſcentibus. *Willd. arb.* 259. *
Meſpilus (*Amelanchier*) inermis, foliis ovato-lanceolatis crenatis, tenellis tomentoſis, adultioribus laevibus nitidis, floribus corymboſis. *Walt. carol. p.* 184.?
Crataegus ſpicata. *Lamarck. encycl.* 1. *p.* 84.
Rundblättrige Birne. W.
Habitat in America *boreali.* ♄ (v. v.)

Media inter praecedentem et ſequentem, differt a P. Botryapio: foliis *magnis ſubrotundis,* racemis *brevioribus,* calycinis *ſegmentis pubeſcentibus,* petalis *brevioribus obovatis, diſtincta a P. Amelanchier:* foliis *baſi ſubcordatis adultis glabris,* racemis *longioribus,* calycinis *ſegmentis pubeſcentibus,* petalis *obovatis ciliatis. Non eſt varietas ſequentis.* W.

4. PYRUS *Amelanchier.*

P. foliis ſubrotundo-ellipticis acutis ſubtus pubeſcentibus, floribus racemoſis, petalis lanceolatis, germinibus ſubvilloſis, calycinis ſegmentis glabris. W.
P. (*Amelanchier*) inermis, foliis ovalibus obtuſis ſerratis ſubtus tomentoſis calveſcentibus, ſtipulis lateralibus ſubulatis deciduis, racemo ſimplici paucifloro, petalis

ſublinearibus. *Lin. ſuppl.* 256. *Du Roi. harbk.* 2. *p.* 219.

P. foliis ovatis ſerratis ſubtus tomentoſis calveſcentibus *Hall. helv. n.* 1095.

Meſpilus (*Amelanchier*) inermis, foliis ovalibus ſerratis ſubtus hirſutis. *Sp. pl.* 685. *Mill. dict. n.* 5. *ic.* 178. *f.* 1. *Pollich. pal. n.* 475. *Iacq. auſt.* 3 *p.* 55. *t.* 300. *Hoffm. germ.* 172. *Roth. germ. I.* 214. *II.* 547.

Meſpilus inermis, foliis ovalibus ſerratis acutis. *Hort. cliff.* 189. *Roy. lugdb.* 271.

Sorbus (*Amelanchier*) foliis ovalibus ſerratis carinatis fructu globoſo multiloculari. *Crantz. auſtr. p.* 90.

Alni effigie lanato folio minor. *Bauh. pin.* 452.

Vitis idaea. *III. Cluſ. hiſt.* 1. *p* 62.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 651.

Felſen-Birne. W.

Habitat in Germaniae, Helvetiae, Auſtriae, Galliae, *rupeſtribus.* ♄ (v. ſ.)

Frutex *orgyalis inermis.* Ramuli *praeteriti anni epidermide alba ſoluta tecti.* Folia *ovalia obtuſa ſerrata, juniora ſubtus valde tomentoſa, adulta calveſcentia, vix pollicaria.* Stipulae *laterales petiolo adnatae ſubulatae deciduae.* Racemus *ſimplex* 4-*ſ.* 5-*florus tomentoſus.* Pedunculi *bracteati.* Bracteae *ſubulato-filiformes tomentoſae caducae.* Flores *Botryapii, interiores exteriores paulo ſuperantes.* Petala *lineari-lanceolata obtuſa.* Styli *quinque.* Fructus *coeruleo-nigreſcens magnitudine Ribis rubri* 5-*locularis ſucculentus dulcis ſapidus.* Semina 10, *in quovis loculamento* 2 *fuſca.* Gemmae *ovatae pubeſcentes rufae, ſquamis interne villoſis.*

Folia *ſero autumno tomentum exuunt.* W.

*5. PYRUS *cretica.*

P. foliis ſubrotundis emarginatis mucronatis ſubtus lanatis, floribus racemoſis, petalis lanceolatis, germinibus, calycinisque ſegmentis lanato-tomentoſis. W.

Meſpilus cretica folio circinato et quaſi cordiformi. *Tournef. cor.* 43.

Ceraſus Idea. *Alpin. exot. p.* 3. *t.* 2.

Candiſche Birne. W.

Habitat in monte Ida Cretae. ♄ (v. ſ.)

Simillima praecedenti, ſed folia fere orbiculata emarginata mucronata, groſſius dentata ſubtus magis

villoſa, pedunculi germina *et* laciniae *calycis denſo tomento ſunt tectae.* Fructus *magnitudine Peperis quinquelocularis, loculis diſpermis, et calycinis dentibus lanatis coronati.* W.

6. PYRUS *communis.*

P. foliis ovatis ſerratis, pedunculis corymboſis. *Ait. Kew.* 2. *p.* 174. *Willd. arb.* 260.

P. foliis ſerratis laevibus, floribus corymboſis. *Sp. pl.* 686. *Pollich. pal. n.* 477. *Du Roi. harbk.* 2. *p.* 207. *Scop. carn. n.* 598. *Blackw t.* 453. *Hoffm. germ.* 173. *Roth. germ. I.* 215. *II* 548.

P. foliis ovato-lanceolatis, ſerratis, glabris. *Hall. helv. n.* 1096.

P. foliis ſerratis, pomis baſi productis. *Hort. cliff.* 190. *Hort. upſ.* 130. *Fl. ſuec.* 401, 436. *Roy. lugdb.* 266.

Sorbus (*Pyrus*) foliis ſerratis, fructu turbinato 5-loculari. *Crantz. auſtr. p.* 93.

α. Pyrus *Pyraſter.*

P. ſylveſtris. *Bauh. pin.* 439. *Dod. pempt.* 351.

β. Pyrus *falerna.*

P. Bergamotta gallis *Bauh. hiſt.* 1. *p.* 44.

γ. Pyrus *pompejana.*

P. boni criſtiani. *Bauh. hiſt.* 1. *p.* 44.

δ. Pyrus *favonia.*

P. jeſu ſ. moſchatellina rubra. *Bauh. hiſt. p.* 1. 44.

ε. Pyrus *Volema.*

Pyra dorſalia eademque liberalia dicta. *Bauh. hiſt.* 1. *p.* 53.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 784.

Gemeine Birne. W.

Habitat in Europa. ♄ (v. v.)

Varietates hujus 80 *videndae in Milleri dictionario Hortulan.*

7. PYRUS *Pollveria.*

P. foliis ſerratis ſubtus tomentoſis, floribus corymboſis. *Mant.* 244. *Hoffm. germ.* 173. *Roth. germ. I.* 215. *II.* 549.

Pyrus (*irregularis*) foliis ovatis acute ſerratis, floribus cymoſis. *Onomat. bot.* 7. *p.* 611. *Munchhauſ. hauſvat.* 5. *p.* 246. *Du Roi. harbk.* 2. *p.* 216. *Knoop. Pomolog.* 2. *p.* 38. *t.* 4. R.

P. pollvilleriana. *Bauh. hiſt.* 1. *p.* 59.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 786.

Hanbutten - Birne. W.
Habitat in Germania. *L. B. de Munchh.* ♄ (v. v.)

Folia *ovata, ſerrata, ſubtus tomentoſa, etiam pagina ſuperiore praeſertim ſecundum coſtam pubeſcentia.* Petioli *tomentoſi.* Corymbus *elongatus, tomentoſus.* Floribus *numeroſiſſimis, P. communi multo minoribus.*

8. PYRUS *nivalis.*

P. foliis integerrimis, floribus corymboſis. *Lin. ſuppl.* 253.
Pyrus nivalis. *Iacq. auſt.* 2. *p.* 4. *t.* 107.
Schnee - Birne. W.
Habitat in Auſtriae *alpibus.* ♄ (v. ſ.)

Arbor *ramulis craſſiuſculis.* Folia *elliptica ſ. ovalia petiolata integerrima ſubtus ſericea albidaque.* Flores *terminales corymboſi.* Fructus *globoſus acerbiſſimus, ſed maturus putreſcens dulciſſimus.* Iacquin.

9. PYRUS *Malus.*

P. umbellis ſeſſilibus, foliis ovato-oblongis acuminatis ſerratis glabris, unguibus calyce brevioribus, ſtylis glabris. *Ait. Kew.* 2. *p.* 175. *Willd. arb.* 261.
P. foliis ſerratis, umbellis ſeſſilibus. *Mat. med.* 127. *Mill. illuſtr. Pollich. pal. n.* 478. *Du Roi. harbk. p.* 222. *Scop. carn. n.* 599. *Hoffm. germ.* 173. *Roth. germ. I.* 215. *II.* 549.
P. foliis ovatis acuminatis, ſubtus hirſutis, petiolis florigeris breviſſimis. *Hall. helv. n.* 1097.
P. foliis ſerratis; pomis baſi concavis. *Hort. cliff.* 189. *Hort. upſ.* 130. *Fl. ſuec.* 402, 437. *Roy. lugdb.* 266.
Sorbus (*Malus*) foliis ſerratis, fructu ſphaerico quinquelocutari. *Crantz. auſtr. p.* 93.
Malus (*ſylveſtris*) foliis ovatis ſerratis, caule arboreo. *Mill. dict. n.* 1.

α. Pyrus M. *ſylveſtris.*
Malus ſylveſtris. *Bauh. pin.* 443. *Dod. pempt.* 690. *Blackw. t.* 178.
β. Pyrus M. *paradiſiaca.*
Malus pumila, quae potius frutex quam arbor. *Bauh. pin.* 433.
γ. Pyrus Mal. *praſomila.*
Malus praſomila. *Bauh. pin.* 433.
δ. Pyrus M. *rubelliana.*

Malus ſativa, fructu ſanguinei coloris ex auſtero ſubdulci. *Tournef. inſt.* 635. *Blackw. t.* 141.

ε. Pyrus M. *ceſtiana.*

Mala curtipendula dicta. *Bauh. hiſt.* 1. *p.* 21.

ζ. Pyrus M. *cavillea.*

Mala ſativa, fructu magno intenſe rubente, violae odore. *Tournef. inſt.* 635.

η. Pyrus M. *epirotica.*

Poma orbiculata. *Ruell. ſtirp.*

Houttuyn Lin. Pfl. Syſt. 1. *p.* 787.

Apfel-Birne. W.

Habitat in Europa. ♄ (v. v.)

*10. PYRUS *dioica.*

P. foliis ovalibus ſerratis, floribus ſolitariis dioicis, petalis linearibus longitudine calycis. *Willd. arb.* 263.*

P. foliis ovalibus ſerratis, floribus ſolitariis dioicis. *Willd. phytog. faſc.* 1. *p.* 8. *n.* 27. *Moench. weiſſenſt. p.* 87. *t.* 8.

Pyrus apetala. *Münchh. Hausv.* 5. *p.* 247.

Malus non florens, fructificans tamen. *Bauh. pin.* 435.

Grünblühende Birne. W.

Habitat - - - - ♄ (v. v.)

Forte a praecedente orta. Folia *ſubtus pubeſcentia.* Pedunculi *uniflori in foliorum axillis ad ramorum apicem formant fere umbellam.* Petala *flaveſcenti-viridia linearia longitudine laciniarum calycis.* Stamina *nulla vidi.* Styli 5 *filiformes glabri.* Stigmata *clavata.* W.

*11. PYRUS *ſpectabilis.*

P. umbellis ſeſſilibus, foliis ovali-oblongis ſerratis laevibus, unguibus calyce longioribus, ſtylis baſi lanatis. *Ait. Kew.* 2. *p.* 175. *Willd. arb.* 263. *.

Prächtige Birne. W.

Habitat in China. ♄ (v. v.)

*12. PYRUS *prunifolia.*

P. umbellis ſeſſilibus, pedunculis pubeſcentibus, ſtylis baſi lanatis, foliis ovatis acuminatis. *Willd. phytogr. faſc.* 1. *p.* 8. *n.* 26. *Willd. arb.* 264.*

Pyrus Malus β. *Ait. Kew.* 2. *p.* 175.

Crataegus ceraſi folio, floribus magnis. *Mill. ic.* 180. *t.* 269.

Pflaumenblättrige Birne. W.

Habitat in Sibiria? ♄ (v. v.)

13. PYRUS *baccata.*

P. foliis aequaliter ſerrulatis, pedunculis confertis, po-

mis baccatis, calycibus deciduis. *Ait. Kew.* 175. *Willd. arb.* 264.
P. foliis ſerratis, pedunculis confertis, pomis baccatis. *Mant.* 75. *Giſecke ic. faſc.* 1. *n.* 12. *Pall. it.* 3. *p.* 105.
P. foliis ſerratis, umbellis pedunculatis, calycibus deciduis. *Pall. roſſ.* 1. *p.* 23. *t.* 10. *
Crataegus cerafi foliis, floribus magnis. *Amm. ruth.* 274. *t.* 31.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 789.
Beerentragende Birne. W.
Habitat in Sibiria, Dauria *ad* Schilkam *fluvium* ♄ (v. v.)

Arbor *ramis laevibus.* Folia *Corni, ovata, argute ſerrata, utrinque laevia.* Petioli *longitudine folii, baſi annectentes* Stipulas 2, *lineari-lanceolatas.* Pedunculi 4, *filiformes, laeves, axillares, conferti, ſimpliciſſimi, uniflori, longitudine petiolorum.* Germen *inferum, globoſum, glabrum, parvum.* Petala 5, *ſeſſilia, ovalia, concava, alba, majuſcula.* Stamina 20, *piſtillis breviora; Antheris flavis.* Piſtilla 5, *corolla breviora.* Bacca *ſubrotunda, 5-locularis, rubra.* Semina *bina, calloſa.* Calyx *minutus, deciduus.*

Obſ. *Ad* Meſpilum *referrem ob baccas, cum vero* Semina *in quovis loculo* 2. *ſint: nec ſolitaria,* Pyri *ſpecies erit, vel nulla diſtinctio inter Pyrum et Meſpilum.* Fructus *figura, conſiſtentia, colore, ſapore Mali aciduli, ſed minutus. Sero matureſcit.* H. U.

14. PYRUS *coronaria.*
P. foliis cordatis inciſo-ſerratis angulatis glabris, pedunculis corymboſis. *Ait. Kew.* 2. *p.* 176. *Willd. arb.* 265. *
P. foliis ſerrato-angulatis, umbellis pedunculatis. *Sp. pl.* 687. *Du Roi. harbk.* 2. *p.* 229.
Pyrus foliis ſerrato-anguloſis. *Sp. pl.* 1. *p.* 480. *Kalm. it.* 3. *p.* 10.
Malus coronaria. *Mill. dict. n.* 2.
Malus ſylveſtris, floribus odoratis. *Gron. virg.* 55. *Du Ham. arb.* 1.
Houttuyn Lin. Pfl. Syſt. 1. *p.* 790.
Wohlriechende Birne. W.
Habitat in Virginia. ♄ (v. v.)

Umbella *pedunculata.* Pedunculi *glabri.* Calyces *glabri, interne tomentoſi.* Folia *Mali, ſed magis glabra et argutius ſerrata.*

*15. PYRUS *anguſtifolia.*

P. foliis lanceolato-oblongis nitidis dentato-ſerratis baſi attenuatis integris, pedunculis corymboſis. *Ait. Kew.* 2. *p.* 176.

Pyrus coronaria. *Wangenh. amer.* 61. *t.* 21. *f.* 47.

Schmalblättrige Birne. W.

Habitat in America *boreali.* ♄

*16. PYRUS *japonica.*

P. foliis cuneatis crenatis glabris, floribus ſolitariis. *Thunb. jap.* 207.*

Buke. *Kaempf. amoen.* 844.

Japaniſche Birne. W.

Habitat in Japonia. ♄

17. PYRUS *Cydonia.*

P. foliis integerrimis, floribus ſolitariis. *Hort. cliff.* 160. *Mat. med.* 127. *Roy. lugdb.* 267. *Kalm. it.* 3. *p.* 107. *Jacq. auſtr. t.* 342. *Ludw. ect. t.* 56. *Kniph. cent.* 6. *n.* 75. *Hoffm. germ.* 173. *Roth. germ. I.* 215. *II.* 550. *Du Roi. harbk.* 2. *p.* 231. *Willd. arb.* 266.*

Sorbus (*Cydonia*) foliis rotundis integerrimis, fructu lanato 5-loculari. *Crantz. auſt. p.* 93.

Malus cotonea ſylveſtris. *Bauh. pin.* 435.

β. Malus cotonea major. *Bauh. pin* 434.

γ. Malus cotonea minor. *Bauh. pin.* 434.

Cotonea et Cydonia. *Lob. hiſt.* 580.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 791.

Quitten-Birne. W.

Habiat ad ripas petroſas Danubii, *ad ſepes* Germaniae, *quaſi ſponte.* ♄ (v. v.)

Calyx *patens, ſerratus, longitudine corollae.*

Poma *Heſperidum antiquis.*

Variat fructu *oblongo, globoſo et pyriformi, ut et* magnitudine *foliorum. Haec et praecedens ſpecies conveniunt* loculamentis *fructus polyſpermis, ſed non genere differunt.* W.

18. PYRUS *ſalicifolia.*

P. foliis lineari-lanceolatis canis ſubtus albo-tomentoſis, floribus axillaribus ſolitariis ſubſeſſilibus. *Lin. ſuppl.* 255. *Ait. Kew.* 2. *p.* 176. *Pall. itin.* 3. *p.* 734. *t. N. f.* 3. *Pall. roſſ.* 1. *p.* 20. *t.* 9.*

P. ſylveſtris orientalis, folio oblongo incano. *Tournef. cor.* 43.

Weidenblättrige Birne. W.

Habitat in Sibiria *inter fluvios* Terec *et* Cumam, *in promontario* Caucaſi *inque* Armenia. ♄

Arbor *primo vere florens.* Fructus *habet pyriformes turbinatos parvos.*

19. PYRUS *Aria.*

P. foliis ſubrotundo-ovatis inciſo-dentatis ſubtus albido-tomentoſis, floribus corymboſis. *Willd. arb.* 267.*

Pyrus (*Aria*) inermis, foliis ovalibus duplicato-ſerratis tomentoſis, corymbis multifloris planis, pomis ſubgloboſis. *Ehrh. Beitr.* 4. *p.* 20.

Crataegus (*Aria*) foliis ovatis inciſis ſerratis ſubtus tomentoſis. *Sp. pl.* 681. *Hort. cliff.* 187. *Fl. ſuec.* 398, 433. *Virid. cliff.* 43. *Mat. med.* 126. *Roy. lugdb.* 271. *Sauv. monſp.* 306. *Fl. dan. t.* 302. *Mill. dict n.* 1. *Du Roi. harbk.* 1. *p.* 190. *Pollich. palat.* 470. *Hoffm. germ.* 171. *Roth germ. I.* 212. *II.* 542.

Meſpilus foliis ovatis ſerratis, ſubtus tomentoſis. *Hall. helv. n.* 1089.

Meſpilus Aria. *Scop. carn. n.* 591.

Sorbus (*Aria*) foliis ovalibus ſerratis, facie ſplendentibus dorſo tomentoſis, fructu biloculari. *Crantz. auſt.* I. *faſc.* 1. *t.* 2. *f.* 2.

Alni effigie lanato folio major. *Bauh. pin.* 452.

Aria. *Dalech. hiſt.* 202.

Sorbus alpina. *Bauh. hiſt.* 1. *p.* 65.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 759.

Mehl Birne. W.

Habitat in nemoribus Europae. ♄ (v. v.)

20. PYRUS *intermedia.*

P. foliis ovato-lanceolatis inciſo-lobatis dentatis ſubtus niveo-tomentoſis, floribus corymboſis. *Willd. arb.* 268.*

Pyrus (*intermedia*) inermis, foliis ovalibus inciſo-lobatis ſerratis, ſubtus tomentoſis, corymbis multifloris planis, pomis ſubgloboſis. *Ehrh. Beitr.* 4. *p.* 20.

Crataegus Aria β. *Sp. pl.* 681.

Crataegus inermis, foliis ellipticis ſerratis transverſaliter ſinuatis, ſubtus villoſis. *Lin. lapp.* 199.

Oxel-Birne. W.

Habitat in Suecia, Ruſſia. ♄ (v. v.)

Toto habitu a praecedente diverſa. W.

21. PYRUS *torminalis.*

P. foliis cordato-ovatis laciniato-lobatis ſerratis, lobis

infimis divaricatis, floribus corymboſis. *Willd. arb.* 268.

Pyrus (*torminalis*) foliis ſubcordatis ſeptemlobatis, lobis infimis divaricatis. *Ehrh. Beitr.* 6. *p.* 92.

Crataegus (*torminalis*) foliis cordatis ſeptangulis, lobis infimis divaricatis. *Sp. pl.* 681. *Mill. dict. n.* 2. *Fl. dan. t.* 798. *Weig. rug. n.* 312. *Pollich. palat. n.* 471. *Du Roi. harbk.* 1. *p.* 187. *Hoffm. germ.* 171. *Roth. germ. I.* 213. *II.* 543. *Iacq. auſt.* 5. *p.* 21. *t.* 443.

Crataegus foliis cordatis acutis, lacinulis acutis ſerratis. *Hort. cliff.* 187. *Roy. lugdb.* 271. *Sauv. monſp.* 286.

Meſpilus foliis ſerratis ſeptilobis, lobis primis divergentibus. *Hall. helv. n.* 1088.

Sorbus (*torminalis*) folio ſeptangulo, fructu ovato-biloculari. *Crantz. auſt. p.* 85. *n.* 2.

Crataegus foliis laciniatis *Tournef. inſt.* 633. *Du Ham. arb.* 1. *t.* 79.

Meſpilus apii folio ſylveſtris non ſpinoſa, ſeu Sorbus torminalis. *Bauh. pin.* 454. *Raj. hiſt.* 1457.

Sorbus torminalis et Crataegus Theophraſti. *Bauh. hiſt.* 1. *p.* 63.

Sorbus torminalis. *Cam. epit.* 162. *Edw. av.* 212. *Blackw. t.* 173.

Sorbus torminalis Plinii. *Lob. ic.* 2. *p.* 100.

Houttuyn Lin. Pfl. Syſt. 1. *p.* 761.

Elzbeer-Birne. W.

Habitat in Germaniae, Auſtriae, Hungariae, Helvetiae, Angliae, Galliae *et* Orientis *ſylvis.* ♄ (v. v.)

*22. PYRUS *hybrida.*

P. foliis ſubtus pubeſcentibus pinnatis, pinna terminali maxima, pinnatifidis, ſimplicibusque, floribus corymboſis. *Willd. arb.* 269. *

Pyrus (*hybrida*) foliis pinnatis, pinna extima maxima ſubtus lanatis, floribus corymboſis. *Moench. weiſſenſt.* 90. *t.* 6.

Baſtard-Birne. W.

Habitat - - - - ♄ (v. v.)

Planta *hybrida matre P. arbutifolia et parte Sorbo aucuparia.* Folia *ſunt pinnata, pinnatifida et ſimplicia ſ. indiviſa, coſta media ſuperne glandulis eſt inſtructa.* W.

993. TETRAGONIA.

Gen. plant. ed. Schreb. n. 859.

Cal. 3-ſ. 5-partitus. *Petala* 0. *Drupa* infera, nuce 3-ſ. 8-loculari.

1. TETRAGONIA *fruticoſa.*

T. fruticoſa, foliis linearibus, fructibus alatis. *Ait. Kew.* 2. *p.* 176.

T. foliis linearibus. *Sp. pl.* 687. *Roy. lugdb.* 250. *Mill. dict. n.* 1. *et ic t.* 263. *f.* 2. *Hort. cliff.* 126. *Hort. upſ.* 126. *Kniph. cent.* 4. *n.* 87.

T. Tetragonocarpus africana rutilans, foliis longis et anguſtis. *Comm. hort.* 2. *p.* 205. *t.* 103.

Evonymo affinis africana, portulacae folio. *Seb. theſ.* 2. *p.* 13. *t.* 11. *f.* 8.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 655.

Strauchartige Viereckſfrucht. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Alae foliorum foliolis plurimis ſ. ramorum rudimentis. Flores *in racemum terminalem digeſti ſ. ſolitarii.*

*2. TETRAGONIA *decumbens.*

T. fruticoſa pruinoſa, foliis obovatis, fructibus alatis. *Ait. Kew.* 2. *p.* 177.

Tetragonia (*decumbens*) foliis ovatis integerrimis, caule fruticoſo decumbente. *Mill. dict. n.* 3. *Ic.* 176. *t.* 263. *f.* 1.

Liegende Viereckſfrucht. W.

Habitat ad Cap. b. ſpei. ♄

3. TETRAGONIA *herbacea.*

T. herbacea laevis, foliis ovatis petiolatis, fructibus alatis. *Ait. Kew.* 2. *p.* 177.

T. foliis ovatis. *Sp. pl.* 687. *Roy. lugdb.* 450.

Tetragonocarpos africana, radice magna craſſa et carnoſa. *Comm. hort.* 2. *p.* 203. *t.* 102.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 656.

Krautartige Viereckſfrucht. W.

Habitat ad Cap. b. ſpei. ♃.

Alae *ſaepius ſine ramorum rudimentis.* Pedunculi 3, *uniflori, ex alis foliorum.* Folia *obovato-lanceolata.*

4. TETRAGONIA *hirſuta.*

T. herbacea hirſuta procumbens, foliis ovatis villoſis, floribus axillaribus ternis ſeſſilibus. *Lin. ſuppl.* 258.

Haarige Viereckſfrucht. W.
Habitat ad Cap. b. ſpei. *Thunberg.*

5. TETRAGONIA *ſpicata.*

T. glabra herbacea erecta, foliis inferioribus ovatis, ſupremis lanceolatis glabris, floribus racemoſis. *Lin. ſuppl.* 258.
Aehrenblütige Viereckſfrucht. W.
Habitat ad Cap. b. ſpei. *Thunberg.*

*6. TETRAGONIA *echinata.*

T. herbacea, foliis rhombeo-ovatis, fructibus echinatis. *Ait. Kew.* 2. *p.* 177.
Stachliche Viereckſfrucht. W.
Habitat ad Cap. b. ſpei. ⊙ ♂ (v. v.)

Caulis *herbaceus prope radicem diviſus in* Ramos *diffuſos, petiolis decurrentibus angulatos, vix pedales.* Folia *ſucculenta patentia uncialia.* Petioli *foliis dimidio breviores.* Pedunculi *axillares ſolitarii filiformes, papulis nitidis tecti purpurei breviſſimi.* Flores *penduli, papulis cryſtallinis pruinoſi.* Calyx 3-ſ. 4-*phyllus, intus e viridi-flavicans.* Stamina *ſaepius* 3 *raro* 4. Germen *ſubtus planum triquetrum, anguli proceſſibus conicis pluribus echinati.* Styli *tres.* Nux *trilocularis.*

*7. TETRAGONIA *expanſa.*

T. herbacea, foliis ovato-rhombeis, fructibus quadricornibus. *Ait. Kew.* 2. *p.* 178.

T. (*expanſa*) herbacea, ramis expanſis elongatis, foliis parabolicis, floribus plerumque ſolitariis, fructu cornuto. *Murr. in comment. goett.* 1783. *p.* 13. *t.* 5.* *Scop. delic. inſub.* 1. *p.* 32. *t.* 14. *Thunb. Act. ſoc. Lin. Lond.* 2. *p.* 335.

T. (*halimifolia*) herbacea papuloſa, foliis elliptico-rhombeis petiolatis, pedunculis axillaribus unifloris ſubſolitariis, fructu cornuto. *Forſt. prod. n.* 223. *Forſt. pl. eſc. p.* 67. *n.* 37.*

T. (*halimifolia*) herbacea divaricata, foliis carnoſis alternis petiolatis ſubcordatis, floribus axillaribus ſolitariis ſubſeſſilibus. *Roth. Abhand. p.* 48. *t.* 8.*

T. (*japonica*) foliis ſubcordatis ovatis punctatis, floribus axilliaribus ſolitariis. *Thunb. jap.* 208.*

Demidovia tetragonoides. *Pall. hort. demidor. p.* 150. *tab.* 1.
Ausgebreitete Viereckſfrucht. W.

Habitat in Nova Zeelandia, *inſulis* Amicorum *et* Japonia. ⊙ (v. v.)

*8. TETRAGONIA *cryſtallina.*
T. herbacea pruinoſa, foliis ovatis ſeſſilibus, fructibus inermibus. *Ait. Kew.* 2. *p.* 178.
T. (*cryſtallina*) foliis ovatis cryſtallino-papilloſis, floribus axillaribus. *l'Herit. ſtirp.* 1. *p.* 81. *t.* 39. *
Cryſtallartige Vierecksfrucht. W.
Habitat in Peru. ⊙

994. MESEMBRYANTHEMUM.

Gen. pl. ed. Schreb. n. 860.

Cal. 5-fidus. *Petala* numeroſa, linearia, baſi cohaerentia. *Capſ.* carnoſa, infera, polyſperma.

* *Acaulia.*

1. MESEMBRYANTHEMUM *criniflorum.*
M. acaule, foliis petiolatis ovatis, ſcapis unifloris, corolla pentapetala. W.
M. foliis ovatis, ſcapis unifloris. *Lin. ſuppl.* 259.
Haarblütige Zaſerblume. W.
Habitat ad Cap. b. ſpei. *D. Montin.*

Planta *magnitudine Bellidis perennis.* Folia *ovato-oblonga petiolata, punctis cryſtallino-papilloſis.* Scapus *pollicaris uniflorus* Calyx *profunde quinquefidus, laciniis oblongis obtuſiſſimis.* Petala 5 *oblonga pedicellata calyce longiora alba.* Stamina *ſetacea alba longiſſima.*

*2. MESEMBRYANTHEMUM *ſpathulatum.*
M. acaule, foliis petiolatis ſpathulatis papuloſis, ſcapo uniflora. W.
Meſembryanthemum ſpathulatum. *Thunb.*
Spathelblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙ (v. ſ.)

Planta *parva uncialis.* Radix *ſimplex filiformis.* Folia *ſemiuncialia plana lanceolata ſpathulata obtuſa petiolata, utrinque papuloſa.* Scapus *papuloſus teres filiformis longitudine foliorum.* Flos *magnitudine Bellidis albus.* Calyx *campanulatus 5-fidus laciniis oblongis obtuſis, unica majore margine membranaceo.* Petala

numeroſa linearia leviſſime emarginata calyce duplo longiora. Filamenta *calyce breviora.* W.

*3. MESEMBRYANTHEMUM *teſticulare.*

M. acaule foliis quatuor decuſſatis inferne convexis ſuperne planis. W.

M. foliis quatuor decuſſatis ſupra planis. *Ait. Kew.* 2. *p.* 181. *Haworth. Meſemb.* 133. *n.* 17. *

Hodenförmige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♃

Corolla *alba.* W.

4. MESEMBRYANTHEMUM *linguiforme.*

M. acaule, foliis linguiformibus altero margine craſſioribus impunctatis, flore ſeſſili, calyce laevi, petalis emarginatis. W.

M. acaule, foliis linguiformibus altero margine craſſioribus impunctatis. *Sp. pl.* 699. *Hort. cliff.* 219. *n.* 8. *Hort. upſ.* 128. *Roy. lugdb.* 282. *Mill. dict. n.* 44. *Knorr. del.* 1. *t.* 5. *G. b. n.* 2.

M. (*ſcalpratum*) foliis ſcalpratiformibus latiſſimis, altero margine craſſioribus, baſi interiore cartilagineo-gibboſis calycibus glaberrimis, floribus maximis ſeſſilibus, petalis apice crenulatis. *Haworth Meſemb.* 187. *n.* 47. *

M. folio ſcalprato. *Dill. elth.* 235. *t.* 183. *f.* 224.

Ficoides afra acaulos, foliis aloes latiſſimis craſſiſſimis lucidis, flore aureo amplo ſine pedunculo. *Boerh. lugdb.* 1. *p.* 292.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 127.

Zungenblättrige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♃ (v. v.)

Folia *uno latere emarginata.* Flores *lutei.* W.

*5. MESEMBRYANTHEMUM *latum.*

M. acaule, foliis linguiformibus obtuſiſſimis altero margine craſſioribus impunctatis, flore ſubſeſſili, petalis acutis. W.

M. (*latum*) foliis linguiformibus breviuſculis latis aequalibus, floribus ſubſeſſilibus. *Haworth Meſemb.* 186. *n.* 46.

Meſembryanthemum linguiforme. *β.* *Sp. pl.* 699.

M. linguiformi folio latiore. *Dill. elth.* 236. *t.* 184. *f.* 224. *Knorr. del. t. G.* 7. *n.* 2.

Ficoides afra acaulos, foliis latiſſimis craſſiſſimis lucidis conjugatis, flore aureo amplo ſine pedunculo. *Boerh. lugdb.* 1. *p.* 292.

Breitblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃

Aetate ſub caulescens evadit. Folia *obtuſiſſima.* Flores *lutei.* W.

*6. MESEMBRYANTHEMUM *obliquum.* W.

M. acaule, foliis linguiformibus apice oblique acutis, coſta laciniarum calycis carinata ciliato-pubeſcente, pedunculo floris longitudine, petalis acutis. W.

M. (*linguiforme*) foliis linguiformibus latis obtuſis, junioribus anguſtis brevioribus, floribus breviſſime pedunculatis, calycibus laciniis ſubtuberculatis carinatis, carinis ciliato-pubeſcentibus. *Haworth Meſembr.* 182. *n.* 45.*

Meſembryanthemum linguiforme. γ. *Sp. pl.* 699.

M. folio linguiformi anguſtiori. *Dill. elth.* 238. *t.* 185. *f.* 226.

Ficoides afra acaulos, foliis latiſſimis craſſiſſimis lucidis conjugatis brevioribus, flore aureo amplo pedunculo brevi. *Boerh. lugdb.* 1. *p.* 292.

Schiefblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃

Flores lutei. W.

*7. MESEMBRYANTHEMUM *longum.*

M. acaule, foliis linguiformibus apice oblique acutis, coſta laciniarum calycis carinata glabra, pedunculis flore duplo longioribus, petalis acutis. W.

M. (*longum*) foliis linguiformibus longiſſimis ſubaequalibus, pedunculis elongatis. *Haworth Meſembr.* 177. *n.* 43.*

Meſembryanthemum. δ. *Sp. pl* 699.

M. folio linguaeformi longiore. *Dill. elth.* 238. *t.* 187. *f.* 227. *Knorr. del* 1. *t.* G. 7. *n.* 1.

Langblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃

Calycis *laciniae carinatae glabrae absque ciliis.* Petala *flava.* W.

8. MESEMBRYANTHEMUM *roſtratum.*

M. acaule, foliis ſemicylindricis connatis externe tuberculatis. *Sp. pl.* 696. *Hort. cliff.* 219. *n.* 23. *Roy. lugdb.* 285. *Mill. dict. n.* 40. *Kniph. cent.* 10. *n.* 61. *Haworth Meſembr.* 152. *n.* 29.*

M. roſtrum. ardeae referens. *Dill. elth.* 240. *t.* 186. *f.* 229.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 124.
Schnabelförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃
Flores *flavi.* W.

*9. MESEMBRYANTHEMUM *compactum.*
M. acaule, foliis connatis punctatis ſemiteretibus apice triquetris ſubreflexis acutis, floribus ſeſſilibus, calyce ſubcylindraceo ſexfido. *Ait. Kew.* 2. *p.* 191.
Dichtblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃
Flores *flavi.*

10. MESEMBRYANTHEMUM *caninum.*
M. acaule, foliis impunctatis oblongis apice triquetris et dentatis, dentibus obtuſis, floribus pedunculatis, pedunculis foliis multo longioribus. W.
M. (*caninum*) ſubacaule, foliis opacis apice triquetris, floribus pedunculatis, pedunculis longiſſimis baſi bracteatis. *Haworth Meſemb. p.* 159. *n.* 34. *
Meſembryanthemum ringens α caninum. *Sp. pl.* 698.
M. ſubacaule, foliis ciliato-dentatis. *Hort. cliff.* 218. *Mill. dict. n.* 39. *ic. t.* 77. *f.* 2.
M. rictum caninum referens. *Dill. elth.* 241. *t.* 188. *f.* 231.
Ficoides capenſis humilis, folio triangulo prope ſummitatem dentato, flore luteo. *Bradl. ſucc.* 2. *p.* 8. *t.* 17.
Hunderachenartige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃ (v. v.)

11. MESEMBRYANTHEMUM *felinum.*
M. acaule, foliis punctatis oblongis ſubtus convexis, apice dentatis, dentibus acuminatis, flore ſeſſili. W.
M. (*felinum*) acaule, foliis ſubtus ſemi-cylindricis ciliatis, ciliis longis. *Haworth Meſembr. p.* 161. *n.* 35. *
M. (*ringens*) ſubacaule, foliis ciliato-dentatis punctatis. *Sp. pl.* 698. *Roy. lugdb.* 293.
Meſembryanthemum ringens β. felinum. *Sp. pl.* 298.
M. rictum felinum repraeſentans. *Dill. elth.* 240. *t.* 87. *f.* 220. *Knorr. del.* 1. *t. G. b. n.* 1.
Ficoides afra, folio triangulari enſiformi craſſo brevi ad margines laterales multis majoribus ſpinis aculeato. *Mart. cent.* 30. *t.* 30.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 125.
Katzenrachenförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃ (v. v.)

Diverſiſſima a praecedente: foliis *punctatis*, *ſubtus convexis*, *nec apice triquetris*, dentibus *ad apicem acuminatis longioribus*, flore*que ſeſſili*, *qui et luteus eſt ut in praecedente.* W.

*12. MESEMBRYANTHEMUM *tigrinum.*

M. acaule, foliis maculatis oblongis ſubtus convexis apice ciliato-dentatis, flore ſeſſili. W.

M. (*tigrinum*) acaule foliis maculatis ciliatisque, ciliis longiſſimis. *Haworth Meſembr. p.* 164. *n.* 36. *

Getiegerte Zaſerblume. W.

Habitat ad Cap. b. ſpei, ♃

Corolla *magna lutea.* W.

*12. MESEMBRYANTHEMUM *murinum.*

M. ſubacaule, foliis punctatis oblongis baſi ſubtus convexis, apice triquetris, trifariam dentatis, flore ſubſeſſili. W.

M. (*murinum*) ſubacaule, foliis punctatis baſi ſemicylindricis attenuatis, apice triquetrisque trifariam denticulatis, floribus ſubſeſſilibus. *Haworth Meſembr. p.* 165. *n.* 37.

Mauſezahnartige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♃

Corolla *parva lutea.* W.

14. MESEMBRYANTHEMUM *albidum.*

M. acaule, foliis triquetris integerrimis. *Sp. pl.* 699. *Mill dict. n.* 45. *Haworth Meſembr. p.* 156. *n.* 32. *

M. foliis robuſtis albicantibus. *Dill. elth.* 243. *t.* 189. *f.* 232.

Ficoides africana, folio triangulari enſiformi glauco craſſo, flore amplo aureo. *Bradl. ſucc.* 5. *p.* 20. *t.* 43.

Ficoides africanum triangulari robuſtiſſimo folio. *Herm. prodr.* 172.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 126.

Weiſsblättrige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♃

15. MESEMBRYANTHEMUM *bellidiflorum.*

M. acaule, foliis triquetris linearibus impunctatis apice trifariam dentatis. *Sp. pl.* 690. *Hort. cliff.* 218. *n.* 13. *Roy. lugdb.* 283. *Mill. dict. n.* 9. *Knorr. del.* 1. *tab. n.* 1. *G. a. Haworth Meſembr. p.* 143. *n.* 27. *

Meſembryanthemum bellidiflorum. *Dill. elth.* 244. *t.* 189. *f.* 233.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 106.

Maſsliebartige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃
Variat corollis purpureis, rubicundis et albis. W.

* * *Subacaulia, caule breviſſimo.*

*16. MESEMBRYANTHEMUM *limpidum.*
M. ſubacaule, foliis ſcabrido-punctatis obovato-lanceolatis planis, pedunculis fructiferis reflexis, calycinis laciniis inaequalibus ſpathulatis. W.
M. (*limpidum*) foliis oppoſitis ſpathulatis obtuſis ſcabridis, papulis oblongis, foliolis calycinis oblongis obtuſis medio coarctatis. *Ait. Kew.* 2. *p.* 183. *Haworth Meſembr. p.* 116. *n.* 3.
Meſembryanthemum cuneifolium. *Iacq. ic. rar.* 3. *t.* 488. *Collect.* 2. *p.* 319.*
Keilblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙
Corolla *magna rubicunda.* W.

17. MESEMBRYANTHEMUM *Tripolium.*
M. ſubcauleſcens foliis impunctatis oblongis planis, calycibus pentagonis, laciniis calycinis ovatis aequalibus. W.
M. foliis alternis lanceolatis planis impunctatis, caulibus laxis ſimplicibus, calycibus pentagonis. *Sp. pl.* 690. *Hort. cliff.* 217. *n.* 11. *Hort. upſ.* 128. *Roy. lugdb.* 283. *Mill. dict. n.* 8. *Kniph. cent.* 8. *n.* 70. *Haworth Meſembr. p.* 122. *n.* 12.*
M. tripolii folio, flore argenteo. *Dill. elth.* 230. *t.* 179. *f.* 220.
Ficoides africanum procumbens tripolii ſucculento folio, flore candido. *Bradl. ſucc.* 5. *p.* 14. *t.* 47.
Ficoides africanum, Meſembryanthemum ſ. Ficus aizoides major procumbens, tripolii folio magis ſucculento denticulis fimbriato. *Pluk. mant.* 77. *t.* 329. *f.* 4.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 105.
Aſterblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♂
Corolla *mediocris alba.* W.

18. MESEMBRYANTHEMUM *dolabriforme.*
M. ſubacaule, foliis punctatis triquetro-carinatis, carina apice dilatata biloba. W.
M. acaule, foliis dolabriformibus punctatis. *Sp. pl.* 699. *Hort. cliff.* 219. *n.* 19. *Roy. lugdb.* 284. *Mill. dict. n.* 41. *Kniph. cent.* 1. *n.* 51. *Curt. mag.* 32.

M. (*dolabriforme*) foliis dolabriformibus punctatis, caule erecto. *Haworth Meſembr. p.* 366. *n.* 38.
M. folio dolabriformi. *Dill. elth.* 248. *t.* 191. *f.* 237.
Ficoides capenſis humilis, foliis cornu cervi referentibus, petalis luteis, noctiflora. *Bradl. ſucc.* 1. *p.* 11. *t.* 10.
β. M. (*dolabriformoides*) foliis ſubdolabriformibus punctatis, caule deflexo. *Haworth Meſembr. p.* 168. *n.* 29.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 125.
Hobelblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

19. MESEMBRYANTHEMUM *difforme.*
M. ſubacaule, foliis punctatis lanceolatis ſemiteretibus margine difforme cordatis, baſi connatis. W.
M. acaule, foliis difformibus punctatis connatis. *Sp. pl.* 699. *Roy. lugdb.* 287. *Mill. dict. n.* 42. *et ic. t.* 176. *f.* 2. *Haworth Meſembr. p.* 169. *n.* 40.
M. foliis difformibus flore luteo *Dill. elth.* 252. *t.* 194. *f.* 241. 242.
Ficoides Meſembryanthemum craſſimo et valde ſucculento folio nigris punctis notato. *Pluk. alm.* 148. *t.* 325. *f.* 4.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 126.
Ungeſtaltete Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

20. MESEMBRYANTHEMUM *calamiforme.*
M. ſubacaule, foliis impunctatis adſcendentibus ſubteretibus acutis, baſi connatis, floribus octogynis pedunculatis. W.
M. acaule, foliis ſubteretibus adſcendentibus impunctatis connatis, floribus octogynis. *Sp. pl.* 690. *Mill. dict. n.* 7. *et ic. t.* 176. *Haworth Meſembr. p.* 140. *n.* 24. *
M. calamiforme. *Dill. elth.* 229. *t.* 186. *f.* 228.
Ficoides capenſis humilis, cepaeae folio, flore ſtramineo. *Bradl. ſucc.* 2. *p.* 10. *f.* 19.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 105.
Federkielartige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Folia quamvis baſi connata recta, tamen ſaepe horizontaliter patentia, apice recurva, ſuperficie punctato-ſcabrida.

*21. MESEMBRYANTHEMUM *digitatum.*
M. ſubacaule, foliis alternis teretibus obtuſis, floribus axillaribus ſeſſilibus. *Ait. Kew.* 2. *p.* 181.

Fingerförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Corolla *lutea.* W.

* * * *Cauleſcentia aphylla.*

*22. MESEMBRYANTHEMUM *minimum.*
M. caule erecto clavato aphyllo maculato, apice pubeſcente, flore pedunculato. W.
M. (*minimum*) planta carnoſa obconica truncata maculata minima, maculis confluentibus. *Haworth Meſembr. p.* 126. *n.* 14. *
Kleinſte Zaſerblume. W.
Habitat - - - - ♄
Flos *luteus breve pedunculatus.* W.

* * * * *Cauleſcentia foliis planis.*

23. MESEMBRYANTHEMUM *pinnatifidum.*
M. foliis planis oblongis pinnatifidis. *Ait. Kew.* 2. *p.* 193, *Curt. mag.* 67.
M. foliis pinnatifidis. *Lin. ſuppl.* 260. *Haworth Meſembr. p.* 112. *n.* 1. *
Gefiederte Zaſerblume. W.
Habitat ad Cap. b, ſpei. ☉ (v. v.)
Caulis *herbaceus dichotomus craſſitie fili groſſioris depreſſus ſanguineus.* Folia *oppoſita oblonga lyrata ſ. pinnatifida.* Lobi laterales *utrinque tres obtuſi, ſinubus patulis obtuſis, non coſtam attingentibus.* Lobus terminalis *impar major acutior.* Baſis *foliorum hinc dilatata, tanquam orta e ſtipula lata.* Flores *axillares ramorum ſolitarii pedunculati parvi flavi.* Pedunculus *deflexus folio brevior.* Petala *lanceolata.* Fructus *turbinatus pentagonus, vertice plano articulato tamquam quinquevalvi. Tota planta adpreſſa micis veſicularibus, verſus autumnum tota rubra. Planta pulchra et in hoc genere valde ſingularis.*

24. MESEMBRYANTHEMUM *cordifolium.*
M. foliis oppoſitis petiolatis cordatis, calycibus quadrifidis, caule tereti. *Ait. Kew.* 2. *p.* 182. *Iacq. ic. rar.* 3. *t.* 487. *Collect.* 2. *p.* 294. * *Smith. ſpicil. bot.* 1. *p.* 5. *t.* 6. *
M. foliis cordatis obtuſis, caulibus proſtratis. *Lin. ſuppl.* 260. *Gloxin. obſ. bot. p.* 22. *t.* 1. *f. a. Haworth Meſembr. p.* 192. *n.* 49. *

Herzblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)

Caules *herbacei proſtrati pedales, utrinque ſulco exarati.* Folia *oppoſita, oppoſitionibus remotis, petiolata cordata obtuſa pollicaria.* Flores *laterales alterni ſolitarii pedicellati longitudine foliorum purpurei.* Germina *majuſcula.* Calyx *foliolis oppoſitis, majoribus ovatis, reliquis minoribus.*

25. MESEMBRYANTHEMUM *cryſtallinum.*
M. foliis alternis ovatis papuloſis, floribus ſeſſilibus, calycibus late ovatis acutis retuſis. *Lin. ſuppl.* 259. *Haworth Meſembr. p.* 113. *n.* 2. *
M. foliis alternis ovatis papuloſis undulatis. *Sp. pl.* 688. *Hort. cliff* 216. *n.* 1. *Hort. upſ.* 127. *Roy. lugdb.* 281. *Gouan. hort.* 248. *Mill. dict. n.* 2. *Kniph. cent.* 4. *n.* 50.
M. cryſtallinum plantaginis folio undulato. *Dill. elth.* 231. *t.* 180. *f.* 221.
Ficoides africana folio plantaginis undulato micis argenteis adſperſo. *Bradl. ſucc.* 5. *t.* 14. *f.* 48.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 110.
Eisartige Zaſerblume. W.
Habitat in Graecia *prope* Athenas. ☉ (v. v.)

Caulis *primum erectus, digitalis foliis oppoſitis.* Rami *horizontales longiſſimi oppoſiti proſtrati alterne ramoſi.* Foliis *alternis.*

Calyx *formatur ſolummodo e tribus laciniis, reliquae duae tam parvae, ut ab his tribus latis excludantur.* Lin. ſuppl.

Corolla *alba.* W.

*26. MESEMBRYANTHEMUM *humifuſum.*
M. foliis amplexicaulibus ſpathulatis carinatis, papulis conicis ſcabris, petalis mutiſſimis. *Ait. Kew.* 2. *p.* 179.
Geſtreckte Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

Corolla *alba.* W.

27. MESEMBRYANTHEMUM *papuloſum.*
M. foliis oppoſitis diſtinctis ovato - ſpathulatis, papulis ſubgloboſis, calycibus angulatis quinquefidis, ramis angulatis. *Ait. Kew.* 2. *p.* 182.
M. foliis oppoſitis ſpathulato - oblongis papuloſis, floribus pedunculatis, calycibus linearibus. *Lin. ſuppl.* 259.

Meſembryanthemum Aitonis. *Iacq. hort.* 3. *p.* 8. *t.* 7.
Blaſige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♂

Caules *et* Rami *depreſſi teretes papuloſi, ut omnes plantae partes una cum calyce, apice dichotomi.* Folia *oppoſita amplexicaulia fere connata ſpathulato-oblonga obtuſiuſcula.* Flores *ex dichotomiis ſolitarii erecti pedunculati, pedunculis filiformibus deflexis.* Calyx *laciniis inaequalibus lineari-ſubulatis.* Corolla *vix calyce longior lutea.* Capſulae *depreſſae nudae.*

Differt a M. cryſtallino, cui papulis cryſtallinis ſimillimum, caule *graciliore,* foliis *oppoſitis anguſtiorbus,* laciniis *calycinis ſubulatis nec ovatis latiſſimis,* corolla *multo minori lutea, quae in M. cryſtallino magna alba.*

Corolla *eſt rubicunda ſecundum Aitonium.* W.

*28. MESEMBRYANTHEMUM *ſeſſiliflorum.*
M. foliis planis ſpathulatis caulibusque papuloſis, ramis divaricatis, floribus ſeſſilibus. *Ait. Kew.* 2. *p.* 193.
Stiellosblütige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙
Corolla *lutea.* W.

*29. MESEMBRYANTHEMUM *glabrum.*
M. foliis amplexicaulibus diſtinctis ſpathulatis glaberrimis, pedunculis longitudine foliorum, calycibus hemiſphaericis, *Ait. Kew.* 2. *p.* 193.
Glattblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙
Corolla *lutea.* W.

*30. MESEMBRYANTHEMUM *Helianthoides.*
M. foliis ſpathulatis planis laevibus, pedunculis longiſſimis, calycibus baſi planis angulatis. *Ait. Kew.* 2. *p.* 193.
Sonnenblumenartige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙
Corolla *lutea.* W.

31. MESEMBRYANTHEMUM *pomeridianum.*
M. foliis planiuſculis lato-lanceolatis laevibus ſubciliatis diſtinctis, caule pedunculis germinibusque hirtis. *Sp. pl.* 698. *Lin. dec.* 23. *t.* 13.* *Iacq. ic. rar.* 3. *t.* 489.
Ficoides ſ. Ficus aizoides africana major, flore flavo, folio plano latiore. *Herm. lugdb.* 25. *Seb. muſ.* 1. *p.* 29. *f.* 5.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 122.

Ranunkelblütige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ☉

Nova e ſeminibus enata primo anno 1762, *adolevit et floruit.* Caulis *ſubdichotomus herbaceus palmaris teres, pilis patulis confertis albis.* Folia *ſpathulata ſ. lanceolata, petiolo baſi dilatato, diſtincta laevia ſubcarnoſa, margine imprimis petiolorum ſubciliato internodiis longiora patentia, prima terna, reliqua oppoſita.* Pedunculus *terminalis caule longior hirtus uti caulis.* Floris *calyx reliquis major, ſubquinquepartitus, baſi albo-villoſus, laciniis lanceolato-foliaceis corolla longioribus patulis, baſin verſus interius fuſco-membranaceis.* Corolla *ſulphurea multiplex, petalis linearibus acutis numeroſiſſimis, intimis ſetaceis umbilicum floris obtegentibus inſtar cancelli.* Stamina *lutea.* Styli 12 *totidemque fructus loculamenta.* Rami *e ſingulis alis, primo ſimiles, unde dichotoma planta, ſed debilis. Floret horis pomeridianis ab* 1-6 *niſi pluvia moneat corollam citius claudi, aut flos aetate provectior diutius expandatur.*

*32. MESEMBRYANTHEMUM *pallens.*

M. foliis oppoſitis amplexicaulibus diſtinctis oblongo-lanceolatis acutis obtuſe carinatis, papulis minutis. *Ait. Kew.* 2. *p.* 182. *Haworth Meſembr. p.* 197. *n.* 52.
Rinnenblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

Corolla *alba.* W.

33. MESEMBRYANTHEMUM *expanſum.*

M. foliis ovato-lanceolatis diſtinctis oppoſitis alternisve remotis, caule divaricato laxo procumbente. W.
M. foliis planiuſculis lanceolatis impunctatis patentibus diſtinctis, oppoſitis alternisque remotis. *Sp. pl.* 697. *Mill. dict. n.* 36. *Knorr. del.* 2. *t. M.* 3. *Haworth Meſembr. p.* 195. *n.* 50.
M. tortuoſum, foliis ſempervivi expanſis. *Dill elth.* 234. *t.* 182. *f.* 223.
Ficoides capenſis folio lato acuto, flore albo intus luteo. *Pet. gaz. t.* 78. *f.* 10.
Ficoides africana procumbens, foliis planis conjugatis lucidis. *Bradl. ſucc.* 3. *p.* 7. *t.* 16.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 104.
Ausgebreitete Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃

Flores *pentagyni.* Corolla *alba.* W.

84. MESEMBRYANTHEMUM *tortuoſum.*

M. foliis oblongo-ovatis connatis ſubpapilloſis confertis, calycibus triphyllis bicornibus, caule divaricato laxo procumbente. W.

M. foliis planiuſculis oblongo-ovatis ſubpapilloſis confertis connatis, calycibus triphyllis bicornibus. *Sp. pl.* 697. *Mill. dict. n.* 38. *Kniph. cent.* 8. *n.* 69. *Haworth Meſembr. p.* 196. *n* 51. *

M. foliis planis oppoſitis ovatis acuminatis connatis integerrimis. *Hort. cliff.* 217. *n.* 10. *

M. tortuoſum foliis ſempervivi congeſtis. *Dill. elth.* 233. *t.* 181. *f.* 222.

Ficoides capenſis procumbens oleae folio, flore albo medio candido. *Bradl. ſucc.* 2. *p.* 7. *t.* 16.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 121.

Gedrehte Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♃ (v. v.)

Calycis *foliola* 3 *expanſa*, 2 *cornuta.* Corolla *calyce brevior.* Piſtillum *tetragynum.*

Corolla *flaveſcens.* W.

* * * * * *Cauleſcentia, foliis ſubtus convexis.*

85. MESEMBRYANTHEMUM *apetalum.*

M. foliis amplexicaulibus diſtinctis linearibus ſupra planis, internodiis longioribus papuloſis, papulis oblongis, floribus pedunculatis, calycibus quinquefidis. *Ait. Kew.* 2. *p.* 180.

M. foliis lineari-ſublanceolatis ſubtus muricatis, caulibus proſtratis. *Lin. ſuppl.* 258.

Meſembryanthemum copticum. *Iacq. hort.* 3. *p.* 7. *t.* 6.

Kleinblütige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ⊙

Caules *herbacei proſtrati teretes rubentes, punctis obſoletis papuloſis nitentibus, ut tota planta, obtecti.* Folia *oppoſita linearia parumque lanceolata ſucculenta, ſupra canaliculata, ſubtus ſemicylindrica, papulis ſubmuricata obtuſa.* Pedunculi *breves papuloſi uniflori.* Calyx *foliolis inaequalibus ſubulato-cylindricis.* Petala 24 *alba diſtantia ſetacea breviſſima, quaſi totidem filamenta absque antheris, germini incumbentia.* Filamenta *ſubulata, petalis ſimillima, ſed breviora.* Antherae *fuſceſcentes.* Germen *inferum orbiculatum magnum convexum ſupra laeve.* Styli 5 *aproximati breves.* Stigmata *recurva.*

36. MESEMBRYANTHEMUM *copticum.*
M. foliis ſemiteretibus papuloſis diſtinctis, floribus ſeſſilibus axillaribus, calycibus quinquefidis. *Sp. pl.* 688.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 100.
Coptiſche Zaſerblume. W.
Habitat in Aegypto. ⊙
Statura et magnitudo M. nodiflori. Caulis *trichotomus herbaceus.* Folia *oppoſita ſemiteretia papuloſa, ramulos terminantia, nulla in internodiis, longitudine internodiorum.* Flores *ſeſſiles ſolitarii in divaricatura ramorum.* Calyx *quinquefidus, corolla longior.* Capſula *turbinata retuſa quinquangularis, angulis ciliatis compreſſis.* Semina *ſubrotunda ſubcochleata ſtriata.*
Corolla *alba.* W.

*37. MESEMBRYANTHEMUM *ciliatum.*
M. foliis oppoſitis connatis ſemiteretibus, ſtipulis membranaceis reflexis laceris ciliiformibus. *Ait. Kew.* 2. *p.* 179.
Gefranzte Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄.
Corolla *alba.* W.

*38. MESEMBRYANTHEMUM *caducum.*
M. foliis filiformibus ſemiteretibus diſtinctis, papulis ovatis, floribus lateralibus ſeſſilibus, terminalibus pari foliorum cinctis. *Ait. Kew.* 2. *p* 179.
Hinfällige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ⊙

39. MESEMBRYANTHEMUM *geniculiflorum.*
M. foliis ſemiteretibus papuloſis diſtinctis, floribus ſeſſilibus axillaribus, calycibus quadrifidis. *Sp. pl.* 688. *Mill. dict. n.* 3. *Haworth Meſembr. p.* 214. *n.* 59. *
M. ramis undique papilloſis folio craſſioribus. *Hort. cliff.* 218. *n.* 22. *Roy. lugdb.* 285.
M. capenſe geniculiflorum neapolitanum creditum *Dill. elth.* 271. *t.* 205. *f.* 261.
Ficoides neapolitana, flore candido *Boerh. lugdb.* 1. *p.* 291. *Bradl. ſucc.* 5. *t.* 17. *f.* 34.
Ficoides capenſe, folio tereti, flore albido. *Petiv. gaz. t.* 78. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 101.
Winkelblütige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Calyx *quadrifidus*. Styli *quatuor*. Fructus *quadrilocularis*.

Corolla *albido-flaveſcens*. W.

*40. MESEMBRYANTHEMUM *clavatum*.

M. foliis ſubteretibus impunctatis diſtinctis, pedunculis terminalibus aggregatis clavatis. W.

M. foliis ſubteretibus impunctatis diſtinctis, pedunculis clavatis, calycibus quadrifidis. *Iacq. hort. Schoenbr.* 1. *p.* 56. *t.* 108. *

Keulenſtielige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

Pedunculi *craſſi clavati in ramorum apice* 3-5 *aggregati*. Calyx *ſubaequalis quadrifidus*. Corolla *alba, petalis linearibus tenuiſſimis*. Styli *quatuor*. W.

41. MESEMBRYANTHEMUM *noctiflorum*.

M. foliis ſemicylindricis impunctatis diſtinctis, floribus pedunculatis terminalibus ſolitariis, calycibus clavatis. W.

M. foliis ſemicylindricis impunctatis diſtinctis, floribus pedunculatis, calycibus quadrifidis. *Sp. pl.* 689. *Mill. dict. n.* 4. *Knorr. del.* 1. *t. G* 8. *Haworth Meſembr. p.* 251. *n.* 72. *

M. foliis ſubulatis ſemicylindricis laevibus vix connatis apice recurvis. *Hort. cliff.* 220. *n.* 28. *Roy. lugdb.* 286.

M. noctiflorum flore intus albo extus phoeniceo odoratiſſimo. *Dill. elth.* 273. *t.* 206. *f.* 262.

Ficoides, ſeu Ficus aizoides africana erecta arboreſcens lignoſa, flore radiato, primo purpureo, dein argenteo, interdiu clauſo, nocte aperto. *Boerh. lugdb.* 1. *p.* 291.

β. M. (*ſtramineum*) foliis ſemiteretibus falcatis caule arboreſcente, floribus extus ſtramineis. *Haworth Meſembr. p.* 252. *n.* 73.

M. noctiflorum, flore intus candido extus ſtramineo odoratiſſimo. *Dill. elth.* 254. *t.* 206. *f.* 263.

Ficoides africana arboreſcens, folio tereti, flore candido nocte aperto, interdiu clauſo. *Boerh. lugd.* 1. *p.* 291. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 102.

Nächtliche Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Calyx *huic quadrifidus*. Flores *incarnato-albi, noctu fragrantes*. Styli *quatuor*. Fructus *quadrilocularis*.

42. MESEMBRYANTHEMUM *umbellatum*.

M. foliis ſubulatis ſcabrido-punctatis connatis, apice pa-

tulo, caule erecto, corymbo trichotomo. *Sp. pl.* 689. *Mill. dict. n.* 6. *Kniph. cent.* 3. *n.* 63. *Haworth Meſembr. p.* 336. *n.* 107.*

M. fruteſcens, floribus albis umbellatis. *Dill. elth.* 277. *t.* 208. *f.* 266.

Ficoides africana erecta teretifolia, floribus albis umbellatis. *Herm. par.* 166. *t.* 166. *Bradl. ſucc.* 4. *p.* 12. *t.* 34. *Pluk. phyt. t.* 117. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 104.

Doldentragende Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

43. MESEMBRYANTHEMUM *bicolorum.*

M. foliis ſubulatis punctatis laevibus diſtinctis, caule fruteſcente, corollis bicoloribus. *Sp. pl.* 695. *Mill. dict. n.* 28. *et. ic.* 177. *f.* 1. *Curt. mag.* 59.

M. (*coccineum*) foliis ſemicylindrico-compreſſis carinatis obtuſis, pedunculis ſubbracteatis baſi laevigatis, petalis coccineis. *Haworth Meſembr. p.* 247. *n.* 71.*

Ficoides ſ. Ficus aizoides africana minor erecta, triangulari folio viridi, flore intus aureo, foris purpureo. *Herm. lugdb.* 250. *t.* 249. *Moriſ. hiſt.* 3. *p.* 507. *ſ.* 12. *t.* 6. *f.* 4.

β. M. (*bicolorum*) foliis ſubtriquetris ſubulatis ſcabrido-punctatis, pedunculis nudis papuloſo-ſcabridis, corollis luteis. *Haworth Meſembr. p.* 241. *n.* 70.*

M. tenuifolium fruticeſcens flore croceo. *Dill. elth.* 267. *t.* 202. *f.* 258.

Ficoides capenſis fruteſcens, folio tereti punctato, petalis luteis. *Bradl. ſucc.* 1. *p.* 8. *t.* 7.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 118.

Zweyfarbige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Corollae *extus purpureae intus flavae, apice extus purpureae* .Calyx *tuberculatus.*

44. MESEMBRYANTHEMUM *tuberoſum.*

M. foliis ſubulatis papuloſis diſtinctis apice patulis, radice capitata. *Sp. pl.* 693. *Hort. cliff.* 216. *n.* 4.* *Roy. lugdb.* 282. *Mill. dict. n.* 21. *Haworth Meſembr. p.* 271. *n.* 86.*

M. fruticeſcens, radice ingenti tuberoſa. *Dill. elth.* 275. *t.* 207. *f.* 264.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 112.

Knollige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. ſ.)
Flores *parvi pallide purpurei.* W.

45. MESEMBRYANTHEMUM *tenuifolium.*
M. foliis ſubfiliformibus glabris diſtinctis internodio longioribus, caulibus procumbentibus. *Syſt. veg.* 390.
M. foliis ſemiteretibus ſubulatis glabris diſtinctis internodio longioribus. *Hort. cliff.* 220. *n.* 26. *Hort. upſ.* 128. *Roy. lugdb.* 286. *Sp.* 2. *p.* 693. *Mill. dict. n.* 22. *Haworth Meſembr. p.* 239. *n.* 68.
M. tenuifolium procumbens, folio tenuiore viridi, flore coccineo. *Dill. elth.* 264. *t.* 201. *f.* 256.
Ficoides capenſis humilis teretifolia, flore coccineo. *Bradl. ſucc.* 1. *p.* 13. *t.* 9.
Ficoides africana minor procumbens, folio tenuiore viridi, flore coccineo. *Moriſ. hiſt.* 3. *p.* 507. *ſ.* 12. *t.* 8. *f.* 6.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 112.
Dünnblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Caules *lignoſi. procumbentes.* Rami *ex ſingulis alis ſuperioribus.* Flores *coccinei, ſpecioſi multis petalis, medio die, ſole illuſtrante expanſi, diutius durantes.* Folia *glaberrima.*

46. MESEMBRYANTHEMUM *ſtipulaceum.*
M. foliis ſubtriquetris compreſſis incurvis punctatis diſtinctis congeſtis: baſi marginatis. *Sp. pl.* 693. *Hort. cliff.* 220. *n.* 29. *Roy. lugdb.* 287. *Mill. dict.* 23. *Haworth Meſembr. p.* 255. *n.* 77. *
M. fruteſcens, flore purpureo rariore. *Dill. elth.* 279. *t.* 209. *f.* 267, 268.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 113.
Steifſtielige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)
Caulis *florens rigidus, ſubgriſeus.* Folia *linearia, carnoſa, lateribus valde compreſſa et punctis pellucidis adſperſa, incurva, ſupra planiuſcula, mucronata.* Flores *ſubcorymboſi, purpurei,* Staminibus *luteis.*

47. MESEMBRYANTHEMUM *corniculatum.*
M. foliis triquetro-ſemicylindricis ſcabrido-punctatis, ſupra baſin linea elevata connatis. *Sp. pl.* 676. *Mill. dict. n.* 35.
M. (*corniculatum*) foliis imbricatis longis aequalibus,

mollibusque punctatis, baſi ſemicylindricis, apice ſemicylindrico-triquetris. *Haworth Meſembr. p* 226. *n.* 26.*
M. foliis corniculatis longioribus. *Dill. elth.* 262. *t.* 199. *f.* 253. 254.
Ficoides capenſis folio triangulari, flore luteo intus pallido. *Petiv. gaz. t.* 77. *f.* 10.
β. M. (*diverſiphyllum*) foliis ramorum connatis inaequalibus, ſcabrido-punctatis, ſemicylindrico-triquetris, caule proſtrato elongato, petalis ſubtus rubris *Haworth Meſembr. p.* 228. *n.* 63.*
M. foliis corniculatis brevioribus. *Dill. elth.* 261. *t.* 198. *f.* 252.
Ficoides africana reptans, folio triangulari craſſo longiſſimo, flore intus luteo extus aurantio. *Bradl. ſucc.* 4. *p.* 18. *t.* 40.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 121.
Hornförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃ (v. v.)
Flores *decagyni.*

48. MESEMBRYANTHEMUM *loreum.*
M. foliis ſemicylindricis recurvis congeſtis baſi interiore gibbis connatis, caule pendulo. *Sp. pl.* 694. *Mill. dict. n.* 30.
M. loreum. *Dill. elth.* 264. *t.* 200. *f.* 255.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 115.
Riemenartige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♃
Flores rariſſime prodiere apud Engelb. Gother, unde patuit diſtinctam a. M. corniculato plantam eſſe. Flos *ex alis foliorum infimis rami, pedunculo brevi;* Calyce *aequali mediocri;* Corolla *purpurea, petalis lanceolato-linearibus.* Staminibus *albidis.*

49. MESEMBRYANTHEMUM *veruculatum.*
M. foliis triquetro-cylindricis acutis connatis arcuatis impunctatis diſtinctis. *Sp. pl.* 696. *Hort. cliff.* 220. *n.* 24. *Hort. upſ.* 128. *Roy. lugdb.* 285. *Mill. dict. n.* 33. *Kniph. cent.* 10. *n.* 62. *Haworth Meſembr. p.* 260. *n.* 79.
M. foliis veruculiformibus, floribus mellinis umbellatis. *Dill. elth.* 268. *t.* 203. *f.* 259.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 123.
Bogigblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)

Folia *ſubtriquetro-cylindrica pellucida, apice purpuraſcentia.* Flores *adoratiſſimi conferti parvi flavi.* Corolla *vix calyce longior, petalis apice ſubbidentatis.* Stamina *parva*; Filamentis *baſi villoſis.*

*50. MESEMBRYANTHEMUM *echinatum.*

M. foliis oblongo-ovatis ſubtriquetris gibbis ramentaceo-hiſpidis, laciniis calycinis foliiformibus. *Ait. Kew.* 2. *p.* 194. *Haworth Meſembr. p.* 288. *n.* 91.*

α. flore luteo.

β. flore albo.

Stachliche Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

Variat flore luteo et albo. W.

*51. MESEMBRYANTHEMUM *viridiflorum.*

M. foliis ſemicylindraceis papuloſo-piloſis, calycibus quinquefidis hirſutis. *Ait. Kew.* 2. *p.* 196. *Haworth Meſembr. p.* 199. *n.* 53.*

Grünblütige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

Corolla *laete viridis* W.

52. MESEMBRYANTHEMUM *ſplendens.*

M. foliis ſubteretibus impunctatis recurvis diſtinctis congeſtis, calycibus digitiformibus terminalibus. *Sp. pl.* 689. *Mill. dict. n.* 5. *Haworth Meſembr. p.* 209. *n.* 56.*

M. foliis confertis ſplendentibus, flore pallido. *Dill. elth.* 270. *t.* 204. *f.* 260.

Ficoides capenſis fruteſcens, foliis teretibus confertis glaucis, flore albo. *Bradl. ſucc.* 1. *p.* 7. *t.* 6.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 103.

Glänzende Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Caulis *punctis albidis in ramis non ultimis.* Folia *triquetro-teretiuſcula, apice patula acuta, axillaria vero incurva.* Corolla *flaveſcenti-albida.* Calyces *erecti digitiformes mutici.*

53. MESEMBRYANTHEMUM *villoſum.*

M. foliis pubeſcentibus connatis impunctatis, caule piloſo. *Sp pl* 692. *Hort. cliff.* 217. *n.* 6.* *Roy. lugdb.* 282. *Mill. dict. n.* 16. *Haworth Meſembr. p.* 222. *n.* 61.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 109.

Behaarte Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

Folia *lineari-ſemiteretia canaliculato-plana opaca, margine, verſus baſin imprimis, villis adſperſa.* Caulis *pubeſcens villis vix exſtantibus.* Flores *non vidi.*
Flores *rubicundi.* W.

54. MESEMBRYANTHEMUM *micans.*
M. foliis ſubcylindricis papuloſis diſtinctis caule ſcabro. *Sp. pl.* 696. *Hort. cliff.* 220. *n.* 30.* *Roy. lugdb.* 287 *Mill. dict. n.* 37. *Kniph. cent.* 7. *n.* 58. *Knorr. del.* 1. *t. G.* 5. *Haworth Meſembr. p.* 268. *n.* 84.
M. micans, flore phoeniceo, filamentis atris. *Dill. elth.* 292. *t.* 215. *f.* 282.
Ficoides capenſis tereti folio flore croceo. *Petiv. gaz. t.* 3. *f.* 9.
Ficoides capenſis folio tereti argenteo, petalis perplurimis aurantiacis. *Bradl. ſucc.* 1. *p.* 9. *t.* 8.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 119.
Schimmernde Zaſerblume. W.
Habitat ad Cap. b. ſpei ♄

*55. MESEMBRYANTHEMUM *groſſum.*
M. foliis ſubcylindricis confertis papuloſis, caudice baſi incraſſato, ramis diffuſis glabris. *Ait. Kew.* 2. *p.* 191.
Dickſtielige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Corolla *lutea.* W.

* * * * * * *Cauleſcentia, foliis teretibus.*

56. MESEMBRYANTHEMUM *nodiflorum.*
M. foliis alternis teretiuſculis obtuſis baſi ciliatis. *Sp. pl.* 687. *Hort. upſ.* 129. *Mill. dict. n.* 1. *Forſk. aegypt. p.* 98.
Kali craſſulae minoris foliis. *Bauh. pin.* 289. *Moriſ. hiſt.* 2. *p.* 610. *ſ.* 5. *t.* 33. *f.* 7.
Kali neapolitanum aizoides repens. *Col. ecphr.* 2. *p.* 72. *t.* 73.
Kali II. *Alp. aegypt.* 125. *t.* 127.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 99.
Knotenblütige Zaſerblume. W.
Habitat in Aegypto, Neapoli *locis arenoſis editis, quae ab aeſtu maris adſperguntur.* ⊙
Corolla *alba.* W.

57. MESEMBRYANTHEMUM *capillare.*
M. foliis connatis teretibus papuloſis, caule erecto, ramulis unifloris filiformibus glabris. *Lin. ſuppl.* 260.
Haarblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. *Thunberg.*

*58. MESEMBRYANTHEMUM *laeve.*
M. foliis cylindraceis obtuſis, amplexicaulibus laevibus, calycibus quinquefidis, laciniis oblongis obtuſis. *Ait. Kew. p* 187. *Haworth Meſembr. p.* 254. *n.* 75.
Weiſſ ſtielige Zaſerblume. W.
Habitat ad Cap. b. ſpei ♄
Corolla *purpurea.* W.

*59. MESEMBRYANTHEMUM *brachiatum.*
M. caulibus foliisque cylindricis papuloſis, ramis trichotomis. *Ait. Kew.* 2. *p.* 191. *Haworth Meſembr. p.* 254. *n.* 74.
Wendäſtige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Corolla *lutea.* W.

*60. MESEMBRYANTHEMUM *brevifolium.*
M. foliis cylindraceis obtuſiſſimis papuloſis patentibus, ramis diffuſis. *Ait. Kew.* 2. *p.* 188. *Haworth Meſembr. p.* 274. *n.* 87.*
Kurzblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Corolla *purpurea.* W.

61. MESEMBRYANTHEMUM *hiſpidum.*
M. foliis cylindricis papuloſis diſtinctis, caule pedunculisque hiſpidis, calyce glabro papuloſo. W.
M. foliis cylindricis papuloſis diſtinctis, caule hiſpido. *Sp. pl.* 691. *Mill. dict. n.* 15. *et ic. t.* 176. *f.* 3. *Kniph. cent.* 3. *n.* 62.
M. caule hiſpido. *Hort. cliff.* 217. *n.* 7.* *Roy. lugdb.* 282.
M. foliis diſtantibus cylindricis, caule hirto, calycibus glabris papuloſis, ſtaminibus purpuraſcentibus. *Haworth Meſembr. p* 277. *n.* 88.*
M. piloſum micans, flore ſaturate purpureo. *Dill. elth.* 289. *t.* 214. *f.* 278.
Ficoides afra fruticoſa, caule lanugine argentea ornato, folio tereti parvo longo, guttulis argenteis quaſi ſcabro, flore violaceo. *Boerh. lugdb.* 1. *p.* 291.
β. Meſembryanthemum hiſpidum pallidum. *Haworth Meſembr. p.* 279.*
M. piloſum micans, flore purpureo pallidiore. *Dill. elth.* 290. *t.* 214. *f.* 279. 280.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 108.
Borſtige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄

Pedunculus *valde deorſum ſcaber, magis viſu, quam tactu.* Calyx *muticus.* Flos *odoratus ſimillimus M. barbati (a quo forte olim ortum) demta barba foliorum, ſed cauli inſperſa.*

*62. MESEMBRYANTHEMUM *ſtriatum.*

M. foliis cylindricis papuloſis diſtinctis, caule pedunculisque hiſpidis, calyce lanuginoſo. W.

M. (*ſtriatum*) foliis cylindraceis, caule hiſpido, pedunculis calycibusque lanuginoſis. *Haworth Meſembr. p.* 280. *n.* 89. *

Meſembryanthemum hiſpidum. *γ. Sp. pl.* 691.

M. piloſum micans, flore purpureo ſtriato. *Dill. elth.* 291. t. 215. *f.* 281.

Ficoides ſ. Ficus azoides africana folio longo tenui, flore aurantio. *Brad. ſucc.* 4. *p.* 13. *t.* 35.

Ficus aizoides teretifolia, foliis cryſtallino rore eleganter conſperſis, floribus dilute roſeis. *Volk. heſp.* 221. *t.* 224. *f.* 2.

Geſtreifte Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄.

63. MESEMBRYANTHEMUM *barbatum.*

M. foliis ſubovatis papuloſis diſtinctis apice barbatis. *Hort. cliff.* 216. *n.* 5. * *Hort. upſ.* 127. *Roy. lugdb.* 282. *Mill. dict. n.* 13. *ic. t.* 176. *f.* 3. *Kniph. cent.* 10. *n.* 60.

M. foliis ſubovatis papuloſis diſtinctis apice barbatis ramulis prolixis recumbentibus, calycibus pentaphyllis glabris. *Haworth Meſembr. p.* 293. *n.* 92. *

M. radiatum, ramulis prolixis recumbentibus. *Dill. elth.* 245. *t.* 190. *f.* 234.

Ficoides capenſis teretifolia apicibus hirtis. *Petiv. gaz. t.* 77. f. 6.

Ficoides capenſis fruteſcens, folio tumido, extremitate ſtellata, flore purpureo. *Bradl. ſucc.* 1. *p.* 6. *t.* 5. *et* 15.

Ficus aizoides, folio tereti in villos radiatos abeunte flore rubro. *Volk. heſp. p.* 222. *t.* 124. *f.* 6.

β. M. (*hirſutum*) ramulis hirſutis, foliis ſemicylindrico-turgidis apice barbatis, baſi ciliatis ſcabrido papuloſis, calycibus ſex-vel ſeptemphyllis inaequalibus. *Haworth Meſembr. p.* 297. *n.* 93. *.

M. radiatum humile, foliis minoribus. *Dill. elth.* 246. *t.* 235. *Willich. obſ. n.* 56.

γ. M. (*denſum*) ſubacaule, foliis denſiſſime imbricatis connatis ſemicylindricis, ſupra concavis apice barbatis, calycibus hexaphyllis ſubaequalibus hiſpidis. *Haworth Meſembr. p.* 302. *n.* 94. *

M. radiatum humile, foliis majorbus. *Dill. elth.* 248. *t.* 190. *f.* 236.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 107.

Bärtige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Calycis *foliola apice barbata, aeque ac folia plantae demonſtrant* Perianthii *ortum.* Petala *minima interiora, quae cingunt ſtamina alba. Varietas* β. *hybrida forte differt ab* α. Calyce *hexaphyllo nec pentaphyllo.* Stylis Loculiis*que* 6, *nec* 5 *ſ.* 8. Pedunculis *hiſpidis piloſis nec laevibus.* Foliorum *oppoſitionibus imbricatis, nec remotis.*

* * * * * * * *Cauleſcentia, foliis triquetris.*

64. MESEMBRYANTHEMUM *falcatum.*

M. foliis ſubacinaciformibus incurvis punctatis diſtinctis, ramis teretibus. *Sp. pl.* 694. *Hort. cliff.* 219. *n.* 19. * *Roy lugdb.* 285. *Knorr. del.* 2. *t. M.* 4. *Haworth. Meſembr. p.* 360. *n.* 118. *

M. falcatum minimum, flore purpureo parvo. *Dill. elth.* 288. *t.* 213. *f.* 276.

Ficoides afra, folio triangulari enſiformi breviſſimo, flore dilute purpuraſcente filamentoſo. *Boerh. lugdb.* 1. *p.* 290. *Bradl. ſucc.* 5. *p. t.* 42.

Sichelblättrige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

65. MESEMBRYANTHEMUM *glomeratum.*

M. foliis teretiuſculis compreſſis punctatis diſtinctis, caule paniculato multifloro. *Sp. pl.* 694. *Mill. dict. n.* 26. *Haworth Meſembr.* 356 *n.* 117. *

M. falcatum minus, flore carneo minore. *Dill. elth.* 281. *t.* 213. *f.* 274.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 114.

Knaulblättrige Zaſerblume. W.

Habitat ad Cap. b. ſpei. ♄ (v. v.)

Caulis *diffuſus paniculatus.* Folia *angulis obtuſis trigona glauca, punctis diaphanis.* Ramulorum *rudi-*

menta ex alis, foliis confertiſſimis incurvis. Calyx *longius pedunculatus;* Germine *ſubverrucoſo, punctis eminentibus obſoletis.* Laciniis *calycinis apice membranaceis ſub apice ſpina patula.* Corolla *purpurea M. deltoidis.*

*66. MESEMBRYANTHEMUM *reptans.*

M. foliis triquetris acutis ſcabris, caule reptante. *Ait. Kew.* 2. *p.* 185. *Haworth Meſembr. p.* 348. *n.* 113.*
Kriechende Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Corolla *purpurea.* W.

*67. MESEMBRYANTHEMUM *deflexum.*

M. foliis triquetris acutis glaucis, punctis obſoletis ſcabriuſculis, laciniis calycinis interioribus membranaceis. *Ait. Kew.* 2. *p.* 187. *Haworth Meſembr. p.* 347. *n.* 112.*
Niedergebengte Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Petala *pallide-purpurea valde anguſta.* W.

*68. MESEMBRYANTHEMUM *auſtrale.*

M. foliis ſubtriquetris punctulatis connatis obtuſiuſculis, caule tereti repente, pedunculis obtuſe ancipitibus. *Ait. Kew.* 2. *p.* 187. *Haworth Meſembr.* 349. *n.* 114.
Südliche Zaſerblume. W.
Habitat in Nova Zeelandia. ♃
Corolla *purpurea.* W.

69. MESEMBRYANTHEMUM *ſpinoſum.*

M. foliis tereti-triquetris punctatis diſtinctis, ſpinis ramoſis. *Sp. pl.* 693. *Hort. cliff* 216. *n.* 3.* *Roy. lugdb.* 281. *Haworth Meſembr. p.* 338. *n.* 108.*
M. fruticeſcens, ramis triacanthis. *Dill. elth.* 276. *t.* 208. *f.* 265.
Ficoides africana, aculeis longiſſimis et foliolis naſcentibus e foliorum alis. *Tournef. act.* 1705. *p.* 316. *Bradl. ſucc.* 41. *t.* 39.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 111.
Dornige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄. (v. v.).
Corolla *purpurea.* W.

70. MESEMBRYANTHEMUM *craſſifolium.*

M. foliis ſemicylindricis impunctatis connatis, apice tri-

quetris, caule repente ſemicylindrico. *Sp. pl.* 693. *Hort. cliff.* 217. *n.* 9.* *Roy. lugdb.* 283. *Mill. dict. n.* 24. *Knorr. del.* 2. *t. M.* 4. *Haworth Meſembr. p.* 350. *n.* 115.*
M. craſſifolium repens flore purpureo. *Dill. elth.* 266. *t.* 201. *f.* 257.
Ficoides africana reptans, folio triangulari viridi, flore ſaturate purpureo. *Bradl. ſucc.* 4. *p.* 16. *t.* 38.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 114.
Dickblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

*71. MESEMBRYANTHEMUM *molle.*
M. foliis triquetris connatis erectis glaucis impunctatis, ramis ſemiteretibus, pedunculis axillaribus compreſſis. *Ait. Kew* 2 *p.* 192. *Haworth Meſembr. p.* 261. *n.* 80.*
Weiche Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Flores *lutei.* W.

72. MESEMBRYANTHEMUM *glaucum.*
M. foliis triquetris acutis punctatis diſtinctis, calycinis foliolis ovato-cordatis. *Sp. pl.* 696. *Hort. cliff.* 220. *n.* 27. *Roy lugdb.* 183. *Mill. dict. n.* 34. *Haworth Meſembr. p.* 329. *n.* 104.*
M. ſcabrum, flore ſulphureo convexo. *Dill. elth.* 256. *t.* 196. *f.* 248.
Ficoides africana fruteſcens, folio tringulari breviori glauco. *Moriſ. hiſt.* 3. *p.* 507. *ſ.* 12. *t.* 6. *f.* 3.
Ficoides afra, caule lignoſo, erecto, folio triangulari enſiformi ſcabro, flore luteo magno. *Bradl. ſucc.* 4. *p.* 15. *t.* 37.
Ficoides ſ. Ficus aizoides africana minor erecta, folio triangulari glauco, flore luteo. *Herm. lugdb.* 247. *t.* 248.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 120.
Meergrüne Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. ſ.)
Folia *mucronata., rore glauca, punctis ſparſis non ſcabra, niſi carina.*

*73. MESEMBRYANTHEMUM *ſpectabile.*
M. foliis triquetris acutis connatis arcuatis punctatis glaucis laevibus, caule adſcendente. W.
M. (*ſpectabile*) foliis perfoliatis longiſſimis glaucis punctatis integerrimis triquetris apice ſubulatis, caule lignoſo

adſcendente. *Haworth Meſembr. p.* 385. *n.* 126. *
Curt. mag. 396.
Anſehnliche Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄
Flores *magni coccineo-purpurei.* Pedunculi *longi uniflori ſolitarii terminales, medio bracteis binis connatis inſtructi.* W.

74. MESEMBRYANTHEMUM *emarginatum.*
M. foliis ſubulatis congeſtis ſubſcabris, calycibus ſpinoſis, petalis emarginatis. *Sp. pl.* 692. *Haworth Meſembr. p.* 340. *n.* 109. *
M. purpureum ſcabrum, ſtaminibus expanſis. *Dill. elth.* 259. *t.* 197. *f.* 250.
Ficoides capenſis triangulari folio acuto, flore purpureo. *Petiv. gaz. t.* 77. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 100.
Ausgerandete Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

75. MESEMBRYANTHEMUM *aureum.*
M. foliis cylindrico-triquetris punctatis diſtinctis piſtillis atropurpureis. *Syſt. nat. ed.* 10. *p.* 1060. *Curt. mag.* 262. *Haworth Meſembr. p.* 333. *n.* 106. *
Goldfarbene Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)
Corolla *crocea.* W.

76. MESEMBRYANTHEMUM *ſerratum.*
M. foliis ſubulatis triquetris punctatis diſtinctis angulo carinali retrorſum ſerratis. *Sp. pl.* 696. *Hort. cliff.* 218. *n.* 15. *Roy. lugdb.* 284. *Mill. dict. n.* 31.
M. foliis ſubulatis triquetris punctatis diſtinctis, angulo carinali tetrorſum ſerrato, flore acetabuliformi luteo. *Haworth Meſembr. p.* 381. *n.* 124. *
M. ſerratum flore acetabuliformi luteo. *Dill. elth.* 249. *t.* 192. *f.* 238.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 119.
Geſägte Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

77. MESEMBRYANTHEMUM *ſcabrum.*
M. foliis ſubulatis diſtinctis, ſubtus undique punctato-muricatis, calycibus muticis. *Sp. pl.* 692. *Hort. cliff.* 219. *n.* 20. * *Hort. upſ.* 129. *Mill. dict. n.* 17. *Haworth Meſembr. p.* 345. *n.* 111. *

M. purpureum ſcabrum, ſtaminibus collectis. *Dill. elth.* 260. *t.* 197. *f.* 251.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 109.
Scharfblättrige Zaſerblume. W.
Habiat ad Cap. b. ſpei. ♄ v. ſ.)

*78. MESEMBRYANTHEMUM *bracteatum.*
M. foliis ſubacinaciformibus punctatis, apice recurvo, bracteis amplexicaulibus lato-ovatis carinatis. *Ait. Kew.* 2. *p.* 185. *Haworth Meſembr. p.* 319. *n.* 101.*
M. uncinatum ſcabrum, petalis purpureis circumactis. *Dill. elth.* 258. *t.* 197. *f.* 249.
Beblätterte Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

79. MESEMBRYANTHEMUM *uncinatum.*
M. articulis caulinis terminatis foliis connatis acuminatis ſubtus dentatis. *Sp. pl.* 692. *Hort. cliff.* 218. *n.* 16. *Roy. lugdb.* 284. *Mill. dict. n.* 18.
M. articulis caulinis terminatis foliis connatis acuminatis brevibus punctatis triquetris, ſubtus diacanthis. *Haworth Meſembr. p.* 309. *n.* 96. *
M. perfoliatum, foliis minoribus diacanthis. *Dill. elth.* 250. *t.* 193. *f.* 239.
Ficoides afra, folio triangulari glauco perfoliato breviſſimo apice ſpinoſo. *Bradl. ſucc.* 3. *p.* 8. *t.* 27. 26. *Burm. afr. t.* 26. *f.* 3.
β. M. (*perfoliatum*) articulis caulinis terminatis foliis connatis acuminatis punctatis glaucis triquetris, ſubtus triacanthis. *Haworth Meſembr. p* 307. *n.* 95. *
M. perfoliatum, foliis majoribus triacanthis. *Dill. elth.* 251. *t.* 193. *f.* 240.
Ficoides africana fruteſcens perfoliata, folio triangulari glauco punctato. *Bradl. ſucc.* 5. *p.* 13. *t.* 46.
Houttuyn Lin. Pfl. Syſt. 7 *p.* 110.
Hackenförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)
Corolla *purpurea.* W.

80. MESEMBRYANTHEMUM *pugioniforme.*
M. foliis alternis confertis ſubulatis triquetris longiſſimis impunctatis. *Sp. pl.* 699. *Hort. cliff.* 216. *n.* 2. * *Hort. upſ.* 129. *Roy. lugdb.* 281. *Mill. dict. n.* 46. *Haworth Meſembr. p.* 387. *n.* 227. *
M. folio pugioniformi, flore aureo ſtramineo. *Dill. elth.* 280. *t.* 210. *f.* 269.

Ficus capenſis caryophylli folio, flore aureo ſpecioſo. *Bradl. ſucc.* 2. *p.* 5. *t.* 14.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 127.
Dolchförmige Zaſerblume. W.
Habitat ad Cap. ſpei. ♄

Caules *apice ramoſi, ramis ternis.* Folia *ex apice ramorum conferta, ad exortum ramulorum florentium apice uncinata.* Pedunculi *longi.* Flores *decagyni, in hoc genere maximi.*

*81. MESEMBRYANTHEMUM *aequilaterale.* †

M. foliis ſubconnatis acutis glabris arcuatis confertis aequilateri-triquetris caule, angulato repente. W.
M. (*aequilaterum*) foliis confertis ſubconnatis longis obſolete punctatis ſubaequilateri-triquetris integerrimis, caule angulato ſubradicante proſtrato. *Haworth Meſembr. p.* 390. *n.* 128. *
Gleichſeitige Zaſerblume. W.
Habitat in Nova Hollandia. ♄.

Flores *non vidit Clariſſ. Haworth.* W.

82. MESEMBRYANTHEMUM *filamentoſum.*

M. foliis aequilateri-triquetris acutis ſubpunctatis connatis, angulis ſcabris, ramis hexagonis. *Sp. pl.* 694. *Haworth Meſembr. p.* 380. *n.* 123. *
M. falcatum majus, flore purpureo mediocri. *Dill. elth.* 285. *t.* 212. *f.* 273.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 115.
Fädige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄

Folia *punctata vix connata.* Flores *ſeſſiles pentagyni.* Corollae *miſerae parvae, obſolete violaceae, petalis ſubulatis nec lanceolatis.*

83. MESEMBRYANTHEMUM *acinaciforme.*

M. foliis acinaciformibus impunctatis connatis, angulo carinali ſcabris, petalis lanceolatis. *Sp. pl.* 695. *Hort. cliff.* 219. *n.* 18. *Roy. lugdb.* 284. *Mill. dict. n.* 19. *Kniph. cent.* 10. *n.* 59. *Haworth Meſembr. p.* 397. *n.* 131. *
M. acinaciforme flore ampliſſimo purpureo. *Dill. elth.* 282. *t.* 211. *f.* 270. *et. t.* 212. *f.* 271.
Sebelförmige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)

84. MESEMBRYANTHEMUM *forficatum.*

M. foliis acinaciformibus obtuſis impunctatis connatis apice

ſpinoſis, caule ancipiti. *Sp. pl.* 695. *Iacq. hort.* 1. *p.* 9. *t.* 26.* *Haworth Meſembr. p.* 394. 130.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 117.
Scheerenblättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)
Caulis *debilis.* Corolla *purpurea.*

85. MESEMBRYANTHEMUM *edule.*
M. foliis aequilatetri-triquetris acutis ſtrictis impunctatis connatis, carina ſubſerratis, caule ancipiti. *Sp. pl* 695. *Mill. dict. n.* 27. *Haworth Meſembr. p.* 392. *n.* 126.
M. falcatum majus, flore amplo luteo. *Dill elth.* 283. *t.* 212. *f.* 272.
Ficoides ſ. Ficus aizoides africana major procumbens triangulari folio, fructu maximo eduli. *Herm. lugdb.* 244. *t.* 245. *Moriſ. hiſt.* 3. *p.* 506. *ſ.* 12. *t.* 7. *f.* 1.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 117.
Genieſsbare Zaſerblume. W.
Habitat in arenoſis maritimis. Cap. b. ſpei. ♄
Folia *craſſitie digiti nitida recta triquetra, aequilatera longa baſi connata, carina cartilaginea ſcabra.*

86. MESEMBRYANTHEMUM *deltoides.*
M. foliis deltoidibus triquetris dentatis impunctatis diſtinctis. *Sp. pl.* 690. *Hort. cliff.* 218. *n.* 14.* *Roy. lugdb.* 284. *Mill. dict. n.* 11. *Knorr. del.* 1. *t. G. b. n.* 1.
M. foliis deltoidibus triquetris, dorſo et lateribus muricatis, impunctatis diſtinctis. *Haworth Meſembr. p.* 364. *n.* 120.*
M. deltoides et dorſo et lateribus muricatis minus. *Dill. elth.* 255. *t.* 195. *f.* 246.
Ficus aizoides africana erecta, folio triangulari breviuſculo fimbriato, floribus roſeis odoratis. *Volk. heſp.* 223. *t.* 224. *f.* 5.
β. M. deltoides et dorſo et lateribus muricatis majus. *Dill. elth.* 244. *t.* 196. *f.* 247. *Knorr. del.* 1. *t. G.* 5. *b. n.* 2.
γ. M. (*cauleſcens*) foliis albidis deltoidibus, lateribus minime dentatis, angulo carinali integris. *Haworth Meſembr. p.* 367. *n.* 121.*
M. deltoides non dorſo, ſed lateribus muricatis. *Dill. elth.* 253. *t.* 195. *f.* 243. 244.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 107.
Deltablättrige Zaſerblume. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)

Pedunculi *uniflori diphylli.* Corollae *dilute purpureae odoratae, non calyce longiores.* Stamina *erecta in conum, alba;* Antheris *flavis.*

995. AIZOON.

Gen. plant. edit. Schreb. n. 861.

Cal. 5-partitus. *Petala* o. *Capſ.* ſupera, 5-locularis, 5-valvis.

1. AIZOON *canarienſe.*

A. foliis cuneiformi-ovatis, floribus ſeſſilibus. *Hort. upſ.* 127. *Mill. dict. n.* 1. *Kniph. cent.* 1. *n.* 1.

A. foliis obverſe ovatis. *Hort. cliff.* 215. *Roy. lugdb.* 221.

Veſlingia Heiſteri. *Fabric. helmſt. p.* 363.

Glinus cryſtallinus. *Forskaehl. fl. aegypt. p.* 95. *n.* 97. *et ic. t.* 14.

Ficoides procumbens, portulacae folio. *Niſſ. act.* 1711. *p.* 422. *t.* 13. *f.* 1.

Kali aizoides canarienſis procumbens. *Pluk. alm.* 202. *t.* 303. *f.* 4. *Volk. norib.* 236. *t.* 236.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 130.

Canariſches Immergrün. W.

Habitat in Canariis. ⊙ (v. v.)

2. AIZOON *perfoliatum.*

A. tomentoſum, foliis obovatis connatis cryſtallino-punctatis, floribus pedunculatis. *Lin. ſuppl.* 261.

Durchwachſenes Immergrün. W.

Habitat ad Cap. b. ſpei.

3. AIZOON *Glinoides.*

A. foliis ſubrotundo-cuneiformibus piloſis, floribus ſeſſilibus, calycibus hirſutis. *Ait. Kew.* 2. *p.* 196.

A. hirſutum herbaceum procumbens, foliis ovatis, floribus ſeſſilibus diſtinctis. *Lin. ſuppl.* 261.

Glinusartiges Immergrün. W.

Habitat ad Cap. b. ſpei. ♄ (v. ſ.)

4. AIZOON *ſecundum.*

A. hirſuto-canum herbaceum procumbens, foliis ovatis, floribus ſeſſilibus imbricatis ſecundis. *Lin. ſuppl.* 261.

Einſeitiges Immergrün. W.

Habitat ad Cap. b. ſpei.

5. AIZOON *rigidum.*

A. fruticoſum procumbens tomentoſum, foliis ovatis, floribus ſeſſilibus remotis. *Lin. ſuppl.* 261.
Steifes Immergrün. W.
Habitat ad Cap. b. ſpei. ♄

6. AIZOON *hiſpanicum.*

A. foliis lanceolatis, floribus ſeſſilibus. *Sp. pl.* 700. *Mill. dict. n.* 2.
Ficoidea hiſpanica annua, folio longiore. *Dill. elth.* 143. *t.* 117. *f.* 143.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 131.
Spaniſches Immergrün. W.
Habitat in Hiſpania. ☉ (v. v.)

7. AIZOON *fruticoſum.*

A. fruticoſum erectum glabrum, foliis lanceolatis, floribus ſeſſilibus. *Lin. ſuppl.* 261.
Strauchartiges Immergrün. W.
Habitat ad Cap. b. ſpei. ♄

8. AIZOON *paniculatum.*

A. hirſutum, foliis lanceolatis, floribus ſeſſilibus, ramis erectis. *Lin. ſuppl.* 261.
Riſpenſtieliges Immergrün. W.
Habitat ad Cap. b. ſpei.

9. AIZOON *lanceolatum.*

A. foliis lanceolatis, floribus paniculatis. *Murr. ſyſt. veg.* 14. *p.* 471.
Aizoon paniculatum. *Sp. pl.* 700. *Mill. dict. n.* 3.
A. foliis lanceolatis ſubtus hirſutis. *Roy. lugdb.* 221.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 131.
Lanzettenförmiges Immergrün. W.
Habitat ad Cap. b. ſpei. ♂

10. AIZOON *ſarmentoſum.*

A. foliis linearibus filiformibus, panicula dichotoma, floribus ſolitariis pedunculatis. *Lin. ſuppl.* 260.
Kriechendes Immergrün. W.
Habitat ad Cap. b. ſpei

Caules *proſtrati ſarmentoſi pedales.* Folia *linearia oppoſita remota laevia, baſi approximata latiuſcula perſiſtente.* Flores *verſus ſummitatem caulis alterni axillares, pedunculo unifloro longitudine foliorum, pubeſcentes.*

996. SPIRAEA.

Gen. plant. edit. Schreb. n. 862.

Cal. 5-fidus. *Petala* 5. *Capſ.* polyſpermae.

* *Fruticoſae.*

1. SPIRAEA *laevigata.*

S. foliis lanceolatis integerrimis ſeſſilibus, racemis compoſitis. *Mant.* 244. *Willd. arb.* 370.

S. (*altaica*) foliis oblongis integerrimis ſeſſilibus, racemis terminalibus compoſitis. *Pall. roſſ.* 1. *p.* 37. *t.* 23.* *itin.* 2. *app. p.* 739. *n. III. tab. T.*

S. (*altaienſis*) foliis lanceolatis integerrimis glabris, ad baſin anguſtatis, ſeſſilibus, floribus racemoſis, racemis ſimplicibus. *Laxm. Nov. Act. Petrop. Vol. XV. p.* 555. *t.* 29. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 657.

Glatte Spierſtaude. W.

Habitat in Sibiriae *alpibus* Altaicis. ♄ (v. v.)

Frutex *ramis teretibus.* Folia *alterna, ſeſſilia, lanceolata, integerrima, obtuſiuſcula, mucronata, laeviſſima.* Racemi *terminales, ſolitarii, erecti, compoſiti e racemulis alternis, erectis,* Bractea *parva ſuffultis.* Flores *pedicello longiores, albi, alterni.* Stamina *corollis longiora.*

2. SPIRAEA *ſalicifolia.*

S. foliis oblongis ſerratis glabris, racemis decompoſitis. *Ait. Kew.* 2. *p.* 197. *Willd. arb.* 370.*

S. foliis lanceolatis obtuſis ſerratis nudis, floribus duplicato-racemoſis. *Hort. cliff.* 191. *Hort. upſ.* 131. *Roy. lugdb.* 277. *Gmel ſib.* 3. *p.* 188. *n.* 47. *t.* 39. *Roth. germ. I.* 215. *II.* 550. *Hoffm. germ.* 157. *Mill. dict. n.* 1. *Du Roi. harbk.* 2. *p.* 428. *n.* 1. *Kniph cent.* 3. *n.* 91. *Weig. obſ.* 31. *Pall. roſſ.* 1. *p.* 36. *t.* 21.* *Wangenh. amer.* 119.

α. *carnea* foliis lanceolatis, racemis ſubſpicatis, petalis carneis, cortice ramorum luteſcente. *Ait. l. c.*

S. caulibus fruticoſis, foliis ovato-lanceolatis undique ſerratis glabris, racemis duplicatis terminalibus. *Ehrh. Beitr.* 7. *p.* 136.

S. Theophraſti forte. *Cluſ. hiſt.* 1. *p.* 80.

Frutex ſpicatus, foliis ſerratis ſalignis. *Bauh. pin.* 475.

Spiraea ſalicis folio longiore ſerrato, floribus rubris.

Amm. ruth. 265. *Tournef. inſt.* 618. *Du Ham arb.* 2. *t.* 75.

β. *alpeſtris*, foliis lanceolatis obtuſis ſimpliciter ſerratis caneſcentibus, racemis ſubſpicatis breviſſimis, petalis carneis. W.

Spiraea ſalicifolia β. *Pall. roſſ.* 1. *p.* 36. *t.* 22.

S. ſalicis folio breviore latiuſculo crenato, floribus rubris. *Amm. ruth. p.* 188. *t.* 28.

γ. *paniculata*, foliis lanceolatis, racemis paniculatis divaricatis, petalis albis cortice ramorum rubro. *Ait. Kew.* 2. *p.* 198.

S. (*alba*) foliis lanceolatis acutis apice ſerratis, floribus duplicato-racemoſis. *Du Roi. harbk.* 2. *p.* 430

S. (*alba*) caulibus fruticoſis, foliis cuneato-lanceolatis, ſuperne ſerratis glabris, racemis duplicatis terminalibus. *Ehrh. Beitr.* 7. *p.* 137.

S. foliis lanceolatis acute ſerratis, floribus paniculatis, caule fruticoſo. *Mill. dict. ic.* 171. *t.* 257. *f.* 2.

δ. *latifolia*, foliis ovato-oblongis, racemis paniculatis, petalis albis, cortice ramorum rufeſcente. *Ait. l. c.*

Houttuyn Lin. Pfl. Syſt. 3. *p.* 657.

Weidenblättrige Spierſtaude. W.

Habitat in Sibiria, Tartaria β. *in alpibus* Sibiriae, γ *et* δ *in* America *boreali.* ♄ (v. v.)

3. SPIRAEA *callosa.*

S. foliis lanceolatis acutis ſerratis ſubvilloſis, panicula decompoſita ſubfaſtigiata. *Thunb. jap.* 209. *

S. (*japonica*) fruticoſa, caule pubeſcente, foliis lanceolatis acutis ſerratis nudis, corymbo terminali compoſito. *Lin. ſuppl.* 262.

Japaniſche Spierſtaude. W.

Habitat in Japonia, ♄

Differt a S. tomentoſa: foliis *villoſis, nec tomentoſis, floribus paniculatis*; *a S. ſalicifolia* :foliis *manifeſte petiolatis ellipticis acutioribus longioribus*, ſerraturis *foliorum remotis*, panicula *faſtigiata nec ſpicata*, foliis *ſubtus*, caule *et* pedunculis *villoſis.* Thunb. l. c.

4. SPIRAEA *tomentoſa.*

S. foliis lanceolatis inaequaliter ſerratis ſubtus tomentoſis, floribus duplicato-racemoſis. *Mill. dict. n.* 5. *Du Ham. arb.* 6. *Du Roi. harbk.* 2. *p.* 433. *Wangenh. amer.* 118. * *Willd. arb.* 371. *

Ulmaria pentacarpos, integris ſerratis foliis parvis ſub-

tus incanis, virginiana. *Pluk alm.* 393. *t.* 321. *f.* 5. *Raj. ſuppl.* 330. *Moriſ. hiſt.* 3. *p.* 323.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 658.
Filzige Spierſtaude. W.
Habitat in Philadelphia. *Kalm.* ♄ (v. v.)

Differt a praecedenti: Foliis *magis profunde et inaequaliter ſerratis, ſubtus valde tomentoſis adeoque caſſis,* Racemo *terminali compoſito denſiori longiori,* Floribus *demum magis rubris, et* Corollis *octuplo minoribus.* Fructus *villoſi.*

5. SPIRAEA *argentea.*

S. fruticoſa, racemis compoſitis, foliis ſericeis cuneiformibus lineatis, apice ſerratis ſubplicatis. *Lin. ſuppl.* 261.
Silberglänzende Spierſtaude. W.
Habitat in Nova Granada. *Mutis.* ♄

Frutex *ramis ſtriatis erectis, ramulis brevibus.* Folia *alterna petiolata lanceolata cuneiformia utrinque albo-ſericea lineata apice ſerrata.* Racemi *compoſiti terminales ramulis longiores.* Flores *reliquis ſpeciebus minores pentagyni, germinibus villoſis.*

*6. SPIRAEA *alpina.*

S. foliis lineari-lanceolatis denticulatis glaberrimis, corymbis lateralibus. W.
S. foliis lanceolatis ſerrulatis glabris, corymbis terminalibus. *Pall. roſſ.* 1. *p.* 35. *t.* 20.*
Spiraeae crenatae varietas. *Georgi itin. p.* 217.
Alpen-Spierſtaude. W.
Habitat in Sibiriae *alpinis ſylvaticis frigidis* Transbaicalenſibus *et circa lacum* Baical. ♄

7. SPIRAEA *hypericifolia.*

S. foliis obovatis integerrimis, umbellis ſeſſilibus. *Hort. upſ.* 131. *Mill. dict. n.* 3. *Du Roi. harbk.* 2. *p.* 434. *Wangenh. amer.* 119. *Willd. arb.* 372.*
S. foliis integerrimis, pedunculis ſimplicibus. *Hort. cliff.* 190. *Roy. lugdb.* 277.
S. hypericifolio non crenato. *Tournef. inſt.* 618. *Du Ham. arb.* 3.
Pruno ſylveſtri affinis canadenſis. *Bauh. pin.* 517. *Pluk. alm.* 408. *t.* 218. *f.* 5.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 659.
Johanniskrautblättrige Spierſtaude. W.
Habitat in Canada. ♄ (v. v.)

8. SPIRAEA *chamaedrifolia.*

S. foliis obovatis apice inciſo-dentatis, corymbis pedunculatis. *Willd. arb.* 372. *

S. foliis ovatis inciſo-ſerratis glabris, umbellis pedunculatis. *Sp. pl.* 701. *Du Roi. harbk.* 2. *p.* 432.

S. foliis oblonge ovatis, ſupra ſaepe crenatis, pedunculis ſimplicibus. *Gmel. ſib.* 3. *p.* 189.

Spiraea chamaedryos foliis. *Amm. ruth.* 269.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 659.

Gamanderblättrige Spierſtaude. W.

Habitat in Sibiria, Hungaria. ♄ (v. v.)

*9. SPIRAEA *ulmifolia.*

S. foliis ovato-lanceolatis duplicato-dentatis, corymbis pedunculatis. *Willd. arb.* 372. *

S. (*ulmifolia*) fruticoſa, foliis ovatis acutis planis duplicato-dentatis, corymbis florum terminalibus. *Scop. carn. n.* 600. *t.* 22.

Spiraea chamaedrifolia. *Jacq. hort.* 2. *t.* 140.

S. (*betulifolia*) foliis lato-ovatis ſerratis, corymbis terminalibus compoſitis faſtigiatis. *Pall. roſſ.* 1. *p.* 33. *t.* 16.

S. foliis ovatis crenatis. *Gmel. ſib.* 3. *p.* 189.

Rüſterblättrige Spierſtaude. W.

Habitat in Carniolia, Sibiria. ♄ (v. v.)

Diverſiſſima a praecedente, foliis *ovato-lanceolatis groſſe duplicato-dentatis, nec obovatis apice tantum inciſis*, floribus *majoribus*, cauleque *altiore.* W.

10. SPIRAEA *crenata.*

S. foliis obovatis acutis apice dentatis trinerviis, corymbis confertis pedunculatis. W.

S. foliis oblongiuſculis apice ſerratis, corymbis lateralibus. *Mill. dict. n.* 4. *Gouan. illuſtr.* 31? *Gmel. it.* 1. *p.* 112. *Kniph. cent.* 11. *n.* 94. *Pall. roſſ.* 1. *p.* 35. *t.* 19. *

S. hiſpanica, hyperici folio crenato. *Barr. rar.* 1376. *t.* 564?

S. hyperici folio ampliore, modo integro modo dentato. *Amm. ruth.* 270.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 660.

Gekerbte Spierſtaude. W.

Habitat in Sibiria, Hungaria. (v. ſ.)

Folia *ovato-oblonga, alia apice crenata ſ. ſerrata,*

alia integerrima. Corymbi *terminales, perfectiſſimi, albi, totum fruticem ſua copia niveum reddentes.*

Varietas hiſpanica mihi diverſa videtur. Differt haec a S. chamaedifolia, qua cum ſaepius confuſa: foliis *apice ſerratis, trinerviis, nec apice inciſis venoſis,* corymbis *confertioribus ſubcapitatis,* floribus *parum minoribus.* W.

11. SPIRAEA *triloba.*

S. foliis ſubrotundis ſubcordatis obtuſe-lobatis dentatis, umbellis pedunculatis. W.

S. foliis lobatis crenatis, umbellis terminalibus. *Mant.* 244. *Pall. roſſ.* 1. *p.* 33. *t.* 17. *

S. ramoſiſſima, parvo ſubrotundo opuli folio. *Amm. ruth.* 271.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 661.

Dreylappige Spierſtaude. W.

Habitat in Sibiriae *alpibus* Altaicis. *D. Laxmann.* ♄

Frutex *ramis teretibus.* Folia *alterna, pitiolata, ſubrotunda, lobata, obtuſa, crenata, laevia.* Umbellae *terminales, ſolitariae: Pedunculis numeroſis, unifloris, aequalibus.* Flos *pedunculatus, praeterea unus alterne ex alis foliorum ſummis.* Stamina *corollis breviora.*

*12. SPIRAEA *thalictroides.*

S. foliis obovatis obtuſis ſubtrilobis, umbellis lateralibus ſeſſilibus. W.

S. (*thalictroides*) foliis oblongis lobatis, umbellis ſeſſilibus. *Pall. roſſ.* 1. *p.* 34. *t.* 18. *

Spiraea aquilegifolia. *Pall. itin.* 3. *app.* 834. *n.* 94. *tab. P. fig.* 3.

S. foliis variis per faſciculos congeſtis. *Amman. ruth. p.* 188.

Thalictrumartige Spierſtaude. W.

Habitat in alpibus Davuricis Sibiriae. ♄

Folia *ſubtus glauca ſubtiliſſime trinervia. Habitus S. hypericifoliae ſed* foliis *diſtinctiſſima.* W.

13. SPIRAEA *opulifolia.*

S. foliis ovatis trilobis ſerratis, corymbis pedunculatis. W.

S. foliis lobatis ſerratis, corymbis terminalibus. *Syſt. veg.* 393. *Willd. arb.* 374. *

S. foliis inciſis angulatis: floribus corymboſis. *Hort.*

cliff. 190. *Gron. virg.* 55. *Roy. lugdb.* 277. *Mill. dict. n.* 2. *Willich. obſ. n.* 37. *Du Roi. harbk.* 2. *p.* 436. *Wangenh. amer.* 119.

S. opuli folio. *Tournef. inſt.* 618. *Du Ham. arb.* 5.

Evonymus virginiana, ribeſii folio, capſulis eleganter bullatis. *Comm. hort.* 1 *p.* 169. *t.* 87.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 661.

Schneeballblättrige Spierſtaude. W.

Habitat in Virginia, Canada. (v. v.)

Trigyna, arborea.

14. SPIRAEA *ſorbifolia.*

S. foliis pinnatis; foliolis uniformibus ſerratis caule fruticoſo, floribus paniculatis. *Sp. pl.* 702. *Mill. dict. n.* 6. *Gmel. ſib.* 3. *p.* 190. *t.* 46. *Du Roi. harbk.* 2. *p.* 436. *Pall. roſſ.* 1. *p.* 38. *t.* 24. * *Willd. arb.* 375. *

S. foliis pinnatis. *Amoen. acad.* 2. *p.* 352. *

S. ſorbi folio tenuiter crenato, floribus in thyrſo albis. *Amm. ruth.* 186.

β. *alpina* caule ſuffruticoſo, floribus corymboſis duplo majoribus. *Pall. roſſ.* 1. *p.* 34. *t.* 25. *

Houttuyn Lin. Pfl. Syſt. 3. *p.* 662.

Ebereſchenblättrige Spierſtaude. W.

Habitat in Sibiriae *orientalis et* Camtſchatcae *uliginoſis, varietas* β. *ad lacum* Baical. ♄ (v. v.)

* * *Herbaceae.*

15. SPIRAEA *Aruncus.*

S. foliis ſupradecompoſitis, ſpicis paniculatis, floribus dioicis. *Gmel. ſib.* 3. *p.* 192. *Crantz. auſtr. p.* 109. *Scop. carn. n.* 601. *Pollich. pal. n.* 479. *Kniph. cent.* 3. *n.* 89. *Hoffm. germ.* 173. *Roth. germ. I.* 216. *II.* 551.

S. foliis bipinnatis, floribus hermaphroditis maſculisque. *Gouan. illuſtr.* 31.

Filipendula dioica, foliis duplicato - pinnatis, pinnis quinquefoliis, trifoliis et ſimplicibus. *Hall. helv. n.* 1134.

Aruncus. *Hort. cliff.* 463. *Roy. lugdb.* 278. *Gron. virg.* 121.

Barba caprae floribus oblongis. *Bauh pin.* 163.

Barba caprae. *Cam. hort.* 26. *t.* 9.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 132.

Geisbart - Spierſtaude W.

Habitat in Auſtriae, Alvorniae, Germaniae *montanis; in* Pyrenaeis. ♃ (v. v.)

Vidit hanc plantam ſemper polygamo - monoicam. *Gouan. l. c.* R.
Trigyna. Capſula *ſubdiſperma.*

16. SPIRAEA *Filipendula.*

S. foliis pinnatis: foliolis uniformibus ſerratis, caule herbaceo, floribus corymboſis. *Mat. med* 128. *Gmel. ſib.* 3. *p.* 191. *n.* 52. *Crantz. auſtr. p.* 109 *n.* 3. *Scop carn. n.* 602. *Pollich. pal. n* 480. *Fl. dan. t.* 935. *Ludw. ect. t.* 60. *Blackw. t.* 467. *Kniph cent* 3 *n* 90. *Hoffm. germ.* 174. *Roth. germ I.* 216. *II.* 552.
Filipendula foliis pinnatis, minimis intermixtis, extrema trilobata. *Hall. helv n.* 1136
Filipendula foliis pinnatis: foliolis uniformibus. *Hort. cliff.* 191. *Fl. ſuec.* 404., 439. *Roy. lugdb.* 277.
Filipendula vulgaris. *Bauh. pin* 163.
Filipendula. *Cam. epit.* 608.
β. Filipendula minor. *Bauh. pin.* 167. *prodr.* 85.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 133.
Knollige Spierſtaude. W.
Habitat in Europae *paſcuis et ſylvis.* ♃ (v. v.)

17. SPIRAEA *Ulmaria.*

S. foliis pinnatis ſubtus tomentoſis, impari majore trilobo, lateralibus indiviſis, corymbis proliferis. W.
S. foliis pinnatis: impari majore lobato floribus cymoſis. *Fl. lapp.* 201. *Hort. cliff.* 191. *Fl. ſuec.* 405, 440. *Mat. med.* 129 *Roy. lugdb.* 277. *Oed. dan. t.* 547. *Crantz. auſtr. p.* 109. *Scop. carn n.* 603. *Pollich. pal. n.* 481. *Ludw. ect.* 23. *Kniph. cent.* 1. *n.* 84. *Hoffm. germ.* 174. *Roth. germ. I.* 216. *II.* 553.
Filipendula foliis pinnatis, pinnis acute ſerratis, minimis intermiſtis, extrema trilobata maxima. *Hall. helv. n.* 1135.
Ulmaria. *Cluſ. pann.* 700. *Blackw. t.* 465.
Barba caprae floribus compactis. *Bauh. pin.* 164.
Regina prati. *Dod. pempt.* 57.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 135.
Sumpf-Spierſtaude. W.
Habitat in Europae *pratis uliginoſis, umbroſis.* ♃ (v. v.)

*18. SPIRAEA *digitata.* W.

S. foliis pinnatis ſubtus tomentoſis, impari majore ſeptemlobo, lateralibus quinquelobis, corymbo ramoſo coarctato. W.

S. (*palmata*) herbacea, foliis pinnatis, impari palmato multifido, floribus cymoſis. *Pall. roſſ.* 1. *p.* 40. *t.* 27. * *itin.* 3. *app.* 735. *n.* 95. *tab. Q. fig.* 1.
S. folio impari majore multifido. *Gmel. ſib.* 3. *p.* 192.
Ulmaria foliis profunde laciniatis. *Amman. ruth.* 74.
Fingerblättrige Spierſtaude. W.
Habitat in ſubalpinis humidis Srbiriae *orientalis.* ♃
Similis ſequenti, tamen diſtincta: foliis *ſubtus tomentoſis*, laciniis *acuminatis*, floribus *albis in corymbo coarctato ramoſo.* W.

19. SPIRAEA *lobata.*
S. foliis pinnatis, glabris, impari majore ſeptemlobo, lateralibus trilobis, corymbis proliferis. W.
Spiraea lobata. *Murr. ſyſt. ed.* 14. *p.* 472. *Iacq. hort.* 1. *p.* 38. *t.* 88.
S. (*palmata*) foliis pinnatis, impari lateralibusque lobabatis biſerratis, floribus cymoſis. *Lin. ſuppl.* 262.
S. (*palmata*) foliis pinnato-ternatis, foliolo impari quinquelobo-palmato. *Syſt. veg. ed.* 13. *p.* 395.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 137.
Rothblütige Spierſtaude. W.
Habitat in America *boreali?* ♃ (v. v.)
Statura et facies S. Ulmariae. Radix *odorata*, Herba *inodora.* Folia *fere S. Ulmariae, foliolis lobatis.* Flores *rubri.* Fructus *S. Filipendulae nec contortus ſed rectus.*

*20. SPIRAEA *camtſchatica.*
S. foliis quinquelobis, petiolis auriculatis, caule hirſuto, corymbis proliferis. W.
S. herbacea, foliis ſimpliciter palmatis, petiolis appendulatis, caule hirſuto, floribus cymoſis. *Pall. roſſ.* 1. *p.* 41. *t.* 28. *
S. foliis multifidis angulatis, fructibus erectis hiſpidis, petiolis appendiculatis. *Gmel. ſib.* 3. *p.* 192.
Camtſchatiſche Spierſtaude. W.
Habitat in Camtſchatca *et* Beringii *inſula.* ♃
Petiolus *appendicibus ſubrotundis foliaceis ſuperne eſt inſtructus. Affinis videtur ſequenti.* W.

*21. SPIRAEA *palmata.*
S. foliis palmatis ſerratis, panicula ſupradecompoſita. *Thunb. jap.* 212.
Fünflappige Spierſtaude. W.
Habitat in Iaponia. ♃

Caulis *herbaceus ſtriatus erectus totus glaber.* Folia *alterna petiolata palmata inciſo-quinqueloba vel ſeptemloba glabra, ſubtus pallida venoſo-reticulata palmaria.* Lobi *oblongi acuminati acute et duplicato-ſerrati.* Petiolus *ſtriatus, folio quadruplo brevior.* Flores *terminales albi vel rubri paniculati.* Panicula *ſupradecompoſita. Hanc floribus rubris tetragynam inveni. Differt a S. opulifolia, cui* foliis *proxime accedit:* foliis *quinque-vel ſeptemlobatis,* ſerraturis *acutis,* panicula *decompoſita,* caule *herbaceo.* Thunb. l. c.

22. SPIRAEA *trifoliata.*

S. foliis ternatis ſerratis ſubaequalibus, floribus ſubpaniculatis. *Sp. pl.* 702. *Mill. dict. n.* 7. *et ic. t.* 256.

Filipendula foliis ternatis. *Hort. cliff.* 191, *Hort. upſ.* 131. *Roy. lugdb.* 277. *Gron. virg.* 55.

Ulmaria major trifolia, flore amplo pentapetalo, virginiana. *Pluk. alm.* 393. *t.* 236. *f.* 5. *Raj. ſuppl.* 330.

Ulmaria virginiana trifolia, floribus candidis amplis longis et acutis. *Moriſ. hiſt.* 3. *p.* 323.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 137.

Dreyblättrige Spierſtaude. W.

Habitat in Virginia, Canada. ♃ (v. v.)

POLYGYNIA.

997. ROSA.

Gen. pl. ed. Schreb. n. 863.

Petala 5. *Cal.* urceolatus, 5-fidus, carnoſus, collo coarctatus. *Sem.* plurima, hiſpida, calycis interiori lateri affixa.

* *Germinibus ſubgloboſis.* (†)

*1. ROSA *berberifolia.*

R. germinibus globoſis pedunculisque aculeatis, caule aculeis ſubgeminatis uncinatis, foliis ſimplicibus ſubſeſſilibus. W.

(†) *Tubus calycis, germinis inferi formam in Roſis mentiens, dicitur a Linnaeo perperam germen. Terminum non mutavi, quum Botanophilis hujus ſenſus pateat.* W.

R. (*berberifolia*) ſpinis recurvatis, foliis ſimplicibus ſeſſilibus, ſpinae geminae interjectis. *Pall. Nov. Act. petrop.* 10. *p.* 379. *t.* 10. *f.* 5. *

R. (*ſimplicifolia*) foliis ſimplicibus. *Saliſb. prod. ſtirp. hort. ad Chapel allerton. p.* 359. * *Juſſ. gen. ed Uſt.* 372.

Einfachblättrige Roſe. W.

Habitat in Perſia *boreali.* ♄

Caulis *pubeſcens aculeis validis uncinatis ſubgeminatis.* Folia *ab omnibus hucusque cognitis Roſis diſtincta, ſimplicia ovata breviſſime petiolata dentata glauca.* Pedunculi *in ramulis terminales breves uniflori pubeſcentes aculeisque aliquot obſiti.* Germen *globoſum aculeis numeroſis tectum.* Petala *obovata flava macula puniceа ad baſin notata.* W.

2. ROSA *lutea.*

R. germinibus globoſis pedunculisque glabris, calycibus petiolisque ſpinuloſis, aculeis ramorum rectis. *Ait. Kew.* 2. *p.* 200. *Willd. arb.* 303. *

R. (*Eglanteria*) germinibus globoſis pedunculisque glabris, caule aculeis ſparſis rectis, petiolis ſcabris, foliolis acutis. *Sp. pl.* 703. *Scholl. barb. n.* 399. *Moench haſſ. n.* 418. *Kniph. cent.* 7. *n.* 77. *Leyſ. hall. n.* 489. *Hoffm. germ.* 174. *Roth. germ. I.* 217. *II.* 553. *Retz. obſ.* 1. *p.* 20.

R. (*lutea*) caule aculeato, foliis pinnatis, foliolis ovatis ſerratis utrinque glabris; pedunculis breviſſimis. *Mill. dict. n.* 11.

R. (*lutea*) calycibus ſemipinnatis, germinibus globoſis pedunculisque glabris: foliolis ovatis glabris ſerratis, ſerraturis petiolisque villoſis glanduloſis: caule aculeato. *Du Roi. harbk.* 2. *p.* 344.

R. (*chlorophylla*) foliis obovatis duplicato-ſerratis glanduloſis glutinoſis nitide-virentibus, petiolis aculeatis, ſtipulis ſerraturis glanduloſis, pedunculis glabris, germinibus ſubgloboſis glabris, laciniis calycinis ſemipinnatis glanduliferis. *Ehrh. Beitr.* 2. *p.* 69.

Roſa foetida. *Allion pedem. n.* 1792.

Roſa lutea. *Dalech. hiſt.* 126. *Lob. ic.* 2. *p.* 209. *Tabern. hiſt.* 1495. *Bauh. hiſt.* 2. *p.* 47.

R. lutea ſimplex. *Bauh. pin.* 483. *Du Ham. arb.* 36. *Beſl. eyſt. vern.* 6. *t.* 5. *f.* 1.

β. Roſa bicolor. *Iacq. hort.* 1. *p.* 1. *t.* 1. *et* 3. *p.* 1.

R. (*punicea*) calycibus integris, germinibus globoſis glanduloſis, pedunculis glabris, foliolis ovatis glabris ſer-

ratis, ſerraturis petiolisque glanduloſis, floribus bicoloribus. *Du Roi. harbk.* 2. *p.* 347.

Roſa punicea. *Corn. canad.* 11.

R. ſylveſtris auſtriaca flore phoeniceo. *Park. theatr.* 1019.

Gelbe Roſe. W.

Habitat in Germania, Anglia. ♄ (v. v.)

Flores *flavi.* Folia *odoratiſſima.*

*2. ROSA *ſulphurea.*

R. germinibus globoſis, petiolis cauleque aculeatis, aculeis caulinis duplicibus majoribus, minoribusque numeroſis, foliis ovalibus. *Ait. Kew.* 2. *p.* 201. *Willd. arb.* 305.*

R. (*glaucophylla*) foliolis obovatis, ſimpliciter ſerratis eglanduloſis pubeſcentibus glaucis, petiolis aculeatis, ſtipulis laceratis ſerratis, ſerraturis glanduloſis, pedunculis glabris, germinibus hemiſphaericis glanduloſis, laciniis calycinis ſemipinnatis glanduliferis. *Ehrh. Beitr.* 2. *p.* 69.

R. lutea multiplex. *Bauh. pin.* 483. *Du Ham. arb.* 37. *Hort. angl. p.* 66. *t.* 18. *Knorr. del.* 1. *t. R. Du Roi. harbk.* 2. p. 346.

R. flava pleno flore. *Cluſ. cur. poſt.* 6.

Schwefelgelbe Roſe. W.

Habitat in Oriente. ♄ (v. v.)

*4. ROSA *blanda.*

R. germinibus globoſis glabris, caulibus adultis pedunculisque laevibus inermibus. *Ait. Kew.* 2. *p.* 202.

Hudſonsbay-Roſe. W.

Habitat in Terra Nova *et ſinu* Hudſonis. ♄ (v. ſ.)

Caules *adulti laeves inermes, juniores ſeu primi anni aculeis rectis ſubreflexis tenuibus armati.* Rami *teretes inermes nitidi rubicundi.* Folia *pinnata.* Foliola *plerumque ſeptem oblonga argute et ſubaequaliter ſerrata glabra.* Petioli *glabri, perumque una alterave ſpinula armati.* Ait. l. c.

5. ROSA *cinnamomea.*

R. germinibus globoſis pedunculisque glabris, caule aculeis ſtipularibus, petiolis ſubinermibus. *Sp. pl.* 703. *Mill. dict. n.* 21. *Leers. herb. n.* 381. *Hoffm. germ.* 174. *Roth. germ. I.* 217. *II.* 554.

R. calycibus integris: germinibus globoſis pedunculisque glabris, caule aculeis ſtipularibus: petiolis villo-

ſis, foliolis ſubrotundis villoſis. *Du Roi. harbk.* 2. *p.* 348.

R. (*collincola*) foliolis ovalibus obtuſis pubeſcentibus eglanduloſis, ſubtus cinereis, petiolis ſubaculeatis, pedunculis ſubſolitariis glaberrimis, germinibus globoſis glaberrimis, laciniis calycinis ſubulatis apice ſaepe foliaceis pubeſcentibus erectis. *Ehrh. Beitr.* 2. *p.* 70.

R. (*majalis*) germinibus globoſis, pedunculis petiolisque inermibus, caule aculeis geminis axillaribus, calycibus integris ſpathulatis. *Hermann Diſſ. de Roſa p.* 8. *n.* 3. *Retz. obſ.* 3. *p.* 33. *

R. (*foecundiſſima*) germinibus globoſis pedunculisque glabris, caule petiolisque aculeatis, aculeis axillaribus ovato-oblongis ſubtus villoſis. *Roth. germ. I.* 218. *II.* 557. *Du Roi. harbk.* 2. *p.* 343.

R. minor rubello flore, quae vulgo a Menſe Majo, majalis dicitur. *C. Bauh. pin.* 483. *Rupp. jen.* 139.

R. odore cinnamomi ſimplex. *Bauh. pin.* 438. *Du Ham. arb.* 33.

R. cinnamomea, floribus ſubrubentibus, ſpinoſa. *Bauh. hiſt.* 2. *p.* 39.

R. ſaxatilis flore ruberrimo. *Cam. epit.* 99.

β. R. (*fluvialis*) germinibus globoſis, pedunculis petiolisque inermibus, caule aculeis ſparſis, foliolis ovatis acutis, calycibus linearibus inciſis. *Retz. prod. ſcand. ed* 2. *n.* 619. *Fl. dan. t.* 868.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 665.

Zimmt-Roſe. W.

Habitat in Europa *auſtrali.* ♄ (v. v.)

6. ROSA *arvenſis.*

R. germinibus ſubgloboſis glabris, pedunculis hiſpidis, caule repente petiolisque aculeatis, floribus ſubſolitariis. W.

R. germinibus globoſis pedunculisque glabris, caule petiolisque aculeatis, floribus cymoſis. *Mant.* 245. *Hudſ. angl* 192. *Scholl. barb. n.* 400. *Hoffm. germ.* 176. *Roth. germ. I.* 217. *II.* 554.

R. (*Herporhodon*) caulibus plerisque repentibus, aculeis ſparſis recurvis, foliolis ſeptenis ovatis glabris, petiolis aculeatis, pedunculis hiſpidis, germinibus ſubgloboſis glabris, ſegmentis calycinis ſubpinnatifidis, ſtylis pubeſcentibus longitudine ſtaminum. *Ehrh. Beitr.* 2. *p.* 71.

R. (*ſylveſtris*) germinibus ovatis glabris, pedunculis ſcabris, calycis foliolis ovatis ſubintegris tenuique mucrone terminatis, ſtylis elongatis, caule petiolisque aculeatis. *Roth. catalect.* 1. *p.* 59. * *Herm. diſſ. Roſa. p.* 10. *n.* 5. *Pollich. pal. n.* 485.

R. ſpinis recurvis, foliis glabris ſeptenis, calycibus tomentoſis, ſegmentis ſubpinnatis, tubis longis barbatis. *Hall. helv. n.* 1102.

R. ſpinoſiſſima. *Oed. dan. t.* 398.

R. aruenſis candida. *Bauh. pin.* 484.

R. ſylveſtris altera minor, flore albo. *Raj. angl.* 3. *p.* 455.

R. canpeſtris repens alba. *Bauh. pin.* 484.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 665.

Feld-Roſe. W.

Habitat in Anglia, Suecia, Helvetia, Germania, Dania. ♄ (v. v.)

7. ROSA *pimpinellifolia.*

R. germinibus globoſis pedunculisque glabris, caule aculeis ſparſis rectis, petiolis ſcabris, foliolis obtuſis. *Sp. pl.* 703. *Pall. it.* 2. *p.* 517. *Hoffm. germ.* 176. *Willd. arb.* 308. *Retz. obſ.* 4. *p.* 27.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 666.

Pimpinellenblättrige Roſe. W.

Habitat in Europa *auſtrali.* ♄ (v. v.)

8. ROSA *ſpinoſiſſima.*

R. germinibus globoſis glabris, pedunculis hiſpidis, caule petiolisque aculeatiſſimis. *Sp. pl.* 705. *Pollich. pal. n.* 487. *Scholl. barb. n.* 401. *Mill. dict. n.* 2. *Leers. herb. n.* 384. *Grim. fl. iſen. nov. Act. A. N. C. tom.* 3. *app. p.* 323. *Hoffm. germ.* 176. *Roth. germ. I.* 217. *II.* 555.

R. ſpinis rectis, confertis, foliis novenis, glabris, pinnis et petiolis ſubſpinoſis. *Hall. helv. n.* 1106.

R. germinibus globoſis glabris, pedunculis caule petiolisque aculeis confertis. *Crantz auſt. p.* 84.

R. calycibus integris: germinibus globoſis glabris, pedunculis hiſpidis, caule petiolisque aculeatiſſimis: foliolis lanceolato-ovatis ſerratis glabris. *Du Roi. harbk.* 2. *p.* 339.

R. caule petiolisque aculeatis calycis foliolis indiviſis. *Fl. ſuec.* 407, 442.

R. campeſtris ſpinoſiſſima, flore albo odorato *Bauh. pin.* 483.

R. campeſtris odora. *Cluſ hiſt.* 1. *p.* 116.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 666.
Vielſtachliche Roſe. W.
Habitat in Germaniae, Helvetiae *collibus.* ♄ (v. v.)

Caulis *aculeis ſubulatis*, *rectis*, *horizontalibus*, *confertiſſimis*, *inaequalibus.* Foliola *minuta*, *obovata*, *novena.* Petioli *inermes.* Pedunculi *nunc inermes*, *nunc aculeati.* Fructus *globoſus*, *maturus niger!* Gemmina *baſi ſaepius aculeata* Petala *alba*, *baſi flav-ſcentia.*

Varietas *R. pimpinellifoliae ſecundum Hallerum.*

*9. ROSA *parviflora.*

R. germinibus depreſſo-globoſis pedunculisque hiſpidis, petiolis pubeſcentibus ſubaculeatis, caule glabro, aculeis ſtipularibus rectis, foliolis ellipticis, floribus ſubgeminatis. *Willd. arb* 309. *

R. (*parviflora*) caule aculeis rectis ſtipularibus, foliis pinnatis, foliolis elliptico lanceolatis, ſimpliciter ſerratis glabris, petiolis ſubaculeatis, ſtipulis ſerratis, pedunculis hiſpidis, germinibus globoſis hiſpidis, foliolis calycinis hiſpidis. *Ehrh. Beitr.* 4. *p* 21.

R. (*carolina*) calycibus ſubintegris petalis longioribus, germinibus globoſis, pedunculisque ſubhiſpidis, caule aculeis rectis ſtipularibus, petiolis glabris, foliis ovatis ſerratis glabris. *Du Roi. harbk.* 2 *p.* 355.

R. (*penſylvanica*) germinibus globoſis hiſpidis, aculeis ſtipularibus geminis, foliolis ovatis acutis medio ſerratis, ſubtus albicantibus, flore pleno. *Wangenh. amer. p.* 113.

Roſa humilis. *Marſh. arb.* 285.
Kleinblumige Roſe. W.
Habitat in America *boreali.* ♄ (v. v.)

Haec binis ſequentibus eſt ſimillima, differt vero: caule *bipedali*, petiolis *ſuperne piloſis*, floribus *geminatis. R. lucida vero eſt diſtincta:* caule 4-ſ. 5-*pedali*, petiolis *glabris*, foliis *nitidis*, floribus *geminatis. Denique R. carolina eſt diverſa:* caule 5-ſ. 6-*pedali*, foliis *opacis nunquam nitidis*, *magis lanceolatis*, *et* floribus *corymboſis.* W.

*10. ROSA *lucida.*

R. germinibus depreſſo-globoſis pedunculisque ſubhiſpidis, petiolis glabris ſubaculeatis, caule glabro, aculeis ſtipularibus rectis, foliolis oblongo-ellipticis nitidis glabris, floribus ſubgeminatis. *Willd. arb.* 310. *

R. (*lucida*) foliis pinnatis, foliolis ovato-lanceolatis obtu-

ſiuſculis groſſe ſerratis glaberrimis nitidis, petiolis ſubaculeatis glabris, corymbis paucifloris, pedunculis ſubhiſpidis, germinibus depreſſo-globoſis ſubhiſpidis, foliolis calycinis integris. *Ehrh. Beitr.* 4. *p.* 11.

R. carolina fragrans, foliis mediotenus ſerratis. *Dill. elth.* 325 *t.* 245 *f.* 316.

Glänzende Roſe. W.

Habitat in America *boreali.* ♄ (v. ſ.)

11. ROSA *carolina.*

R. germinibus globoſis pedunculisque ſubhiſpidis, petiolis piloſis ſubaculeatis, caule glabro, aculeis ſtipularibus ſubuncinatis, foliolis oblongo-lanceolatis, floribus corymboſis. *Willd. arb.* 311. *

R. germinibus globoſis hiſpidis, pedunculis ſubhiſpidis, caule aculeis ſtipularibus, petiolis aculeatis. *Sp. pl.* 703. *Wangenh. amer.* 112. *t.* 31. *f.* 71.

R. germinibus globoſis pedunculisque hiſpidis, petiolis aculeatis, caule glabro, aculeis ſtipularibus, foliis glabris. *Ait. Kew* 2. *p.* 203. (*excluſis varietatibus*).

R. (*corymboſa*) foliis pinnatis, foliolis ovato lanceolatis acutis argute ſerratis, ſubtus pubeſcentibus glaucis, petiolis ſubaculeatis piloſis, corymbis multifloris planis, pedunculis pubeſcentibus apice ſubhiſpidis, germinibus globoſis ſubhiſpidis, foliolis calycinis integris. *Ehrh Beitr.* 4. *p.* 21.

R. (*virginiana*) calycibus integris petalis longioribus, germinibus globoſis hiſpidis, pedunculis glabris, petiolis carinatis villoſis, foliolis ovatis glabris, floribus umbellatis. *Du Roi harbk.* 2. *p.* 353.

Houttuyn Lin. Pfl Syſt. 3. *p.* 667.

Caroliniſche Roſe. W.

Habitat in America *ſeptentrionali.* ♄ (v. v.)

Caulis *laevis. Aculei ſtipulares, duo.* Folia *foliolis* 7, *oblongo-ovatis, glabris, ſerratis, ſubtus pallidioribus. Petiolis aculeatis.* Pedunculi *plures, ramoſi, inermes, adſperſi pilis glandulofis* Calycis *foliola indiviſa ſubtus hiſpida.* Petala *obcordata, rubicunda.* Flos *ſerotinus.*

12. ROSA *villoſa.*

R. germinibus globoſis pedunculisque hiſpidis, caule aculeis ſparſis, petiolis aculeatis, foliis tomentoſis. *Sp. pl.* 704 *Mill. dict. n.* 3. *Leers. herb. n.* 385. *Pollich pal n.* 483 *Hoffm. germ.* 176. *Roth. germ. I.* 217. *II.* 556. *Retz. obſ.* 1. *p.* 19.

R. ſpinis rectis, foliis quinis tomentoſis, pinnis rotundis, ſpinoſis. *Hall. helv. n.* 1105.

R. calycibus ſemipinnatis: germinibus globoſis pedunculisque hiſpidis; caule petiolisque aculeatis: foliolis ovatis tomentoſis. *Du Roi. harbk.* 2. *p.* 341.

R. foliis utrinque villoſis, fructu ſpinoſo. *Flor. ſuec.* 2. *p.* 1295. *

R. pomo ſpinoſo, folio et caule hirſuto. *Bauh. hiſt.* 2. *p.* 38. *Hall.* R.

R. ſylveſtris pomifera major. *Bauh. pin.* 484. *Du Ham. arb.* 42.

β. R. (*molliſſima*) germinibus ſubgloboſis glabris pedunculis hiſpidis, caule petiolisque aculeatis, foliis tomentoſis. *Willd. prod. n.* 1237.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 667.

Weichhaarige Roſe. W.

Habitat in Europa. ♄ (v. v.)

Caulis *laevis, aculeis ſub genicula* 2. *ſ.* 4.' *approximatis.* Folia *obtuſa, tomentoſa.* Pedunculi *hiſpidi.* Germina *hiſpida.* Petala *rubra.*

*13. ROSA *rugoſa.*

R. germinibus globoſis glabris, pedunculis caule petiolisque aculeatis, foliis ſubtus tomentoſis. *Thunb. jap.* 213.

Runzliche Roſe. W.

Habitat in Japonia. ♄

Caulis *fruticoſus.* Rami *teretes ſubtomentoſi aculeati, aculeis majoribus minoribusque denſiſſimis patentibus albis.* Folia *pinnata cum impari quadrijuga.* Foliola *ovata obtuſa cum acumine ſerrata, ſupra viridia rugoſa, ſubtus tomentoſa venoſo-rugoſa pollicaria.* Petiolus *tomentoſus aculeatus, aculeis ſparſis patulis albis.* Calyx *intus tomentoſus extus hirſutus.* Germen *globoſum. Differt facile ab omnibus Roſae ſpeciebus:* Foliolis *obtuſis cum acumine rugoſis tomentoſis.* Ramis *maxime aculeato-hiſpidis.* Thunb. l. c.

*14. ROSA *provincialis.*

R. germinibus ſubrotundis, pedunculis petiolisque hiſpidis, aculeis ramorum ſparſis ſubreflexis, foliolis ovatis ſubtus villoſis, ſerraturis glanduloſis. *Ait. Kew.* 2. *p.* 204. *Willd. arb.* 314. *

R. (*provincialis*) calycibus ſemipinnatis, germinibus

globoſis, pedunculis petiolisque hiſpidis, foliis ovato-acuminatis ſubtus villoſis, ſerraturis glanduloſis. *Du harbk.* 2. *p.* 349.

β. foliis floribusque duplo minoribus. W.
Roſa provinicalis varietas. *Curt. mag.* 407.
Roſe de Champagne. *Hortulan.*
Provencer Roſe. W.
Habitat in Bohemia, Italia, Hiſpania, Gallia. ♄ (v. v.)

Varietas β. differt caule foliis et floribus duplo minoribus, ceterum convenit ſtructura omnium partium. W.

* * *Germinibus ovatis.*

15. ROSA *centifolia.*

R. germinibus ovatis pedunculisque hiſpidis. caule hiſpido aculeato, petiolis inermibus. *Sp. pl.* 704. *Mat. med. p.* 128. *Mill. dict. n.* 14. *Kniph. cent.* 1. *n.* 75. *Knorr. del.* 1. *t. R. Regn. bot. Willd. arb.* 315. *
R. crule aculeato pedunculis hiſpidis, calycibus ſemipinnatis glabris. *Sp. pl.* 1. *p.* 91.
R. calycibus ſemipinnatis: germinibus ovatis pedunculisque hiſpidis: caule hiſpido aculeato: petiolis glandulоſis: foliolis ovatis ſerratis ſubtus piloſis. *Du Roi harbk.* 2. *p.* 367.
R. multiplex media. *Bauh. pin.* 482. *Du Ham. arb.* 15.
R. centifolia batavica. 11. *Cluſ. hiſt.* 1. *p.* 114.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 669.
Garten-Roſe. W.
Habitat - - - - ♄ (v. v.)

Flos *quaſi tornatus.*

Varietates numeroſiſſimae in omnibus hortis coluntur. W.

16. ROSA *gallica.*

R. germinibus ovatis pedunculisque hispidis, caule petiolisque hiſpido-aculeatis. *Sp. pl.* 704. *Gmel. tub. p.* 148. *Blackw. t.* 82. *Regn. bot. Hoffm. germ.* 176. *Roth. germ. I.* 218. *II.* 359. *Willd. arb.* 315.
R. (*auſtriaca*) germinibus ovatis pedunculisque hiſpidis, petiolis medioque caule aculeatis. *Crantz. auſtr. p.* 86. R.
R. calycibus ſemipinnatis: germinibus ovatis pedunculisque hiſpidis; caule petiolisque hiſpido-aculeatis; foliolis ovatis ſubtus villoſis. *Du Roi. harbk.* 2. *p.* 363.

R. rubra multiplex. *Bauh. pin.* 481. *Du Ham. arb.* 2. *t.* 53.
R. rubra flore ſemipleno. *Bauh. hiſt.* 2. *p.* 34.
β. Roſa *gallica, verſicolor.*
R. praeneſtina variegata plena. *Mill. dict. t.* 221. *f.* 2.
R. verſicolor. *Bauh. pin.* 481.
Roſa praeneſtina alba et verſicolor. *Bauh. pin.* 2. *p.* 2.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 669.
Zucker-Roſe. W.
Habitat in Europa. ♄ (v. v.)
Caulis *laevis ſ. parum aculeatus.* Pedunculi *hiſpidi.* Folia *ſupra nuda, ſubacuta, ſubtus vix tomentoſa.* Germen *ovatum, baſi hiſpidum.* Flores *rubri ſ. albi.*

*17. ROSA *damaſcena.*
R. calycibus ſemipinnatis, germinibus ovatis turgidis pedunculisque hiſpidis, caule petiolisque aculeatis, foliolis ovatis acuminatis ſubtus villoſis. *Du Roi. harbk.* 2. *p.* 369. *Ait. Kew.* 2. *p.* 205. *Willd. arb.* 316. *Retz. obſ.* 3. *p.* 33.
Damaſcener Roſe. W.
Habitat in Europa *auſtrali.* ♄ (v. v.)

18. ROSA *ſempervirens.*
R. germinibus ovatis calycibus pedunculisque hiſpidis, caule petiolisque aculeatis, floribus ſubumbellatis, bracteis lanceolatis reflexis. *Ait. Kew.* 2. *p.* 205.
R. germinibus globoſis pedunculisque hiſpidis, caule petiolisque aculeatis, floribus ſubumbellatis. *Sp. pl.* 704. *Mill. dict. n.* 9. *Hoffm. germ.* 176. *Roth. germ. I.* 218. *II.* 556.
R. caule aculeato, foliis quinis glabris perennantibus. *S. pl.* 1. *p.* 92.
R. germinibus globoſis pedunculisque hiſpidis, caule petiolisque aculeatis, foliolis lanceolatis ſubcarnoſis perennantibus. *Du Roi. harbk.* 2. *p.* 358.
R. ſempervirens Iungermanni. *Cluſ. hiſt.* 2. *app. alt. Dill. elth.* 326. *t.* 246. *f.* 318.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 868.
Immergrüne Roſe. W.
Habitat in Germania. ♄ (v. v.)

19. ROSA *pumila.*
R. germinibus ovatis pedunculisque hiſpidis, petiolis cauleque aculeato, foliis ſubtus glaucis, ſerraturis glanduloſis fructibus pyriformibus. *Ait. Kew.* 2. *p.* 206.

R. germinibus ovatis, petiolis pedunculisque hiſpidis, caule ſuperne aculeatiſſimo. *Lin. ſuppl.* 262. *Iacq. auſt.* 2. *p.* 59. *t.* 198. *Allion. pedem. n.* 1852.
R. ſylveſtris pumila rubens. *Bauh. pin.* 483.
Roſa VI. pumila. *Cluſ. hiſt.* 1. *p.* 117.
Niedrige Roſe. W.
Habitat in Auſtira, Italia. ♄ (v. ſ)

Ramuli *aculeis copioſiſſimis, qui in caulibus decidunt. Fructus magni pyriformes.*

*20. ROSA *turbinata.*

R. germinibus turbinatis pedunculisque piloſis, petiolis villoſis, aculeis ſparſis recurvis. *Ait. Kew.* 2. *p.* 206.
R. (*campanulata*) caule aculeis recurvis, foliis villoſis, petiolis aculeatis, pedunculis hiſpidis, calyce hiſpido, tubo campaniformi. *Ehrh. Beitr.* 6. *p* 97.
R. inapertis floribus, alabaſtro craſſiore, francofurtenſis quibusdam. *Tournef. inſt. p.* 639. ?
Roſa francofurtana. *Munchh. hausv.* 5. *p.* 24.
Kreiſelförmige Roſe. W.
Habitat in Europa. ♄

21. ROSA *rubiginoſa.*

R. germinibus ovatis pedunculisque hiſpidis, petiolis cauleque aculeatis, aculeis recurvis, foliolis ovatis ſubtus glanduloſo-piloſis. *Ait. Kew.* 2. *p.* 206. *Willd. arb.* 317.
R. germinibus globoſis aculeatisque, aculeis recurvis, foliis ſubtus rubiginoſis. *Mant.* 564. *Iacq. auſt.* 1. *p.* 31. *t.* 50. *Hoffm. germ.* 174. *Roth. germ. I.* 218. *II.* 558. *Retz obſ.* 1. *p.* 20.
R. caule aculeis ſparſis recurviuſculis, foliis pinnatis ſubtus glanduloſo-hiſpidis rubiginoſis, foliolis ovatis duplicato-ſerratis, petiolis aculeatis, pedunculis glanduloſo-hiſpidis, germinibus ovatis ſubhiſpidis, foliolis calycinis ſemipinnatis. *Ehrh. Beitr.* 4. *p.* 22.
R. (*Eglanteria*) calycibus ſemipinnatis, germinibus globoſis glabris, pedunculis petiolisque hiſpidis glanduloſis, caule aculeis ſparſis curvis, foliolis ſubrotundis ſerraturis glanduloſis. *Du Roi. harbk.* 2. *p.* 336.
Roſa Eglanteria. *Mill. dict. n.* 4. *Gouan. hort.* 245. *Herm. diſſ. de Roſa p.* 17. *n.* 12.
Roſa ſuavifolia. *Lightfoot ſcot.* 262. *Fl. dan. t.* 870.
R. ſpinis aduncis, foliis ſubtus rubiginoſis. *Hall. helv. n.* 1103.

R. ſylveſtris foliis odoratis. *Bauh. pin.* 483.
R. ſylveſtris odorata. *Dod. pempt.* 187.
R. eglanteria. *Tabern. ic.* 1087.
R. foliis odoratis Eglantina dicta. *Bauh. hiſt.* 2. *p.* 41.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 664.
Wein-Roſe. W.
Habitat in apricis Europae. ♄ (v. v.)

Ramis *laeves, ſed aculei ſparſi majores recurvi.* Folia *ſepteno-pinnata ovata acuta ſparſa, ſubtus atomis reſinoſis purpuraſcentibus.* Petioli *hiſpidi, aculeis minutis recurvis, uti Bracteae glandulis minutis pedicellatis.* Germen *ſubgloboſum, rarius aculeolis praeſertim baſi.* Penduculus *aculeis minutiſſimis.* Flos *purpureus.*

*22. ROSA *muſcoſa.*

R. germinibus ovatis, calycibus pedunculis petiolis ramulisque hiſpidis glanduloſo-viſcoſis, ſpinis ramorum ſparſis rectis. *Ait. Kew.* 2. *p.* 207. *Willd. arb.* 318.
R. (*muſcoſa*) calycibus ſemipinnatis, germinibus ovatis, pedunculis foliorumque ſerraturis hiſpido-viſcidis, caule petiolisque aculeatis, foliolis ovatis ſerratis glabris. *Du Roi. harbk.* 2. *p.* 368. *Curt. mag.* 69.
R. rubra plena ſpinoſiſſima, pedunculo muſcoſo. *Mill. ic.* 148. *t.* 221. *f.* 1.
R. provincialis ſpinoſiſſima pedunculo muſcoſo. *Hort. angl.* 66. *t.* 18.
Moos-Rose. W.
Habitat - - - - ♄ (v. v.)

Hybrida ſine dubio proles, an a R. provinciali orta? W.

*23. ROSA *moſchata.*

R. germinibus ovatis pedunculisque villoſis, caule petiolisque aculeatis, foliolis oblongis acuminatis glabris, paniculis multifloris. *Ait. Kew.* 2. *p.* 207. *Willd. arb.* 319.
R. foliolis quinis ovatis ſerratis acutis laevibus, floribus corymboſis, calycibus oblongis, laciniis integris. *Desfont. fl. atlant.* 1. *p.* 400. *
R. calycibus ſemipinnatis glanduloſis, germinibus ovatis pedunculisque hiſpidis, caule petiolisque aculeatis, foliolis ovatis acuminatis glabris, floribus corymboſis. *Du Roi. harbk.* 2. *p.* 365.
R. (*Opſoſtemma*) caulibus erectis, foliolis ellipticis acu-

minatis glabris, coſta pubeſcente, petiolis aculeatis, corymbis multifloris, pedunculis hiſpidis, germinibus ovatis hiſpidis, laciniis calycinis ſemipinnatis ſubglanduloſis hiſpidis, longitudine petalorum, ſtylis pubeſcentibus longitudine ſtaminum. *Ehrh. Beitr.* 2. *p.* 72.

R. moſchata minor flore ſimplici. *Bauh. hiſt.* 2. *p.* 45.
Biſam-Roſe. W.

Habitat in ſepibus Barbariae. ♄ (v. v.)

Colitur a Tunetanis. Oleum eſſentiale odoratiſſimum e petalis deſtillatione obtinent. Desfont. l. c.

*24. ROSA *rubrifolia.*

R. germinibus ovatis pedunculisque glabris glauceſcentibus, petiolis aculeatis, caule aculeis ſparſis uncinatis, foliolis oblongis argute ſerratis glabris, floribus ſubcorymboſis. W.

R. (*rubrifolia*) petiolis ſpinoſis, foliolis 7-9 glabris acutis, germinibus umbellatis glabris, calycibus foliaceis. *Villars. delph.* 3. *p.* 549.

Rothblättrige Roſe. W.

Habitat in montibus Delphinatus, Helvetiae, Salisburgae. ♄ (v. ſ.)

Foliorum *venae ſubtus rubrae.* Caulis pedunculi *et* germina *rore glauco tecta.* W.

*25. ROSA *lagenaria.*

R. germinibus obovatis glabris, pedunculis petiolisque glanduloſo-hiſpidis, caule inermi, foliolis ovalibus glabris. W.

R. (*lagenaria*) germinibus oblongis collo attenuatis, foliolis novenis ellipticis ſerratis, fructibus ſubcymoſis pendulis. *Villars. delph.* 3. *p.* 553.

Keulenfrüchtige Roſe. W.

Habitat in montibus Delphinatus, Helvetiae. ♄ (v. ſ.)

Sequentibus affinis, ſed germine obovato aliisque notis diſtincta, a R. pumila diverſa: foliis *ſubtus non glaucis,* petiolis cauleque *inermibus.* W.

26. ROSA *alpina.*

R. germinibus ovatis glabris, pedunculis petiolisque hiſpidis, caule inermi. *Gmel ſib.* 3. *p.* 177. *Iacq. auſtr. t.* 279. *Willd. arb.* 320. *Hoffm. germ.* 176. *Roth. germ. II.* 559.

R. inermis, foliis ſeptenis glabris, calycis ſegmentis indiviſis. *Hall. helv. n.* 1107.

R. (*inermis*) germinibus ovatis, caule pedunculisque glabris inermibus, petiolis ſcabris. *Turra. diar. Act. p.* 128.

R. (*inermis*) caule inermi, pedunculis hiſpidis, calycis foliolis indiviſis, fructibus oblongis. *Mill. dict. n.* 6.

R. (*rupeſtris*) germinibus glabris, pedunculis hiſpidis, caule petiolisque inermibus. *Crantz. auſtr. p.* 85. *n.* 6. R.

R. campeſtris, ſpinis carens, biflora. *Bauh. pin.* 484.

R. rubello flore ſimplici non ſpinoſo. *Bauh. hiſt. p.* 39.

R. non ſpinoſa. *Hall. opuſ.* 218.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 670.

Alpen-Roſe. W.

Habitat in alpibus Helvetiae. ♄ (v. v.)

Folia *glabra.* Calyces *ſimplices.* Petala *obcordata biloba.*

*27. ROSA *pyrenaica.*

R. germinibus ovatis pedunculisque hiſpidis coloratis, petiolis hiſpido-aculeatis, calycibus omnino foliofis. *Gouan. illuſt.* 31. *t.* 19. *

Roſa hiſpida. *Krock. ſileſ. n.* 783.

Pyrenäiſche Roſe. W.

Habitat in Pyreneicis, Helveticis, Sileſiacis *alpibus.* ♄ (v. ſ.)

28. ROSA *pendulina.*

R. inermis, germinibus oblongis, pedunculis petiolisque hiſpidis, caule ramisque glabris, fructibus pendulis. *Ait. Kew.* 2. *p.* 208. *Willd. arb.* 322.

R. germinibus ovatis glabris, pedunculis cauleque hiſpidis, petiolis inermibus, fructibus pendulis. *Sp. pl.* 705. *Du Roi. harbk.* 2. *p.* 371.

R. ſanguiſorbae majoris folio, fructu longo pendulo. *Dill. elth.* 325. *t.* 245. *f.* 317.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 671.

Hängende Roſe. W.

Habitat in America *boreali.* ♄ (v. v.)

*29. ROSA *montana.*

R. germinibus oblongis pedunculisque hiſpidis, petiolis aculeatis, caule aculeis ſtipularibus uncinatis, foliolis glabris obovatis glanduloſo-ſerratis. W.

R. (*montana*) petiolis aculeatis, foliis ſeptenis ſubrotundis, germinibus hiſpidis, calycibus piſtillisque perſiſtentibus villoſis. *Villars. delph.* 3. *p.* 547. *

Habitat in montibus Delphinatus, Helvetiae ♄ (v. ſ.)

Petiolus *aculeatus pilisque glanduliferis raris praecipue baſin verſus obſitus.* Foliola *orbiculato-obovata glabra concoloria duplicato-ſerrata, ſerraturis glandulosis.* Caulis *glaber aculeis plerumque gemini uncinatis ſub foliorum ortu.* Petala *obcordata alba.* W.

*30. ROSA *multiflora.*

R. germinibus ovatis pedunculisque inermibus villoſis, caule petiolisque aculeatis. *Thunb. jap.* 214.

Habitat in Japonia. ♄

Caulis *fruticoſus erectus ramoſus.* Rami *teretes purpurei aculeati glabri erecti.* Aculei *ſparſi recurvi.* Folia *alterna petiolata pinnata.* Foliola *oppoſita, rarius alterna ſeſſilia ovata ſerrata, ſupra viridia glabra, ſubtus pallida villoſa.* Petioli *villoſi aculeati, aculeis ſparſis minutis recurvis.* Flores *terminales paniculati, magnitudine florum Rubi.* Panicula *decompoſita patens.* Pedunculi *et* Pedicelli *villoſi inermes.* Calyx *valde hirſutus villo albo, imprimis margine.* Corolla *alba.* Germen *villoſum ovatum inerme* *Differt optime ab omnibus aliis Roſis:* floribus *parvis paniculatis,* pedunculis *villoſis.* Thunb. l. c.

31. ROSA *canina.*

R. germinibus ovatis pedunculisque glabris, caule petiolisque aculeatis. *Mat. med.* 129. *Pollich. pal. n.* 186. *Fl. dan. t.* 555. *Ludw. ect. t.* 70. *Blackw. t.* 8. *Kniph. cent.* 7. *n.* 76. *Hoffm. germ.* 176. *Roth. germ. I.* 218. *II.* 560. *Willd. arb.* 321.

R. ſpinis aduncis, foliis ſeptenis, calycibus tomentoſis, ſegmentis pinnatis et ſemipinnatis, tubis breviſſimis. *Hall. helv. n.* 1101.

R. calycibus ſemipinnatis villoſis, germinibus ovatis pedunculisque glabris: petiolis ſubhiſpidis: foliolis ovatis mucronatis. *Du Roi. harbk.* 2. *p.* 359

R. caule aculeato, petiolis inermibus, calycibus ſemipinnatis. *Fl. ſuec.* 406, 441.

R. ſylveſtris vulgaris, flore odorato incarnato. *Bauh. pin.* 483.

R. canina vulgo dicta. *Dod. pempt.* 187.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 670.

Hecken-Roſe. W.

Habitat in Europa. ♄ (v. v.)

Caulis *laevis: internodiis aculeis* 2 *alternis.* Petioli

aculeati. Folia *acutiuſcula, nuda.* Pedunculi *glabri.* Germen *glabrum.* Petala *biloba, incarnata.* Bracteae 2 *oppoſitae, ciliatae.*

*32. ROSA *collina.*

R. germinibus ovatis ſubglabris, pedunculis petiolisque glanduloſo-hirſutis, caule aculeato. *Murr. ſyſt. ed.* 14. *p* 474. *Iacq. auſt.* 2. *t.* 197. *Hoſt. ſynop.* 280.

Hügel-Roſe. W.

Habitat in collibus Auſtriae. ♄ (v. ſ.)

Folia *ſubtus villoſa.* W.

*33. ROSA *parviſolia.*

R. germinibus ovatis ſubglabris, pedunculis glanduloſis, petiolis cauleque tenuiſſime aculeatis, foliolis rugoſis ſubtus villoſiuſculis ovatis glanduloſo-ſerratis. W.

R. (*parvifolia*) aculeis rectis minutis, foliis rugoſis glanduloſis, ſubtus villoſis, petiolis aculeatis, pedunculis glanduloſis, calycis tubo ovato nudiuſculo. *Ehrh. Beitr.* 6. *p.* 97.

Burgunder Roſe. W.

Habitat in Europa. ♄ (v. v.)

Frutex *pygmaeus.* Flores *parvi.* Foliola *quina parva ovata acuta.* W.

34. ROSA *ſemperflorens.*

R. germinibus oblongis pedunculisque hiſpidis, caule petiolisque aculeato-hiſpidis, foliis ſubternatis aculeatis. W.

R. (*ſemperflorens*) caule aculeato, foliis ſubternis, pedunculisque ſubunifloris aculeatis hiſpidis, calycis foliolis integris. *Curt. mag.* 284.

R. (*ſinica*) germinibus ſubgloboſis glabris, pedunculis aculeatis hiſpidis, caule petiolisque aculeatis, calycinis foliolis lanceolatis ſubpetiolatis. *Syſt. veg.* 394.?

Immerblütige Roſe. W.

Habitat in China. ♄ (v. v.)

Flores *ſaturate rubri.* Calycis *laciniae apice foliaceae, unica reliquis majore.* Germina *et* Pedunculi *interdum at rarius glabri.* W.

*35. ROSA *chinenſis.*

R. germinibus ovatis pedunculisque glabris, petiolis cauleque aculeatis, foliolis ovato-lanceolatis ſubternatis ſerrulatis glabris. W.

R. (*chinenſis*) germinibus ovatis, pedunculis, foliolisque utrinque glabris, ramis petiolisque aculeatis. *Iacq. obſ.* 3. *p.* 7. *t.* 55. *

Chineſiſche Roſe. W.

Habitat in China. ♄

Valde affinis praecedenti; an varietas? W.

36. ROSA *indica.*

R. germinibus ovatis pedunculisque glabris, caule ſubinermi, petiolis aculeatis. *Sp. pl.* 705.

R. ſubinermis, foliolis quinis ſubtus tomentoſis: impari majori, ſtipulis obſoletis. *Sp. pl.* 1. *p.* 402.

R. cheuſan glabra, juniperi fructu. *Petiv. gaz.* 57. *t.* 35. *f.* 11.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 671.

Indiſche Roſe. W.

Habitat in China. ♄

Rami *inermes; rarius armati una alterave ſpina tenuiſſima verſus folia vel in petiolis.* Folia *pinnata: foliolis quinis, ſubtus tomentoſis, ſupra glabris, ſerratis: extimo duplo majore.* Pedunculi *longi, nudi, ſimplices.* Calyx *inciſus, laevis.* Fructus *magnitudine Sorbi aucupariae.*

*37. ROSA *longifolia.* W.

R. germinibus ovatis glabris, pedunculis glanduloſo-ſubaculeatis, caule ſubinermi, petiolis aculeatis, foliolis glabris ovatis acuminatis. W.

Langblättrige Roſe. W.

Habitat in India *orientali.* ♄ (v. ſ.)

Caulis *forte magnus, glaber absque aculeis, videtur itaque ex toto inermis, ramis pedalibus.* Folia *pinnata:* Foliolis *quinis ovatis acuminatis utrinque glabris remote ſimpliciter ſerratis, lateralibus ſesquiuncialibus, terminali biunciali.* Petiolus *pilis glanduliferis ſparſis ſuperne inferne, uno alterove aculeo uncinato obſitus.* Flores *magnitudine R. caninae ſubcorymbſi.* Pedunculi *pilis apice glanduliferis hiſpidi.* Germen *oblongum glabrum.* Calycis *foliola apice foliacea acuminata dentata externe glabriuſcula, interne tomentoſa. A praecedente mihi diverſa videtur.* W.

*38. ROSA *bracteata.*

R. germinibus obovatis, pedunculis bracteatis, ramulisque villoſis, caule petiolisque aculeatis, foliolis glabris ſubrotundis crenatis ſubaculeatis. W.

R. (*bracteata*) germinibus pyriformibus ſericeis bracteatis, caule ſericeo aculeato, foliis pinnatis aculeatis, foliolis ovatis crenatis glabris ſubaculeatis, floribus ſolitariis. *Wendl. obſ. p.* 50. *

Beblätterte Roſe. W.

Habitat in China. ♄ (v. v.)

Caulis *villoſus aculeis ſparſis rectis.* Folia *pinnata, foliolis quinis ſeu ſeptenis ſubrotundo-ellipticis ſubcoriaceis obtuſis crenatis glabris, ſuperne nitidis.* Flores *fragrantes terminales ſolitarii.* Pedunculus *breviſſimus bracteis lanceolatis germen obovatum tegentibus ſericeo-villoſis, obſitus.* Calycis *foliola indiviſa ovato-lanceolata acuminata ſericeo-villoſa.* Petala *albo-flaveſcentia obcordata.* W.

39. ROSA *alba.*

R. germinibus ovatis glabris, pedunculis hiſpidis, caule petiolisque aculeatis. *Mat. med.* 129. *Mill. dict. n.* 16. *Crantz. auſtr. p.* 85. *n.* 5. *Willd. arb.* 322. *Knorr. del.* 1. *t. R.* 6. *Roth. germ. I.* 219. *II.* 561. *Retz.* 3. *p.* 34. *Hoffm. germ.* 176.

R. caule aculeato, pedunculis hiſpidis, calycibus ſemipinnatis glabris. *Sp. pl.* 1. *p.* 492.

R. calycibus ſemipinnatis: germinibus ovatis glabris: pedunculis hiſpidis: caule petiolisque villoſis aculeatis: foliolis ovatis ſubtus villoſis. *Du Roi. harbk.* 2. *p.* 361.

R. alba vulgaris major. *Bauh. pin.* 482. *Du Ham. arb.* 16.

R. ſativa. 1. *Dod. pempt.* 186.

β. R. alba flore pleno. *Beſl. eyſt. vern.* 6. *t.* 3. *f.* 1. *Blackw. t.* 73. *Kniph. cent.* 3. *n.* 76.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 672.

Weiſse Roſe. W.

Habitat in Europa, Auſtria. ♄ (v. v.)

998. RUBUS.

Gen. plant. edit. Schreb. n. 864.

Cal. 5-fidus. *Petala* 5. *Bacca* compoſita acinis monoſpermis.

* *Fruteſcentes.*

*1. RUBUS *roſaefolius.*

R. foliis quinato-pinnatis ternatisque utrinque viridi-

bus, caule petiolisque aculeatis, floribus ſolitariis. *Smith. ic. ined. faſc.* 3. *p.* 60 *t.* 60. *

Roſenblättrige Himbeere. W.

Habitat in inſula Mauritii. ♄

Foliola *ovato-lanceolata utrinque viridia ſubpiloſa.* Pedunculi *ſolitarii in ramorum apice foliis oppoſiti.* W.

*2. RUBUS *pinnatus.* W.

R. foliis quinato-pinnatis ternatisque rugoſis, utrinque glabris, caule petiolis pedunculisque aculeatis, racemo terminali. W.

Gefiederte Himbeere. W.

Habitat - - - - ♄ (v. ſ.)

Rami *villoſi virides aculeis uncinatis obſiti.* Folia *quinato-pinnata, ſuperiora ternata.* Foliola *ovato-lanceolata viridia utrinque glabra rugoſo-venoſa argute duplicato-ſerrata, ut petioli ſubtus, ad coſtam mediam aculeta.* Racemus *terminalis ſimplex.* Pedunculi *villoſi valde aculeati.* Calycis *villoſi petalis longiores. Affinis videtur ſequenti, ſed flores hermaphroditi, et racemus terminalis.* W.

*3. RUBUS *auſtralis.* †

R. fruticoſus dioicus, foliis ternatis quinato-pinnatisque, caule petiolisque aculeatis racemis axillaribus ſimplicibus. *Forſt. prodr. n.* 224.

Südliche Himbeere. W.

Habitat in Nova Zeelandia. ♄

4. RUBUS *Idaeus.*

R. foliis quinato-pinnatis ternatisque, caule aculeato, petiolis canaliculatis. *Fl. ſuec.* 408, 446. *Mat. med. p.* 130. *Crantz. auſtr. p.* 81. *Willd. arb.* 323. *Pollich. pal. n.* 488. *Scop. carn. n.* 611. *Du Roi. harbk.* 2. *p.* 376. *Hoffm. germ.* 177. *Blackw. t.* 289. *Knorr. del.* 2. *t. R.* 1. *Roth. germ. I.* 219. *II.* 562.

R. caule erecto hiſpido, foliis ternatis. *Fl. lapp.* 204. *Hort. cliff.* 192. *Hort. upſ.* 133. *Roy. lugdb.* 273.

R. caule ſpinoſo, ſubrecto, foliis quinatis et ternatis, ſubtus tomentoſis, fructibus hirſutis. *Hall. helv n.* 1108.

R. idaeus ſpinoſus. *Bauh pin.* 479. *Du Ham. arb.* 9.

β. Rubus idaeus fructu albo. *Bauh. pin.* 479. *Du Ham. arb.* 10.

γ· Rubus idaeus laevis. *Bauh. pin.* 479.
R. (*glaber*) foliis ternatis, ſubtus tomentoſis, caule glabro. *Mill. dict. n.* 4. R.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 673.
Gemeine Himbeere. W.
Habitat in Europae *lapidoſis.* ♄ (v. v.)

5. RUBUS *occidentalis.*

R. foliis ternis ſubtus tomentoſis, caule aculeato, petiolis teretibus. *Sp. pl.* 706. *Kalm. it.* 2. *p.* 283. *Mill. dict. n.* 5. *Willd. arb.* 324.
R. caule ſuberecto leviter aculeato, foliis ternatis, fructu nigro. *Gron. virg.* 2. *p.* 78.
R. idaeus, fructu nigro, virginianus. *Dill. elth.* 327. *t.* 287. *f.* 319.
R. americanus magis erectus, ſpinis rarioribus, ſtipite caeruleo. *Pluk. alm.* 325.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 675.
Schwarze Himbeere. W.
Habitat in Canada. *Kalm.* ♄ (v. v.)

Caulis *teres, laevis, aculeis recurvis, nebula caerulea obductus.* Petioli *teretes aculeis recurvis.* Folia *ternata intermedio pedicellato, ovata, inciſo-ſerrata, ſubtus albo-tomentoſa.* Racemus *terminalis.* Fructus *niger; variat etiam fructu rubro, magis acido et grato quam R. idaei.* Differt *imprimis a R. idaeo cui ſimillimus:* Foliis *ſaepius ternis, quamvis lateralia foliola ſaepe duo evadant et tum ternatis, nec pinnatis:* Foliolis *lateralibus ſaepius angulatis:* Aculeis *alternis, recurvis, nec ſparſis et confertis.* Petioli *etiam aculeati, ſed primo intuitu nebula caerulea caulis dignoſcitur.*

*6. RUBUS *triphyllus.*

R. foliis ternatis ſubtus tomentoſis, foliolis ovatis inciſis dentatis, ramis petiolis pedunculisque villoſis aculeatisque. *Thunb. jap.* 215.
Dreyblättrige Himbeere. W.
Habitat in Japonia. ♄

Caulis *ſuffruticoſus glaber aculeatus.* Rami *debiles flexuoſo-erecti aculeati villoſi.* Folia *petiolata ternata.* Foliola *obovata ſubangulata inciſa dentata, dentibus magnis ſubſetiferis, ſupra glabra, ſubtus tomentoſo-alba.* Petioli *et* Pedunculi *villoſi aculeato-hiſpidi.* Flo-

res *in ramulis terminales ſubpaniculati.* Calyces *tomentoſi hiſpidi.* Thunb. l. c.

*7. RUBUS *tomentoſus.* W.

R. foliis ternatis obovatis acutis inaequaliter dentatis utrinque tomentoſis, lateralibus ſubinciſis. W.

Filzige Himbeere. W.

Habitat in Germania, Helvetia. ♄ (v. ſ.)

Rami *glabri ſtriati aculeis uncinatis ſparſis obſiti.* Folia *ternata utrinque tomentoſa mollia.* Foliola *lateralia obovata acuta groſſe inaequaliter dentata, latere exteriore ſublobata; terminale obovatum acutum baſi integerrimum.* Aculei *parvi in petiolis, coſta foliorum et pedunculis.* Flores *albi in apice ramorum paniculati.* Calycis *foliola tomentoſa reflexa.* Petala *obovata calyce duplo longiora. A R. fruticoſo eſt diverſiſſimus nec cum R. occidentali convenit, ſub quo nomine ſpecimen accepi.* W.

8. RUBUS *hiſpidus.*

R. foliis ternatis nudis, caulibus petiolisque hiſpidiſſimis, ſtrigis rigidulis. *Syſt. veg.* 395. *Mill. dict. n.* 7.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 675.

Vielſtachliche Himbeere. W.

Habitat in Canada. *Kalm.* ♄

Caules *viticulis longis, procumbentibus lignoſis, una cum petiolis adſperſi ſetis regidis.* Folia *ternata, nuda, inciſo-ſerrata; intermedio pedicellato;* *Pedunculi etiam hiſpidi.*

9. RUBUS *parvifolius.*

R. foliis ternatis ſubtus tomentoſis, caule hirto petiolisque aculeis recurvis. *Sp. pl.* 707.

R. moluccanus parvifolius. *Rumph. amb.* 5. *p.* 88. *t.* 47. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 676.

Kleinblättrige Himbeere. W.

Habitat in India. *Osbeck.* ♄

Caules *fruticoſi, teretes, tomentoſi aculeis reflexis copioſis.* Folia *ſubtus albo-tomentoſa, venoſa, petiolis ſubtus recurvato-aculeatis.* Calyces *tomentoſi, in racemum digeſti.*

*10. RUBUS *ſanctus.*

R. foliis ternatis ſimplicibusque ſubtus tomentoſis, caule petiolisque aculeis recurvis. *Schreb. dec. p.* 15. *t.* 8. *

R. creticus triphyllus, flore parvo. *Tournef. cor.* 43.

Heilige Himbeere. ♄
Habitat in Creta, Tauria, Paläſtina. ♄

11. RUBUS *jamaicenſis.*

R. foliis quinatis ternatisve ſubtus tomentoſis, caule petiolis foliisque pubeſcentibus recurvato-aculeatis, paniculis diffuſis. *Swartz obſ.* 205.

R. foliis ternatis ſubtus tomentoſis, caule petiolis foliisque pubeſcentibus recurvato-aculeatis. *Mant.* 75.

R. aculeatus, foliis digitato-quinatis ſerratis, ſubtus argenteis. *Brown. jam.* 342.

R. foliis longioribus ſubtus molli lanugine obductis et incanis, flore et fructu minoribus. *Sloan. jam.* 2. *p.* 109. *t.* 213. *f.* 1. *Raj. dendr.* 76.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 676.

Weſtindiſche Himbeere. W.

Habitat in Jamaica *proximisque.* ♄

Differt a R. parvifolio: Caule *non incano, ſed in junioribus laeviter pubeſcente, alias nudo.* Caulis *ut in R. fruticoſo.* Foiia *ternata, ſupra laevia, ſubtus albo-tomentoſa, inciſo-ſerrata.* Aculei *in caule, petiolis et coſta foliorum.*

Differt a R. fruticoſo: Foliis *inciſo-ſerratis,* Coſtis *aculeatis,* Paniculis *terminalibus diffuſis.* Floribus *et* Baccis *parvis.* Swartz l. c.

12. RUBUS *caeſius.*

R. foliis ternatis ſubnudis: lateralibus bilobis, caule aculeato tereti. *Hort. cliff.* 192. *Fl. ſuec.* 410, 445. * *Roy. lugdb.* 273. *Mill. dict. n.* 2. *Du Roi. harbk.* 2. *p.* 375. *Scop carn. n.* 612. *Pollich. pal. n.* 489. *Willd. arb.* 325. *Hoffm. germ.* 177. *Roth. germ. I.* 219. *II.* 563.

R. caule aculeato proſtrato, foliis ternatis. *Hall. helv. n.* 1110.

R. repens, fructu caeſio. *Bauh. pin.* 479.

R. minor. *Dod. pempt.* 742.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 677.

Kriechende Himbeere. W.

Habitat in Europae *dumetis.* ♄ (v. v.)

Folia *ſubtus non tomentoſa, quamvis ſaepe pubeſcentia.* Caulis *inter ſtrigoſitatem adſperſus aculeis tenuiſſimis.*

13. RUBUS *fruticoſus.*

R. foliis quinato-digitatis ternatisque, caule petiolisque

aculeatis. *Fl. ſuec.* 409, 444. * *Mill. dict. n.* 1. *Scop. carn. n.* 613. *Pollich. pal. n.* 490. *Kniph. cent.* 5. *n.* 77. *Willd. arb.* 326. *Hoffm. germ.* 177. *Roth. germ. I.* 220. *II.* 564.

R. caule aculeato, foliis ternatis ac quinatis. *Hort. cliff.* 192. *Gron. virg.* 163. *Roy. lugdb.* 273.

R. caule ſpinoſo, ſerpente, foliis quinatis et ternatis, ſubtus tomentoſis, baccis laevibus. *Hall. helv. n.* 1109.

R. foliis ternatis et quinatis, coſta ſpinoſa, fructu nigro et laevi. *Crantz auſt. p.* 82. *n.* 2.

R. foliis quinato-pinnatis ternatisque, caule petiolisque aculeatis. *Du Roi. harbk.* 2. *p.* 372.

R. vulgaris ſ. Rubus fructu nigro. *Bauh. pin.* 479. *Du Ham. arb.* 1. *Blackw. t.* 45.

Rubus. *Cam. epit.* 751.

β. R. vulgaris major, fructu albo. *Raj. angl.* 3. *p.* 467.

γ. R. flore albo pleno. *Magn. hort.* 175.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 677.

Strauchartige Himbeere. W.

Habitat in ſepibus praeſertim maritimis Europae. ♄ (v. v.)

Caulis *ſubangulatus, longiſſimus.* Folia *vel utrinque viridia; vel ſubtus alba et tomentoſa.*

*14. RUBUS *villoſus.*

R. foliis quinatis ellipticis acuminatis argute ſerratis utrinque villoſis, caulibus petiolisque aculeatis. *Ait. Kew.* 2. *p.* 210.

Haarige Himbeere. W.

Habitat in America *boreali.* ♄

15. RUBUS *canadenſis.*

R. foliis digitatis denis quinis ternatisque, caule inermi. *Sp. pl.* 707. *Mill. icon. t.* 223.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 679.

Canadiſche Himbeere. W.

Habitat in Canada, *Kalm.*

Caulis *ſubpurpureus.* Foliola *lanceolata, utrinque nuda, tenuiſſima, argute, ſerrata.* Bracteae *lanceolatae.* Stipulae *lineares, aculeatae.*

16. RUBUS *odoratus.*

R. foliis ſimplicibus palmatis, caule inermi multifolio multifloro. *Hort. cliff.* 192. * *Hort. upſ.* 133. *Roy.*

lugdb. 274. *Mill. dict. n.* 6. *et ic. t.* 223. *Willd. arb.* 326.

R. foliis quinquelobis inaequaliter ſerratis: caule inermi multifolio multifloro. *Du Roi. harbk.* 2. *p.* 379.

R. odoratus. *Corn. canad.* 149. *t.* 159. *t.* 150. *Du Hamel. arb.* 14.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 679.

Wohlriechende Himbeere. W.

Habitat in Canada. ♄ (v. v.)

17. RUBUS *moluccanus.*

R. foliis ſimplicibus cordatis ſublobatis, caule aculeato decumbente. *Sp. pl.* 707. *Thunb. jap.* 219.

Rubus moluccanus latifolius. *Rumph. amb.* 5. *p.* 88. *t.* 47. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 680.

Moluccaniſche Himbeere. W.

Habitat in Amboina. ♄

Flores *paniculati.* Petala *calyce breviora.* Germina *hirſuta.* Styli *ſtaminibus longiores.* Folia *ſubtus tomentoſa.*

18. RUBUS *microphyllus.*

R. fruticoſus, aculeatus glaber, foliis ſimplicibus cordatis ovatis obtuſis ſublobatis, pedunculis ſolitariis unifloris. *Lin. ſuppl.* 263.

R. (*palmatus*) foliis cordatis palmato - trilobis glabris, caule petiolisque aculeatis. *Thunb. jap.* 217. *

Ki Itſigo. *Kaempf. amoen.* 787.

Lappige Himbeere. W.

Habitat in Japonia. ♄

Similis R. parvifolio, ſed ille tomento caneſcit et habet folia ſemper ternata ſubtus alba. Caules *glaberrimi teretes rarius aculeati.* Folia *parva petiolata cordata obtuſa ſubtriloba ſerrata inermia.* Petioli *et* Pedunculi *inermes.* Flores *parvi magnitudine Potentillae.*

*19. RUBUS *inciſus.*

R. foliis ſimplicibus cordatis inciſis glabris, caule erecto aculeato. *Thunb. jap.* 217.

Eingeſchnittene Himbeere. W.

Habitat in Japonia. ♄

Caulis *fruteſcens erectiuſculus glaber purpureus aculeatus, aculeis ſparſis patentibus.* Rami *cauli ſimiles,*

aculeis frequentibus. Folia *e gemma plura petiolata cordata ovata rotundata raro acuta inciſa ſublobata ſerrata nervoſa, nervis elevatis, utrinque glabra, inermia pollicaria.* Petioli *glabri recurvato-aculeati patentes longitudine foliorum.* Pedunculi *capillares inermes et aculeati glabri longitudine petiolorum.* Flores *axillares pedunculati ſolitarii.* Calyx *extus glaber, intus albo-tomentoſus.* Thunb. l. c.

20. RUBUS *japonicus.*

R. fruticoſus inermis glaberrimus, foliis ſimplicibus cordatis oblongis acuminatis duplicato-ſerratis, pedunculis ſolitariis unifloris. *Lin. ſuppl.* 263.

R. foliis ſimplicibus cordatis oblongis ſerratis, caule inermi ſuffruticoſo. *Mant.* 245.

Houttuyn Lin. Pfl. Syſt. 3. *p.* 680.

Japaniſche Himbeere. W.

Habitat in Japonia. ♄

Frutex *humilis.* Caulis *erectus. laevis, angulatus, parce ramoſus.* Folia *alterna, petiolata, ſimplicia, ovata, ſ. ovato-oblonga, acuminata, acuta, duplicato-ſerrata, utrinque laevia.* Pedunculus *filiformis, longitudine corollae, terminalis, ſolitarius.* Flos *magnitudine Fragariae:* Petalis *ſubrotundis, albis.* Calyx *parvus.*

Frutex *inermis glaberrimus.* Folia *valde acuminata exacte duplicato-ſerrata.* Petala *calyce multo majora.* Suppl. l. c.

21. RUBUS *corchorifolius.*

R. fruticoſus aculeatus tomentoſus, foliis ſimplicibus oblongis cordatis ſerratis, pedunculis ſolitariis unifloris. *Lin. ſuppl.* 263.

R. (*villoſus*) foliis ſimplicibus cordatis acutis aculeatis, caule erecto petiolisque aculeatis. *Thunb. jap.* 218. *

Corchorusblättrige Himbeere. W.

Habitat in Japonia *inter* Miaco *et* Jedo. ♄

Frutex *in caule, petiolis et raro ſub foliis aculeatus.* Caulis, Folia *ſubtus,* Petioli, Pedunculi, Calycesque *tomento cani.* Petala *calyce vix longiora.*

*22. RUBUS *elongatus.*

R. foliis ſimplicibus cordatis acuminatis duplicato-crenatis ſubtus tomentoſis, caule aculeato, calycibus obtuſis. *Smith. ic. ined. faſc.* 3. *p.* 62. *t.* 62. *

Langriſpige Himbeere. W.

Habitat in Java. ♄

Paniculae *ramis elongatis flexuoſis paucifloris.* Pedunculis *ſubgeminatis.* W.

*23. RUBUS *pyrifolius.*

R. foliis ſimplicibus ovalibus acuminatis ſerratis nudis, caule aculeato paniculato, petalis minutis. *Smith. ic. ined. faſc. 3. p. 61. t. 61.*

Birnenblättrige Himbeere. W.

Habitat in Java. ♄

Panicula *ampla e corymbis compoſita.* Petala *calyce quadruplo breviora.* W.

* * *Herbacei.*

*24. RUBUS *pedatus.*

R. foliis pedato-quinatis inciſis, pedunculis filiformibus medio bracteatis, calycibus glabriuſculis. *Smith. ic. ined. faſc. 3. p. 63. t. 63.*

Anemonenartige Himbeere. W.

Habitat in Americae *borealis tractu occidentali.* ♃

Facies *a Rubis aliena ob folia pedata, convenit potius habitu cum Anemonis et Helleboris.* W.

25. RUBUS *ſaxatilis.*

R. foliis ternatis nudis, flagellis reptantibus herbaceis. *Fl. ſuec.* 411, 447. *Iacq. vind.* 245. *Oed. dan. t.* 134. *Mill. dict. n.* 8. *Crantz. auſtr. p.* 82. *Scop. carn. n.* 614. *Pall. it.* 1. *p.* 72. *Willd. arb.* 327. *Hoffm. germ.* 177. *Roth. germ. I.* 220. *II.* 565.

R. caule herbaceo, foliis glabris, ternatis, acinis pauciſſimis. *Hall. helv. n.* 1111.

R. caule repente annuo, foliis ternatis. *Fl. lapp.* 206. *Roy. lugdb.* 274.

R. ſaxatilis alpinus. *Cluſ. pann.* 115. 116.

Chamaerubus ſaxatilis. *Bauh. pin.* 480.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 138.

Stein-Himbeere. W.

Habitat in Europae *collibus lapidoſis.* ♃ (v. v.)

26. RUBUS *arcticus.*

R. foliis ternatis, caule inermi unifloro. *Fl. ſuec.* 412. 448. * *Mat. med.* 130. *Mill. dict. n.* 9. *Gmel ſib.* 3. *p.* 179. *n.* 22. *Flor. dan. t.* 488. *Willd. arb.* 327.

R. caule unifloro, foliis ternatis. *Fl. lapp.* 207. *t.* 5. *f.* 2. *Hort. cliff.* 192. *Roy. lugdb.* 274.

R. humilis, flore purpureo. *Buxb. cent.* 5. *p.* 13. *t.* 26. *mala.*

R. trifolius humilis non ſpinoſus, ſapore et odore fragariae, fructu rubro polycocco. *Amm. ruth.* 185.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 139.

Nordiſche Himbeere. W.

Habitat in Bothnia Sueciae, Sibiria, Canada. ♃ (v. v.)

*27. RUBUS *trifidus.*

R. foliis ſimplicibus inciſo-trifidis glabris, caule inermi. *Thunb. jap.* 217.

Dreytheilige Himbeere. W.

Habitat in Japonia *prope* Quana. ♃

Fructus *ruber edulis grati ſaporis.* Caulis *herbaceus flexuoſo-erectus teres ſubſimplex purpureus glaber.* Folia *petiolata cordata ſubrotunda trifida utrinque glabra fere palmaria, lobi inciſi inaequaliter ſerrati.* Flores *e gemma cum foliis pedunculati ſolitarii, pedunculo ſimplici, rarius bifido.* Petioli *et* Pedunculi *villoſi digitales.* Calyx *albo-tomentoſus.* Thunb. l. c.

*28. RUBUS *ſtellatus.*

R. foliis ſimplicibus cordatis trilobis, rugoſo-venoſis, caule inermi unifloro erecto, petalis lanceolatis. W.

R. foliis ſimplicibus lobatis: caule inermi erecto uniflo-ro, laciniis calycinis lanceolatis acutis. *Smith. ic. ined. faſc.* 3. *p.* 64. *t.* 64. *

Sternförmige Himbeere. W.

Habitat in Americae *borealis tractu occidentali.* ♃

Flos *purpureus ut in R. arctico.* W.

*29 RUBUS *geoides.*

R. foliis ſimplicibus ternatisque obtuſis ſerratis nudis, foliolo impari maximo. *Smith. ic. ined. faſc.* 1. *p.* 19. *t.* 19.

Geumartige Himbeere. W.

Habitat in freto Magellanico. ♃

Caules *herbacei breviſſimi depreſſi.* Folia *petiolata plerumque ternata, foliolis irregulariter ſerratis obtuſis, utrinque glaberrimis ſubtus venoſis, impari maximo cordato emarginato.* Petioli *ſubvilloſi.* Flores *ſolitarii hermaphroditi, pedunculis brevibus craſſiſſimis pubeſcentibus, petalis ſubrotundis. A R. Dalibarda differt:* foliis *ternatis glabris ſerratis potius quam crenatis,* pedunculis *brevibus craſſis nec filifor-*

mibus longiſſimis, petalis *ſubrotundis nec lanceolatis.* Smith l. c.

30. RUBUS *Chamaemorus.*

R. foliis ſimplicibus lobatis, caule inermi unifloro *Fl. ſuec.* 414, 449. *Mat. med.* 130. *Mill. dict. n.* 10. *Gunn. norv. n.* 6. *Lightf. ſcot.* 1. *p.* 266. *t.* 13. *Retz. obſ.* 1. *p.* 20.

R. caule bifolio unifloro, foliis ſimplicibus. *Fl. lapp.* 208. *t.* 5. *f.* 1. *Hort. cliff.* 192. *Roy. lugdb.* 279.

R. paluſtris humilis. *Tournef. inſt.* 615.

Chamaerubus foliis ribes. *Bauh. pin.* 480.

Morus norvegica. *Till. aboenſ.* 47. *t.* 150. *Fl. dan. t.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 140.

Sumpf-Himbeere. W.

Habitat in Sueciae, Sibiriae, Daniae *paludibus uliginoſis, turfoſis, frequens.* ♃ (v. ſ.)

Herba *dioica*: Radix *et Maris et Faeminae unita ſub terra in monoicam, obſervante D. C. Solandro.*

31. RUBUS *Dalibarda.*

R. foliis ſimplicibus cordatis indiviſis crenatis, ſcapo aphyllo unifloro. *Sp. pl.* 708. *Mill. dict. n.* 11. *Smith. ic. ined. faſc.* 1. *p.* 20. *t.* 20.*

Dalibarda repens. *Sp. pl.* 1. *p.* 401.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 141.

Trockene Himbeere. W.

Habitat in Canada. *Kalm.* ♃

Radix *repens.* Stolones *repentes, herbacei.* Folia (*Violae*) *cordata, obtuſa, crenata, venoſa, pilis raris adſperſa, ſubtus nuda.* Petioli *piloſi, longitudine foliorum.* Scapi *nudi ſ. aphylli, piloſi, longitudine foliorum, uniflori.* Calyx 5-*partitus.* Corolla *alba.* Semina 5, *ovata, nuda.* Stamina *longitudine corollae. Anne* Flores *maſculi et feminei in eadem planta*? Habitus *ſuadet conjunctionem cum Rubo, imprimis Chamaemoro, vel utraque diſtinguenda a Rubis.*

999. FRAGARIA.

Gen. pl. ed. Schreb. n. 865.

Cal. 10-fidus. *Petala* 5. *Receptaculum* ſeminum ovatum, baccatum, deciduum.

1. FRAGARIA *veſca.*

F. calyce fructus reflexo, pubeſcentia petiolorum patentiſſima, pedunculorum adpreſſa. W.

F. flagellis reptantibus. *Hort. cliff.* 192. *Hort. upſ.* 133. *Fl. ſuec.* 414, 450. *Mill. dict. n.* 1. *Gmel. ſib.* 3. *p.* 180. *Scop. carn. n.* 625. *ſub* Potentiſla. *Pollich. pal. n.* 491. *Ludw. ect. t.* 136. *Blackw. t.* 77. *Kniph. cent.* 3. *n.* 38. *Knorr. del.* 1. *t. E.* 1. *Hoffm. germ.* 178. *Roth. germ. I.* 221. *II.* 568.

F. (*vulgaris*) calyce fructus reflexo, pubeſcentia pedunculi adpreſſa. *Ehrh. Beitr.* 7. *p.* 21.

F. foliis ternatis: flagellis reptantibus. *Hall. helv. n.* 1112.

F. flagellis reptans, fructu eduli. *Crantz. auſtr. p.* 81. *n.* 16.

F. vulgaris. *Bauh. pin.* 326. *Fl. lapp.* 209. *Gron. virg.* 56.

β. F. fructu albo. *Bauh. pin.* 326. *Gmel. l. c.*

F. ferens fraga alba. *Bauh. hiſt.* 2. *p.* 394.

γ. F. bis fructum ferens. *Bauh. pin.* 327.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 142.

Wilde Erdbeere. W.

Habitat in Europae *ſylvis ſiccis ſterilibus.* ♃ (v. v.)

Folia *utrinque pubeſcentia, ſerratura terminalis proximis paullo longior.* Ehrh. l. c.

*2. FRAGARIA *elatior.*

F. calyce fructus reflexo, pubeſcentia pedunculorum et petiolorum patentiſſima. W.

F. (*elatior*) calyce fructus reflexo, pubeſcentia pedunculi patentiſſima. *Ehrh. Beitr.* 7. *p.* 23.

Fragaria veſca β. pratenſis. *Sp. pl* 709.

F. ſcabra flore foemineo, fructu purpureo moſchato. *Du Ham. arb.* 1. *p.* 185. *t.* 8.

F. fructu parvi pruni magnitudine. *Bauh. pin.* 327.

Fraga fructu magno. *Beſl. eyſt. ve. n. O.* 7. *t.* 8. *f.* 5.

Fragaria et Fraga. *Lob. ic.* 1. *t.* 697. *Dod. pempt.* 672.

β. F. (*muricata*) caule erecto ſuffruticoſo, foliis hirſutis. *Sp. pl.* 709.

F. arborea flore herbaceo. *Zanon. hiſt.*

F. major veſca flore herbaceo. *Moriſ. hiſt.* 2. *p.* 186.

Garten-Erdbeere. W.

Habitat in America? ♃ (v. v.)

Folia *utrinque pubeſcentia, ſerratura terminalis proximis paullo longior.* Ehrh. l. c.

*3. FRAGARIA *virginiana.*

F. calyce fructus patente, pubeſcentia petiolorum erecta, pedunculorum adpreſſa, foliis ſupra glabriuſculis. W.

F. (*virginiana*) calyce patente, foliis ſupra fere glabris, ſerraturis acuminatis. *Ehrh. Beitr.* 7. *p.* 24.

Fragaria veſca δ. virginiana. *Ait. Kew.* 2. *p.* 211.

F. (*virginiana*) foliis oblongo-ovatis ſerratis inferne incanis, calycibus longioribus, fructu ſubrotundo. *Mill. dict.*

F. glabra, fructu coccineo majore, ſeminibus in cortice loculoſo altius depreſſis, virginiana. *Du Ham. arb.* 1. *p.* 181. *t.* 5.

F. virginiana fructu coccineo. *Moriſ. hiſt.* 2. *p.* 186.

Virginiſche Erdbeere. W.

Habitat in Virginia, Carolina. ♃ (v. v.)

Serratura *foliorum terminalis proximis ſubaequat.* Ehrh. l. c.

*4. FRAGARIA *grandiflora.*

F. calyce fructus erecto, pubeſcentia pedunculorum et petiolorum erecta, foliis ſupra glabriuſculis. W.

F. (*grandiflora*) calyce fructus erecto, foliis obtuſe ſerratis ſupra fere glabris. *Ehrh. Beitr.* 7. *p.* 25.

Fragaria veſca ε. ananas. *Ait. Kew.* 2 *p.* 212.

F. foliis ovatis crenatis nervoſis, calycibus maximis. *Mill. ic.* 192. *t.* 288.

F. flore ampliſſimo, fructu ananae ſaporem et odorem referente. *Du Ham arb.* 1. *p.* 183. *t.* 6.

Ananas-Erdbeere. W.

Habitat in Surinamo. ♃ (v. v.)

Serraturae *foliorum ovatae, terminalis proximis brevior.* Ehrh. l. c.

*5. FRAGARIA *chiloenſis.*

F. calyce fructus erecto, pubeſcentia petiolorum et pedunculorum patentiſſima, foliis utrinque villoſis. W.

F. (*chiloenſis*) calyce fructus erecto, foliis crenato-ſerratis, utrinque pubeſcentibus. *Ehrh. Beitr.* 7. *p.* 26.

F. (*chilenſis*) fructu maximo, foliis carnoſis hirſutis. *Molin. chil. ed germ. p.* 112.

Fragaria veſca γ. chiloenſis. *Sp. pl.* 709.

F. chiloenſis fructu maximo, foliis carnoſis hirſutis. *Dill. elth.* 145. *t.* 120. *f.* 146.

F. pubeſcens flore ampliſſimo, fructu maximo chiloenſis. *Du Hamel.* 1. *p.* 176. *t.* 3.

Chiliſche Erdbeere. W.

Habitat in America *auſtrali.* ♃

Serraturarum *latitudo longitudinem ſuperat.* Ehrh. l. c.

*6. FRAGARIA *collina.*

F. calyce fructus erecto, pubeſcentia pedunculorum erecta, petiolorum patentiſſima, foliis utrinque pubeſcentibus. W.

F. (*collina*) calyce fructus erecto, foliis acute ſerratis utrinque pubeſcentibus. *Ehrh. Beitr.* 7. *p.* 26.

F. fructu majori candicante oblongiuſculo. *Volk. norimb. p.* 172.

F foliis hiſpidis. *Bauh. pin. p.* 327. *Mapp. alſat.* 110. *Rupp. jen. ed. Hall.* 108.

Fraga minor ſ. nigra. *Thal. herc.* 43.

Fraga partim rubra partim candida. *Trag. hiſt.* 500.

Harte Erdbeere. W.

Habitat in collibus ſterilibus ſiccis Germaniae, Helvetiae. ♃

Serraturae *ovato-lanceolatae, extima proximis brevior.* Ehrh. l. c.

7. FRAGARIA *monophylla.* †

F. calyce fructus patente dentato, pubeſcentia petiolorum et pedunculorum erecta, foliis ſimplicibus. W.

F. foliis ſimplicibus. *Syſt. nat.* 13. *p.* 349. *Uſt. neue Annalen* 8. *Stück. p.* 40. *t.* 1.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 145.

Einblättrige Erdbeere. W.

Habitat - - - - ♃ (v. ſ.)

Miſit D. Anton. Nic. Ducheſne. *An varietas F. veſcae? folium enim primum ternatum eſt, ſed* Scapus *longior;* Petala *minora;* Calyces *inciſi. Vide* Ducheſne *monograph. Fragariae.*

Folia *utrinque pubeſcentia groſſe ſerrata, ſerratura terminalis magnitudine reliquarum.* W.

8. FRAGARIA *ſterilis.*

F. petalis obcordatis calycis longitudine, receptaculo ſicco, pubeſcentia petiolorum et pedunculorum patentiſſima, caule decumbente. W.

F. caule decumbente, ramis floriferis laxis. *Pollich. pal. n.* 492. *Neck. gallob. p.* 223. *Moench. haſſ. n.* 430. *Doerr. naſſ. p.* 111. *Hoffm. germ.* 178. *t.* 6.

F. caule proſtrato, foliis ternatis, retuſis, ſericeis. *Hall. helv. n.* 1113.

F. caule decumbente repente. *Roy. lugdb.* 274. *Dalib. pariſ.* 14. *Sauv. monſp.* 177. *Hudſ. angl.* 195.

F. ſterilis. *Bauh. pin.* 327.
Comarum fragarioides. *Roth. germ. I.* 224. *II.* 577.
F. ſterilis ampliſſimo folio et flore, petalis florum cordatis. *Vaill. pariſ.* 55. *t.* 10. *f.* 1.
F. ſterilis ſ. minime veſca hirſuta minime incana. *Moriſ. hiſt.* 2. *p.* 186. *f.* 2. *t.* 19. *f.* 5.
F. ſylveſtris minime veſca ſ. ſterilis. *Lob. ic.* 698.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 145.
Trockene Erdbeere. W.
Habitat in Anglia, Helvetia, Germania. ♃ (v. ſ.)
Surculi *craſſi, depreſſi, tecti* Stipulis *lanceolatis, ferrugineis.* Folia *ternata, obovata, ſerrata, retuſa, laxiora, piloſa, ſericea, ſubtus albida: petiolis valde piloſis.* Caulis *floriferi filiformes, decumbentes, laxi, foliis raris minoribus.* Flores *ſolitarii, pedunculati, albi.* Flagella *nulla.*

1000. POTENTILLA.

Gen. pl. ed. Schreb. n. 866.

Cal. 10-fidus. *Petala* 5. *Semina* ſubrotunda, nuda, receptaculo parvo exſucco affixa.

* *Foliis pinnatis.*

1. POTENTILLA *fruticoſa.*

P. foliis pinnatis, caule fruticoſo. *Hort. cliff.* 193. *Fl. ſuec.* 416, 451. *Roy. lugdb.* 275. *Hudſ. angl.* 195. *Kalm. it.* 2. *p.* 263. *Mill. dict. n.* 1. *Gmel. ſib.* 3. *p.* 180. *n.* 26. *Du Roi. harbk.* 2. *p.* 153. *Kniph. cent.* 5. *n.* 71. *Willd. arb.* 235.
Pentaphylloides rectum fruticoſum eboracenſe. *Moriſ. hiſt.* 2. *p.* 193. *ſ.* 2. *t.* 93. *f.* 5. *Hortul. angl.* 54. *t.* 54. *Du Ham. arb.* 2. *t.* 20.
Pentaphylloides rectum fruteſcens. *Walth. hort.* 95. *t.* 17.
Pentaphylloides fruticoſa elatior, minus hirſuta. *Amm. ruth.* 88. *n.* 114. *t.* 17.
β. Pentaphylloides fruticoſa humilior lanuginoſa et incana. *Amm. ruth.* 89. *n.* 115. *t.* 18. *f.* 1.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 147.
Strauchartiges Fingerkraut. W.
Habitat in Eboraco Angliae, Oelandia *auſtrali,* Sibiria. ♄ (v. v.)

2. POTENTILLA *Anſerina.*

P. foliis interrupte pinnatis ſericeis, foliolis argute ſerratis, caule repente, pedunculis unifloris. W.

P. foliis pinnatis ſerratis, caule repente, pedunculis unifloris. *Fl. lapp.* 210. *Fl. ſuec.* 415, 452. *Hort. cliff.* 193. *Mat. med.* 131. *Roy. lugdb.* 275. *Gunn. norv. n.* 58. *Pollich. pal. n.* 493. *Scop. carn. n.* 615. *Flor. dan. t.* 544. *Hoffm. germ.* 178. *Roth. germ. I.* 221. *II.* 569.

Fragaria caule repente, foliis pinnatis ſerratis, tomentoſis, petiolis longis unifloris. *Hall. helv. n.* 1126.

Fragaria Anſerina. *Crantz auſtr. p.* 71.

Argentina. *Blackw. t.* 6. *Dodon. pempt.* 600.

Potentilla. *Bauh. pin.* 321. *Cam. epit.* 758.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 148.

Gänſerich-Fingerkraut. W.

Habitat in Europae *paſcuis argilloſis.* ♃ (v. v.)

3. POTENTILLA *ſericea.*

P. foliis bipinnatifidis multijugis utrinque tomentoſis, ſegmentis parallelis, approximatis, caule decumbente. W.

P. foliis bipinnatis utrinque tomentoſis: ſegmentis parallelis aproximatis, caulibus decumbentibus. *Sp. pl.* 710.

P. foliis pinnatis hirſutis, pinnis tredecim ovatis crenatis decurrentibus. *Hall. goett.* 2. *p.* 106.

P. foliis duplicato-pinnatis: pinnulis linearibus integerrimis brevibus. *Gmel. mſſ.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 149.

Seidenartiges Fingerkraut.

Habitat in Sibiria. ♃ (v. ſ.)

Habitus *foliorum P. Anſerinae argenteae, quamvis minima. Caulis vero et fructificationes P. vernae.* Radix *ſquamis foliorum ruderibus nigris veſtita.* Folia *pinnata ex* 13 *circiter foliolis ovatis, tomentoſis, ſericeis, pectinato-pinnatifidis, et parallelis.* Pedunculi *piloſi.* Caules *foliis duplo ſ. triplo longiores, decumbentes absque ſarmentis, nec repentes, in medio folio unico minori magisque divaricato inſtructi, teretes, apice* Floribus *aliquot luteis, alternis, pedunculatis.*

Folia *accedunt magis ad P. Anſerinam, quam ad P. multifidam. A P. multifida eſt diverſa:* foliis *radicalibus pinnatis ſexjugis, nec quadrijugis, laciniis foliorum breviſſimis approximatis,* petiolis *longiſſime et valde villoſis.* W.

4. POTENTILLA *multifida.*

P. foliis bipinnatifidis quadrijugis ſuperne glabris, ſubtus tomentoſis, ſegmentis diſtantibus; caule decumbente. W.

P. foliis bipinnatis: ſegmentibus integerrimis diſtantibus ſubtus tomentoſis, caule decumbente. *Gmel. ſib.* 3. *p.* 182. *n.* 30. *Kniph. cent.* 2. *n.* 62.

Fragaria foliis ſubtus tomentoſis, pinnatis, pinnis primis pinnatis, ultimis ſimplicibus. *Hall. helv. n.* 1125.

Pentaphylloides ſupina minor, foliis alatis hirſutis varie diſſectis. *Amm. ruth. p.* 92. *n.* 117.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 150.

Vielſpaltiges Fingerkraut. W.

Habitat in Sibiria, Tataria, Cappadocia, Helvetia. ♃ (v. v.)

Habitus *multa indicat affinia cum P. argentea.* Folia *ex ſeptem foliolis: ſegmentis linearibus ſ. lanceolatis, quinis ſ. ſummum ſeptenis, alternis, at in impari oppoſitis, a ſe invicem hiantibus ſive apice remotis, ſubtus tomentoſis.*

*5. POTENTILLA *verticillaris.*

P. foliis pinnatis ſubtus tomentoſis, foliolis linearibus margine revolutis geminatis, terminali trifido, caule erectiuſculo. W.

Potentilla verticillaris. *Stephan. in litt.*

Potentilla multifida. *Pall. itin.* 3. *p.* 194. *not.*

Pentaphylloides humile, foliis anguſtiſſimis. *Amm. ruth. p.* 86. *n.* 113. *t.* 16.

Wirtelblättriges Fingerkraut. W.

Habitat in Sibiria. ♃ (v. ſ.)

Diſtinctiſſima ab omnibus hucusque cognitis Potentillis: foliis *ſimpliciter pinnatis, foliolis geminatis ut in nonnullis Aſtragalis et Phacis, unde foliola quaſi verticillata.* Foliola *ceterum ſunt anguſta linearia ſuperne viridia villoſa inferne niveo-tomentoſa.* Calyces Pedunculi *ut et* caulis *tomentoſo-villoſi.* Corolla *lutea.* W.

6. POTENTILLA *fragarioides.*

P. foliis pinnatis ternatisque: extimis majoribus, flagellis reptantibus. *Sp. pl.* 710.

P. foliis pinnatis ternatisque hirſutis utrinque et viridibus: lobulis acceſſoriis. *Gmel. ſib.* 3. *p.* 182. *n.* 31. *t.* 34. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 151.

Erdbeerartiges Fingerkraut. W.
Habitat in Sibiria. ♃

Facies *Fragariae ſterilis.* Radix *ſubtuberoſa.* Folia *ſericea petiolis piloſis, pinnata ex* 3. 5. *vel* 7. Foliolis *ovatis, oppoſitis, ſerratis, baſi quo propioribus eo minoribus.* Stolones *decumbentes.*

*7. POTENTILLA *ruthenica.* W.

P. foliis radicalibus pinnatis, caulinis ternatis, foliolis ovatis dentatis, petalis calyce brevioribus. W.
Pentaphylloides erectum fragariae folio. *Moriſ. hiſt.* 2. *p.* 193. *ſ.* 2. *t* 20. *f.* 2.
Ruſſiſches Fingerkraut. W.
Habitat in Sibiria. ♃ (v. v.)

Saepius hanc plantam ſub nomine praecedentis ſpeciei in hortis vidi. Caulis *erectus pedalis bipedalisque dichotomo-ramoſus.* Folia *radicalia ſeu caulina inferiora pinnata e foliolis quinis obovatis profunde dentatis, binis ſuperioribus ſeſſilibus baſi decurrentibus, binis inferioribus breve petiolatis.* Folia *caulina longe petiolata ternata, foliolis oblongo-obovatis profunde dentatis.* Stipulae *oblongae margine exteriore dentatae.* Pedunculi *uniflori in dichotomia caulis ſolitarii.* Petala *flava ſubrotunda calyce minora.* W.

8. POTENTILLA *rupeſtris.*

P. foliis pinnatis alternis: foliolis quinis ovatis crenatis, caule erecto. *Hort. cliff.* 193. *It. wgoth.* 63, 76. *Fl. ſuec.* 2. *p.* 453. *Pall. it.* 2. *p.* 655. *Iacq. auſtr. t.* 114. *Scop. carn n.* 616. *Gouan. illuſtr.* 32. *Crantz. auſtr. p.* 72. *ſub* Fragaria. *Pollich. pal. n* 494. *Kniph. cent.* 11. *n.* 88. *Doerr. naſſ. p.* 186. *Hoffm. germ.* 178. *Roth. germ. I.* 222. *II.* 570.
Fragaria foliis pinnatis, pinnis ſeptenis, ſubrotundis, ſerratis. *Hall. helv. n.* 1127.
Pentaphylloides erectum. *Bauh. hiſt.* 2. *p.* 398. *Moriſ. hiſt.* 2. *p.* 192. *ſ.* 2. *t.* 20. *f.* 1.
Quinquefolium fragiferum. *Bauh. pin.* 326. *Cluſ. hiſt.* 2. *p.* 107. *n.* 5.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 151.
Felſen-Fingerkraut. W.
Habitat ad latera montium Weſtrogothiae, Sibiriae, Germaniae. ♃ (v. v.)

Flores *albi.* Folia *quina, ſeptena, novena. Gou. l. c.*

9. POTENTILLA *bifurca.*

P. foliis pinnatis ſubaequalibus: foliolis oblongis ſubbifidis: extimis confluentibus. *Sp. pl.* 711. *Gmel. it.* 1. *p.* 149. *t.* 27. *f.* 1.

Pentaphylloides orientale ſupinum, foliis alatis bifidis vel trifidis. *Tournef. cor.* 21.

Pentaphylloides repens foliis pinnatis. *Buxb. cent.* 1. *p.* 30. *t.* 49. *f.* 1.

Pentaphylloides ſupinum minus folio glabro non ſerrato. *Amm. ruth.* 116.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 152.

Spaltblättriges Fingerkraut. W.

Habitat in Sibiria, Armenia. ♃ (v. ſ.)

10. POTENTILLA *pimpinelloides.*

P. foliis pinnatis, foliolis ſubrotundis dentatis aequalibus, caule erecto. *Sp. pl.* 711.

Pentaphylloides orientale erectum, folio et facie pimpinellae. *Tournef. cor.* 21. *Buxb. cent.* 1. *p.* 30. *t.* 48.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 152.

Bibernellblättriges Fingerkraut. W.

Habitat in Armeniae *ſaxoſis.* ♃ (v. ſ.)

Planta *ſexuncialis erecta.* Folia *longe petiolata pinnata* 6-8-*juga, foliolis ſubrotundis groſſe dentatis.* Stipulae *profunde* 3-*ſ.* 4-*partitae, laciniis lineari-lanceolatis acuminatis.* W.

*11. POTENTILLA *cicutariaefolia.* W.

P. foliis pinnatis glabriuſculis, foliolis oblongo-cuneiformibus profunde dentatis, ſummis confluentibus, caule erecto, W.

Pentaphylloides orientale erectum, foliis alatis Geranii cicutariae folio facie. *Tournef. cor.* 21.

Schierlingartiges Fingerkraut. W.

Habitat in Galatia. ♃ (v. ſ.)

Radix *lignoſa ſimplex multiceps.* Caulis 4-*ſ.* 5-*uncialis erectus.* Folia *petiolata pinnata glabra, venis et petiolis pubeſcentibus.* Foliola 4-*ſ.* 5-*juga oblonga profunde dentata, ſumma baſi decurrentia.* Stipulae *apice dentatae nervoſae.* Corymbus *terminalis dichotomus.* Folia *iis Erodii cicutarii ſimillima. Differt a praecedente,* foliorum *forma, et ſummis confluentibus, denique* corymbo *dichotomo.* W.

12. POTENTILLA *penſylvanica.*

P. foliis decurſive pinnatis, ſummis ternatis, foliolis profunde dentatis, caule erecto pubeſcente. W.

P. foliis inferioribus pinnatis, ſuperioribus ternatis: foliolis inciſo-ſerratis, caule erecto pubeſcente. *Mant.* 76. *Iacq. hort. t.* 189. *Gmel. ſib.* 3. *t.* 34. *f.* 1.

Pentaphylloides canadenſe, foliis agrimoniae. *Boerh. lugdb.* 1. *p.* 40.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 153.

Penſylvaniſches Fingerkraut. W.

Habitat in Canada, Sibiria. ♃ (v. v.)

Folia Radicalia *pinnata: foliolis oblongis, obtuſiuſculis, ſerrato-dentatis, lineatis, mollibus, villoſis: extimis tribus connatis majoribus; interioribus minoribus alternis.* F. Caulina *ſubdigitata, ſeptena, ut P. recta, ſed pinnatifida.* Petala *vix calyce majora.* Habitus *P. rectae.* H. V.

Planta ſibirica magis eſt glabra; ſed foliorum florumque ſtructura eadem ac in americana. W.

13. POTENTILLA *ſupina.*

P. foliis pinnatis, foliolis oblongis profunde dentatis, caule decumbente dichotomo, pedunculis axillaribus ſolitariis. W.

P. foliis pinnatis, caule dichotomo decumbente. *Hort. cliff.* 193. *Hort. upſ.* 133. *Roy. lugdb.* 275. *Reyg. ged.* 1. *p.* 133. *n.* 2. *Gmel. it.* 1. *p.* 149. *t.* 27. *f.* 2. *Crantz. auſtr. p.* 73. *ſub* Fragaria. *Pollich. pal. n.* 495. *Hoffm. germ* 178. *Roth. germ. I.* 222. *II.* 571.

Pentaphyllum alpinum minus ſupinum. *Pluk. alm.* 285. *t.* 106. *f.* 7.

Pentaphyllum ſupinum quorundam, potentillae facie. *Cluſ. hiſt.* 2. *p.* 107.

Pentaphylli ſpecies flore luteo enneaphyllos hirſuta. *Ambroſ. phyt. t.* 414.

Quinquefolio fragifero affinis. *Bauh pin.* 326.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 153.

Niedriges Fingerkraut. W.

Habitat in Sibiria, Germania, Auſtria. ⊙ (v. v.)

* * *Foliis digitatis.*

14. POTENTILLA *recta.*

P. foliis ſeptenatis quinatisque foliolis lanceolatis groſſe dentatis, petalis obcordatis calyce majoribus, caule erecto. W.

P. foliis ſeptenatis lanceolatis ſerratis utrinque ſubpiloſis, caule erecto. *Sp. pl.* 711. *Mill. dict. n.* 2. *Iacq. auſtr. t.* 383. *Doerr. naſſ. p.* 186. *Scop. carn. n.* 618. *Hoffm. germ.* 179. *Roth. germ. I.* 222. *II.* 571.

Fragaria caule erecto, foliis quinis et ſeptenis, acute ſerratis. *Hall. helv. n.* 1121.

Fragaria foliis ſeptenis quinatisque, ſerratis utrinque piloſis: caule erecto corymboſo. *Crantz. auſtr. p.* 79.

Potentilla foliis digitatis, caule erecto corymboſo. *Hort. upſ.* 133. * *Roy. lugdb.* 275.

Quinquefolium rectum luteum. *Bauh. pin.* 325.

Quinquefolium alterum vulgare. *Dod. pempt.* 116.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 154.

Gerades Fingerkraut. W.

Habitat in Italia, Narbona, Helvetia, Auſtria, Germania *ad muros et ad agrorum margines.* ♃ (v. v.)

Folia *utrinque piloſa; at adulta fere nuda.*

Folia *radicalia et inferiora caulina ſeptenata, ſuperiora quinata, ſumma ternata utrinque pilis raris obſita, interdum fere glabra.* Petala *magna obcordata pallide flava ad albidum colorem vergentia, calyce majora.* W.

*15. POTENTILLA *obſcura.* W.

P. foliis ſeptenatis quinatisque, foliolis lanceolatis groſſe dentatis, petalis obcordatis longitudine calycis, caule erecto. W.

Dunkeles Fingerkraut. W

Habitat in Sibiria. ♃ (v. v.)

Similis praecedenti, ſed caulis *et* folia *magis villoſa,* Petala *flava longitudine calycis.* W.

*16. POTENTILLA *piloſa.* W.

P. foliis quinatis, foliolis obovatis groſſe dentatis, petalis ſubretuſis calyce minoribus, caule erecto. W.

Potentilla hirta. *Allion. pedem. n.* 1478. *t.* 71. *quoad figuram.*

Behaartes Fingerkraut. W.

Habitat in Europa *auſtrali.* ♃ (v. v.)

Differt a binis praecedentibus quibus habitu ſimilis: foliis *quinatis nec ſeptenatis. foliolis obovatis, praecipue in foliis radicalibus,* petalis *obtuſis leviſſime retuſis, nec obcordatis, calyce brevioribus. Ceterum* Corolla *flava et* caulis, folia, calyces*que pilis ſparſis tecti.* W.

17. POTENTILLA *argentea.*

P. foliis quinatis cuneiformibus inciſis ſubtus tomentoſis, caule erecto. *Mill. dict. n.* 3. *Gmel. ſib.* 3. *p.* 184. *n.* 37. *Scop. carn. n.*619. *Pollich. pal. n.*496. *Kniph. cent.* 11. *n.* 86. *Roth. germ. I.* 222. *II.* 572. *Hoffm. germ.* 179

P. foliis digitatis, caule erecto corymboſo. *Fl. ſuec.* 417, 454.

P. foliis digitatis inciſo-ſerratis, caule erecto. *Hort. cliff.* 193.

Fragaria caule erecto, foliis quinatis, apice ſerratis, ſubtus tomentoſis. *Hall. helv. n.* 1124.

Fragaria foliis quinatis cuneiformibus et cuneiformiter inciſis, ſubtus tomentoſis: caule firmo erecto. *Crantz. auſtr. p.* 73.

Quinquefolium petraeum majus. *Tabern.* 122. R.

Quinquefolium folio argenteo. *Bauh. pin.* 325.

Pentaphyllum minus. *Cam. epit.* 760.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 155.

Silberblättriges Fingerkraut. W.

Habitat in Europae *ruderatis.* ♃

*18. POTENTILLA *geranioides.* W.

P. foliis quinatis, foliolis pinnatifidis, laciniis linearibus, caule decumbente. W.

Quinquefolium orientale humifuſum, tenuiſſime laciniatum. *Tournef. cor.* 21.

Storchſchnabelblättriges Fingerkraut. W.

Habitat in Armenia. ♃ (v. ſ.)

Radix *craſſitie pennae anſerinae multiceps.* Caules *decumbentes tri-ſ. quadriunciales.* Folia *fere Geranii diſſecti vel columbini, radicalia quinata viridia villoſa, praeſertim ſubtus.* Foliola *profunde pinnatifida, laciniis linearibus.* Folia *caulina ternata, foliolis ut in folio radicali. Affinis P. multifidae, ſed* foliis *quinatis ſubtus non tomentoſis diverſa.* W.

19. POTENTILLA *intermedia.*

P. foliis radicalibus quinatis, caulinis ternatis, caule erectiuſculo ramoſiſſimo. *Mant.* 76.

Fragaria caule ſuberecto, foliis quinis et ſeptenis ſerratis obtuſis. *Hall. helv. n.* 1120. (*excluſis ſynonymis.*)

Quinquefolium montanum luteum erectum hirſutum. *Bauh. pin.* 325. *prodr.* 139. *Raj. eur.* 213.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 156.
Mittleres Fingerkraut. W.
Habitat in Helvetia, Delphinatu, Hungaria. ♃ (v. ſ.)

Media inter P. argenteam *et* norvegicam; *verbo, eſſet* P. norvegica, *ſi non Radix eſſet perennis, et folia ſubradicalia quinata. Ceterum folia inferiora alterna, ſuperiora oppoſita.* Foliola *obovata, inciſo-ſerrata, non tomentoſa, ſerraturis novem.* F. caulina *ternata, paniculae oppoſita.* H. U.

20. POTENTILLA *hirta.*

P. foliis ſeptenatis quinatisque cuneiformibus inciſis piloſis, caule erecto hirto. *Sp. pl.* 712. *Mattuſch. fil.* 1. *n.* 364. *Kniph. cent.* 4. *n.* 65. *Hoffm. germ.* 179.
Quinquefolium rectum minus, anguſtioribus foliis, monſpelienſe. *Raj. eur.* 213.
Quinquefolium montanum erectum hirſutum luteum. *Magnol. monſp.* 216. (*non C. Bauh.*)
Houttuyn Lin. Pfl. Syſt. 7. *p.* 156.
Haariges Fingerkraut. W.
Habitat Monſpelii; *inque* Pyrenaeis *et* Sileſia. ♃ (v. v.)

Folia *radicalia* 5. *ſ.* 7; *Caulina, omnia alterna, ſublinearia, dentibus* 3. *ſ.* 5. Caulis, Pedunculi *et* Calyces *valde piloſi.*

Foliola *inciſo-dentata, in ſpontanea planta duplo anguſtiora, quam in culta.* W.

*21. POTENTILLA *inclinata.*

P. foliis inferioribus quinatis lanceolatis dentatis ſubtus caneſcentibus, caulinis ternatis, caule adſcendente, petalis calycis longitudiue. W.
P. (*inclinata*) tota hirſuta, foliis quinis ſenisve, caulinis ternatis oblongis, caule aſſurgente ſubdichotomo. *Villars. delph.* 3. *p.* 567. *t.* 45. *
Aufwärtsſteigendes Fingerkraut. W.
Habitat in Delphinatu. ♃ (v. ſ.)

Accedit ad P. hirtam et ad P. opacam et tenet medium inter has. A P. hirta diverſa: foliolis *non profunde inciſis ſubtus caneſcentibus nec pilis longis obſitis,* caule *adſcendente,* ramis *magis divaricatis. A P. opaca vero:* foliis *minoribus rarius ſeptenis et ſubtus caneſcentibus,* caule *dichotomo. Cum P. verna nullam habet ſimilitudinem. Figura Clariſſ. Villarſii non optima.* W.

22. POTENTILLA *ſtipularis.*

P. foliis ſeptenatis ſeſſilibus ſtipulae dilatatae inſidentibus. *Sp. pl.* 712. *Gmel. ſib.* 3. *p.* 185. *t.* 37. *f.* 2.

P. foliis novenis palmatis apice ſerratis. *Hall. goett.* 2. *p.* 196.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 157.

Scheidenblättriges Fingerkraut. W.

Habitat in Sibiria. ♃ (v. ſ.)

Caulis *adſcendens, filiformis, laevis, apice diviſus.* Stipulae *amplexicaules, ſolitariae, emarginatae.* Foliola 8 *ad* 9. *lineari-lanceolata, ſubſerrata, glabra, ex emarginatura ſtipulae pendentia absque petiolo.*

Folia *caulina, excepto infimo, quod breve petiolatum, in ſtipulis ſeſſilia.* Foliola *lineari-cuneiformina apice tri-ſ. quinquedentata.* Petala *flava obcordata.* W.

23. POTENTILLA *opaca.*

P. foliis radicalibus ſeptenatis quinatisque lineari-cuneiformibus dentatis, petalis retuſis calycis longitudine, caulibus filiformibus decumbentibus hirtis. W.

P. foliis radicalibus quinatis cuneiformibus ſerratis, caulinis ſuboppoſitis, ramis filiformibus decumbentibus. *Amoen. acad.* 4. *p.* 274. * *Hudſ angl.* 196. *Pollich. pal. n.* 498. *Iacq. ic. rar.* 1. *t.* 91. *Collect.* 1. *p.* 33. * *Hoffm. germ.* 179. *Roth. germ. I.* 223. *II.* 573.

Fragaria (*rubens*) foliis quinatis anguſtis cuneiformibus et cuneiformiter ſerratis, piloſis, caulibus filiformibus rubentibus. *Crantz. auſtr. p.* 75. *n.* 7.

Quinquefolium minus repens lanuginoſum luteum. *Bauh. pin.* 325.

Quinquifolium IV, flavo flore 2. ſpecies. *Cluſ. hiſt.* 2. *p.* 106. *f.* 106.

Quinquefolio ſimilis enneaphyllos hirſuta. *Bauh. pin.* 325. *prodr.* 139.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 157.

Schattenliebendes Fingerkraut. W.

Habitat in Auſtria, Helvetia, Baldo, Germania. ♃ (v. v.)

Similis ſequenti, tamen diverſa, caulibus *decumbentibus longioribus rubris pilis patentiſſimis hirtis,* foliis *anguſtioribus profunde dentatis non pubeſcentibus, ſed hirtis,* pedunculis *longioribus,* corollis *minoribus tantum verſus meridiem apertis alio tempore clauſis.* W.

24. POTENTILLA *verna.*

P. foliis quinatis obovatis dentatis pubeſcentibus, petalis obcordatis calyce majoribus, caulibus declinatis. W.

P. foliis radicalibus quinatis acute ſerratis retuſis, caulinis ternatis, caule declinato. *Fl. ſuec.* 419, 455. *Pallas it.* 1. *p.* 147. *Scop. carn. n.* 621. *Allion. pedem n.* 1481. *t.* 24. *f.* 2. *Kniph. cent.* 8. *n.* 78. *Pollich. pal. n.* 497. *Hoffm. germ.* 179. *Roth. germ. I.* 233. *II.* 573.

Fragaria caule decumbente, foliis pinnatis ſerratis ſubhirſutis. *Hall. helv. n.* 119.

Potentilla foliis quinatis inciſis, caule aſſurgente. *Fl. lapp.* 212. *Hort. cliff.* 194. *Roy. lugdb.* 276.

Fragaria foliis ternatis quinatisque ſubtus canis, caulinis ternatis. *Crantz. auſtr. p.* 74. *n.* 6. *t.* 1. *f.* 1.

Quinquefolium minus repens luteum. *Bauh. pin.* 325.

Pentaphyllum ſ. Quinquefolium minus. *Tabern. ic* 123.

β. P. (*filiformis*) caule proſtrato, foliis quinatis et ſeptenis oblongis ſerratis, caulinis ternatis ſtipulatis, pedunculis ſubradicalibus elongatis. *Vill. delph.* 3. *p.* 564.

P. (*ſalisburgenſis*) foliis hirſutis inciſo-ſerratis obtuſis, radicalibus quinatis, caulinis ternatis caule adſcendente compreſſo paucifloro, calycibus ſubdentatis. *Iacq. ic. rar.* 3. *t.* 490. *Collect.* 2. *p.* 68. *

γ. P. (*rubens*) foliis quinatis et ſeptenatis, caule ſuberecto, petalis cordatis aureis maculatis. *Vill. delph.* 3. *p.* 566.

Quinquefolium minus repens aureum. *Bauh. pin.* 325.

δ. Potentilla ſubacaulis. *Iacq. ic. rar.* 3. *t.* 491. *Collect.* 2. *p.* 145. *

Houttuyn Lin. Pfl. Syſt. 7. *p.* 158.

Frühlings Fingerkraut. W.

Habitat in Europae *et* Aſiae *borealis paſcuis ſiccis, frigidioribus.* ♃ (v. v.)

Variat macula *ad baſin petalorum fulva et nulla tali.*

Planta *ratione loci natalis admodum varians. Nec limites invenio, licet numeroſa ſpecimina comparaverim, quibus apte diſtingantur omnes varietates; una enim ſenſim in alteram tranſiit. P. ſalisburgenſis ab Haenke in Iacquini collectariis deſcripta, non niſi magnitudine, a P. filiformi eſt diverſa, quae iterum cum aliis varietatibus convenit. P. ſubacaulis ejusdem*

Botanici a ſpeciminibus ſibiricis eſt diverſa, P. ſubacaulis Linnaei foliis tantum gaudet ternatis minutiſſimis, et eſt forte ſola hujus ſpeciei varietas a climate boreali orta. W.

25. POTENTILLA *aurea.*

P. foliis radicalibus quinatis, foliolis cuneiformibus ciliatis apice quinquedentatis, caulinis ternatis ſubſeſſilibus, petalis obcordatis calyce majoribus, caule adſcendente. W.

P. foliis radicalibus quinatis ſerratis acuminatis, caulinis ternatis, caule declinato. *Amoen. acad* 4. *p.* 316.* *Jacq. vind.* 245. *Oed. dan. t.* 114. *Mattuſch. ſil. n.* 366. *Hoffm. germ.* 179.

Fragaria caule reclinato, foliis quinis ſerratis. *Hall. helv. n.* 1122. *t.* 21.

Fragaria foliis radicalibus quinatis ſplendentibus, ora argentea, petalis maculatis. *Crantz. auſtr. p.* 77. *n.* 9.

P. foliis margine ſericeis acute ſerratis, caule declinatio. *Ger. prov.* 468.

Quinquefolium 3, aureo flore. *Cluſ. hiſt.* 2. *p.* 105.

Quinquefolium alpinum ſplendens, aureo flore. *Bauh. hiſt.* 2. *p.* 398. *Hall.*

Quinquefolium minus repens alpinum aureum. *Bauh. pin.* 325.

β. Varietas flore pleno. *Hall. l. c.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 159.

Goldfarbiges Fingerkraut. W.

Habitat in alpibus Helvetiae, Auſtriae, Daniae, Sileſiae. ♃ (v. ſ.)

Planta *a P. verna primo intuitu diſtinguenda*: foliolis *cuneiformibus apice tantum* 3-*ſ.* 4-*dentatis*; pilis *ad foliorum marginem et coſtam mediam arcte adpreſſis nitidis*, caule *adſcendente.* W.

*26. POTENTILLA *aſtracanica.*

P. foliis radicalibus quinatis oblongis dentatis, ſummis tripartitis, corollis calyce majoribus, caule adſcendente. W.

P. foliis radicalibus et caulinis infimis quinatis, caulibus dichotomis villoſis, baſi procumbentibus. *Jacq. ic. rar.* 1. *t.* 92. *Miſc.* 2. *p.* 349 *

Aſtracaniſches Fingerkraut. W.

Habitat ad Aſtracan. ♃ (v. ſ.)

Tota planta pubeſcens. Differt ab affinibus nempe P. recta, obſcura et piloſa, foliis *caulinis ſuperioribus tripartitis apice tantum dentatis, ſummis integerrimis, et* caule *adſcendente.* Petala *ſunt pallide flava. In figura bona Iacquini pictor perperam corollam monopetalam pinxit.* W.

27. POTENTILLA *canadenſis.*

P. foliis quinatis villoſis, caule adſcendente hirſuto. *Sp. pl.* 713.

Quinquefolium canadenſe humilius. *Moriſ. hiſt.* 2. *p.* 188.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 160.

Canadiſches Fingerkraut. W.

Habitat in Canada. *Kalm.*

Habitus *Fragariae.* Folia *radicalia quinata: foliolis obovatis, acute ſerratis, ſupra pubeſcentibus, ſubtus villoſis albo-ſericeis, petiolis valde albo-hirſutis.* Caulis *digitalis, ramoſus, rarioribus foliis.* Corollae *luteae. Similis* P. vernae, *a qua differt Foliis valde piloſis, argute ſerratis, ſubrotundis et caule valde hirſuto.*

28. POTENTILLA *alba.*

P. foliis quinatis apice conniventi-ſerratis, caulibus filiformibus procumbentibus, receptaculis hirſutis. *Hort. cliff.* 194. *Roy. lugdb.* 276. *Iacq. auſtr. t.* 115. *Mattuſchk. ſil. n.* 367. *Kniph. cent.* 8. *n.* 77. *Hoffm. germ.* 179. *Roth. germ. I.* 223. *II.* 574.

Fragaria (*alba*) petiolis et foliis quinatis, caule procumbente fere altioribus. *Crantz. auſtr. p.* 82. *n.* 2.

Quinquefolium album. *Hall. helv. it.* 70.

Quinquefolium album majus alterum. *Bauh. pin.* 325.

Quinquefolium album minus. *Bauh. pin.* 325.

Quinquefolium Tragi. *Trag.* 507.

Pentaphyllum majus, flore albo. *Cluſ. hiſt.* 2. *p.* 105.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 160.

Weiſses Fingerkraut. W.

Habitat in alpibus Stiriae, Auſtriae, Pannoniae; *in ſylvis* Germaniae. ♃ (v. v.)

29. POTENTILLA *caulescens.*

P. foliis quinatis apice conniventi-ſerratis caulibus multifloris decumbentibus, receptaculis hirſutis, petalis obovatis. W.

P. foliis quinatis apice conniventi-ſerratis, caulibus multifloris decumbentibus, receptaculis hirſutis. *Amoen. acad.* 4. *p.* 317.* *Iacq. auſtr. t.* 220.

Fragaria foliis ovatis, acute ſerratis, oris ſericeis, pappo diſci longiſſimo. *Hall. helv. n.* 1123.

Quinquefolium album majus. *Bauh. prodr.* 139. caulefcens. *Bauh. pin.* 325. *Hall. opuſc.* 263. *n.* 50. 47.

Quinquefolium album minus alterum. *Bauh pin.* 325. *Houttuyn Lin. Pfl. Syſt* 7. *p.* 161.

Vielſtieliges Fingerkraut. W.

Habitat in alpinis Helvetiae, Auſtriae, Stiriae, Horto Dei. ♃ (v. ſ.)

Calyces *hirſuti, purpurei.*

Foliola *cuneiformia apice argute ſerrata, dente terminali reliquis breviore.* Corymbus *terminalis multiflorus.* Petala *obovata alba.* W.

*30. POTENTILLA *cluſiana.*

P. foliis quinatis apice conniventi-ſerratis, caulibus multifloris decumbentibus, receptaculis hirſutis, petalis ſubrotundis. *Murr. ſyſt. ed.* 14. *p* 478. *Iacq. auſtr. t.* 116. *Hoſt. ſynop.* 285.

Quinquefolium ſecundum minus albo flore. *Cluſ. hiſt.* 2. *p.* 105.

Cluſiſches Fingerkraut. W.

Habitat in Auſtriae *alpibus.* ♃ (v. ſ.)

Simillma praecedenti, tamen diſtincta: Foliis *apice* 3-*ſ.* 5-*dentatis, dentibus aequalibus*, corymbo 5-*ſ.* 10-*floro, nec* 20-*floro*, petalis *ſubrotundis, denique* ſtatura *minori.* W.

*31. POTENTILLA *lupinoides.* W.

P. foliis quinatis utrinque ſericeo-villoſis, foliolis obovatis apice obtuſe dentatis, petalis calycis longitudine, receptaculis lanatis. W.

Potentilla valderia. *Villars delph.* 3. *p.* 572.* (*excluſis ſynonymis.*)

Feigbohnenblättriges Fingerkraut. W.

Habitat in Delphinatus *alpibus.* ♃ (v. ſ.)

Toto coelo a P. valderia diſtincta, ut ex ſpecimine mecum a Claſſ. Villars communicato vidi. Differt a P. valderia: foliis *utrinque ſericeo villoſis, nec ſubtus tomentoſis*, foliolis *apice tantum dentatis*, ſtatura *duplo minori. A P. caulefcenti diverſa*, foliis *utrinque ſericeo-villoſis*, foliolis *obovatis*, petalis *longitudine ca-*

lycis, ſtatura *minorii*, *a P. nitida*: foliis *quinatis*, *pilis non tam denſis ac nitidis*, caule *multifloro altiore et* petalorum *magnitudine*. W.

32. POTENTILLA *nitida*.

P. foliis ſubternatis tomentoſis conniventi-tridentatis, caulibus unifloris, receptaculis lanatis. *Syſt. veg.* 398.

P. foliis quinatis tomentoſis conniventi-tridentatis, caulibus unifloris, receptaculis lanatis. *Amoen. acad.* 4. *p.* 316. * *Sp. pl.* 2. *p.* 714. *Iacq. auſtr.* 5. *t.* 25.

Heptaphyllum argenteum alpinum trifoliatum ſabaudum. *Bocc. muſ.* 2. *p.* 20. *t.* 9.

Trifolium alpinum argenteum perſici flore. *Bauh. pin.* 328. *Pon. bald.* 222. *Raj. hiſt.* 615.

Comaroides alpina argentea perſici flore. *Segv. ver.* 1. *p.* 497.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 162.

Glänzendes Fingerkraut. W.

Habitat in Baldo, Auſtria. ♃ (v. ſ.)

Folia *ternata eſſe monet D. Turre; habeo unum alterumve exemplar ſiccum foliis infimis tantum quinatis.* Cultura *determinabit folia radicalia.*

33. POTENTILLA *valderia*.

P. foliis ſeptenatis obovatis ſerratis tomentoſis, caule erecto, petalis calyce brevioribus, receptaculis lanatis *Sp. pl.* 714. *Allion. pedem n.* 1454. *t.* 24. *f.* 1.

Quinquefolium pyrenaicum, folio ſubtus argenteo. *Tournef. inſt.* 297.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 262.

Valdenſiſches Fingerkraut. W.

Habitat in alpibus Valderianis *et* Vinadenſibus. ♃ (v. ſ.)

Folia radicalia *ſeptena: foliolis obovatis, conniventi-ſerratis, ſubtus valde tomentoſis.* Caulis *erectus paucis foliis ternatis.* Flores *dichotomi, corymboſi, conferti.* Petala *alba, calyce breviora.*

34. POTENTILLA *reptans*.

P. foliis quinatis, caule repente, pedunculis unifloris. *Mat. med.* 131. *Neck. gallob. p.* 222. *Pall. it.* 1. *p.* 205. *Pollich. pal. n.* 499. *Scop. carn. n.* 622. *Blackw. t.* 454. *Ludw. ect. t.* 116. *Kniph. cent.* 7. *n.* 73. *Hoffm. germ.* 180. *Roth. germ. I.* 223. *II.* 575.

Fragaria foliis quinatis ſerratis, petiolis unifloris, caule reptante. *Hall. helv. n.* 118.

Fragaria Pentaphyllum. *Crantz. auſtr. p.* 80.
Potentilla foliis digitatis longitudinaliter patenti-ſerratis, caule repente. *Hort. cliff.* 194. *Gron. virg.* 163.
Quinquefolium majus repens. *Bauh. pin.* 325.
Quinquefolium majus luteum. *Fuchſ. hiſt.* 624.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 163.
Gemeines Fingerkraut. W.
Habitat in Europae *apricis argilloſis.* ♃ (v. v.)

* * * *Foliis ternatis.*

35. POTENTILLA *monſpelienſis.*

P. foliis ternatis, caule ramoſo erecto, pedunculis ſupra genicula enatis. *Hort. upſ.* 134. * *Mill. dict. n.* 7.
Pentaphylloides majus erectum, flore luteo. *Moriſ hiſt.* 2. *p.* 193. *ſ.* 2. *t.* 20. *f.* 2.
Pentaphylloides fragariae folio. *Magn. monſp.* 304.
Fragaria ſterilis alpina cauleſcens. *Boerh. lugdb.* 1. *p.* 42.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 164.
Franzöſiſches Fingerkraut. W.
Habiat Monſpelii. ☉ (v. v.)
Foliola *ovata, obtuſe et aequaliter ſerrata.*

36. POTENTILLA *norvegica.*

P. foliis ternatis, caule dichotomo, pedunculis axillaribus. *Sp. pl.* 715. *Fl. dan.* 171. *Hoffm. germ.* 180. *Retz. obſ.* 1. *p.* 20.
Fragaria foliis ternatis hirſutis, caulinis tripartitis, ſtipulis ovato lanceolatis. *Hall. helv. n.* 1115?
P. foliis ternatis, caule corymboſo erecto. *Fl. ſuec.* 420, 457. *
P. foliis ternatis inciſis, caule diffuſo. *Fl. lapp.* 211. *Hort. cliff.* 193. *Roy. lugdb.* 475.
Quinquefolium hirſutum luteum paucioribus laciniis. *Loeſel. pruſſ.* 218. *t.* 70.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 164.
Norvegiſches Fingerkraut. W.
Habitat in Norvegiae, Sueciae, Boruſſiae, Canadae, Sibiriae *agris.* ☉ (v. v.)
Foliola *lanceolata, ſerrata, dentata.*

37. POTENTILLA *nivea.*

P. foliis ternatis inciſis ſubtus tomentoſis, caule adſcendente. *Fl. ſuec.* 2. *n.* 458. * *Gunn. norv. n.* 512. *t.* 3. *f.* 1.

P. foliis ternatis utrinque hirſutis, infra candidis, foliolis ovatis ſerratis. *Gmel. ſib.* 3. *p.* 183. *t.* 36. *f.* 1.
P. foliis ampliter crenatis ſubtus tomentoſis. *Hall. goett.* 2. *p.* 108.
Fragaria ſterilis, folio ſubtus argenteo. *Amman. ruth. n.* 111.
β. Potentilla leucophylla. *Pall. itin.* 3. *p.* 194.
Fragaria ſterilis procumbens, foliis betonicae inſtar ſerratis. *Amm. ruth. n.* 109. *t.* 14. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 165.
Schneeweiſses Fingerkraut. W.
Habitat in alpibus Lapponiae, Sibiriae. ♃ (v. ſ.)
Structura *P. vernae, ſeu folia P. argenteae, at terna et latiora, magisque hirſuta, ſubtus tomentoſa nivea.* Corolla *lutea.*

*38. POTENTILLA *ſpecioſa.* W.
P. foliis ternatis ellipticis obtuſis dentatis ſubtus tomentoſis, caule fruticoſo. W.
Fragaria cretica fruticoſa, folio ſubtus argenteo. *Tournef. cor.* 21.
Prächtiges Fingerkraut. W.
Habitat in Creta. ♄ (v. ſ.)
Caulis *fruticoſus humilis, ramis craſſitie pennae cygneae brunneis, cortice rimoſo ſua ſponte decidente tectis.* Folia *magnitudine et ſigura Fragariae veſcae ſuperne pubeſcentia, ſubtus niveo-tomentoſa.* Petioli *albo-tomentoſi.* Stipulae *lanceolato-acuminatae ſcarioſae.* Corymbi *longe pedunculati, foliis binis alternis remotis ſimplicibus inſtructi.* Pedunculi *valde tomentoſi.* Calyx *tomentoſus magnitudine corollae albae.* W.

*39. POTENTILLA *tridentata.*
P. foliis ternatis cuneiformibus apice trifidis. *Ait. Kew.* 2. *p.* 216 *t.* 9. *Vahl. ſymb.* 2. *p.* 59.*
P. (*retuſa*) foliis ternatis hirtis apice retuſo tridentato. *Retz. prodr. fl. ſcand. ed.* 2. *n.* 642. *Fl. dan.* 799.
Dreyzähniges Fingerkraut. W.
Habitat in Groenlandia, Terra Nova. ♃ (v. ſ.)
Corolla *flava.* W.

40. POTENTILLA *grandiflora.*
P. foliis ternatis dentatis utrinque ſubpiloſis, caule decumbente foliis longiore. *Sp. pl.* 715. *Mill. dict. n.* 8.

Fragaria foliis ternatis, ſericeis, caule declinato, petalis calycem ſuperantibus. *Hall. helv. n.* 1114. *t.* 21.

P. foliis ternatis dentatis utrinque ſubvilloſis et viridibus ovatis ſerratis obtuſis. *Gmel. ſib.* 3. *p.* 183. *n.* 32. *t.* 35 *f.* 1. 2.

P. foliis ternatis dentatis, caule ſuperne dichotomo, petalis calyce duplo longioribus. *Ger. prov.* 409.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 166.

Groſsblütiges Fingerkraut. W.

Habitat in Helvetia, Sibiria, Pyrenaeis. ♃ (v. ſ.)

Folia *utraque pagina concoloria, nec inferiori argentea.* Caules *pedales, adſcendentes: rufeſcentes.* Corolla *magna flava. Planta ſibirica occurrit quandoque foliis nonnullis quinatis.* W.

41. POTENTILLA *ſubacaulis.*

P. foliis ternatis dentatis utrinque tomentoſis ſcapo decumbente. *Sp. pl.* 715. *Gmel. ſib.* 3. *p.* 183. *n.* 34. *t.* 36. *f.* 2.

P. (*frigida*) foliis ternatis ſerratis hirſutis, petalis ovatis vix calyce longioribus. *Vill. delph.* 3. *p.* 563.

P. foliis radicalibus ternatis dentatis utrinque tomentoſis, caulibus ſubfolioſis. *Ger. prov.* 469. *Garid. aix.* 190. *t.* 37.

Fragaria ſterilis molli lanugine pubeſcens. *Amm. ruth.* 84. *n.* 110.

Fragaria ſterilis ſylveſtris ſericea ſ. incana. *Moriſ. hiſt.* 2. *p.* 187.

Fragariae affinis ſericea incana. *Bauh. pin.* 327. *prodr.* 139.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 166.

Kurzſtieliges Fingerkraut. W.

Habitat in Sibiria, Gallia *auſtrali*, Granada, Galloprovinciae *rupe victoriae.* ♃ (v. ſ.)

Folia *ternata, rugoſa, denſe tomentoſa, piloſa, coriacea; cuneiformia, obtuſa, dentata.* Stipulae [illegible] *piloſae.* Caulis *ſ. Scapus decumbens, foliis* [illegible] *gior, biflorus ſ. uniflorus.* Corolla *maxima.* [illegible] *petalis emarginatis, baſi flavis.* Stamina [illegible]

Omnia ſynonyma, excepto Villarſii [illegible] *manni, videntur ad varietates P. vernae* [illegible] *ſibirica planta, forte nil niſi climate* [illegible] *eſt?* W.

1001. TORMENTILLA.

Gen. plant. ed. Reich. n. 691.

Cal. 8-fidus. *Petala* 4. *Sem.* ſubrotunda, nuda, receptaculo parvo exſucco affixa.

1. TORMENTILLA *erecta.*

T. caule erectiuſculo, foliis ſeſſilibus. *Sp. pl.* 716. *Pall. it.* 2. *p.* 655. *Pollich. pal. n.* 500. *Blackw. t.* 445. *Ludw. ect. t.* 174. *Fl. dan. t.* 589. *Kniph. cent. n.* 93. *Knorr. del.* 2. *t. T.* 1. *Hoffm. germ.* 180. *Roth. germ. I.* 220. *II.* 566,

Tormentilla. *Cam. epit.* 685. *Fl. lapp.* 213. *Fl. ſuec.* 421, 459. *Mat. med.* 132. *Hort. cliff.* 194. *Roy. lugdb.* 267.

Fragaria tetrapetala, foliis caulinis ſeſſilibus, quinatis. *Hall. helv. n.* 1117.

Fragaria tormentilla officinarum. *Crantz. auſtr. p.* 80.

Potentilla Tormentilla erecta. *Scop. carn. n.* 620.

Potentilla (*ſylveſtris*) caule erectiuſculo, foliis ternis ſeſſilibus: corollis tetrapetalis. *Neck. gallob. p.* 222.

Tormentilla ſylveſtris. *Bauh pin.* 326.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 168.

Aufrechte Tormentill. W.

Habitat in Europae *paſcuis ſiccis.* ♃ (v. v.)

2. TORMENTILLA *reptans.*

T. caule repente, foliis petiolatis. *Sp. pl.* 716. *Hudſ. angl.* 197. *Hoffm. germ.* 180. *Roth. germ. I.* 220. *II.* 567.

Pentaphyllum minus viride, flore aureo tetrapetalo, radiculas in terram e geniculis dimittens. *Moriſ. hiſt.* 2. *p.* 190.

Pentaphyllum reptans alatum, foliis profundius ſerratis. *Plot. oxf.* 6. §. 7. *t.* 9. *f.* 5.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 169.

Liegende Tormentill.

Habitat in Angliae, Germaniae *nemoribus umbroſis.* ♃ (v. v.)

Tormentillae non differunt a Potentillis niſi numero partium conſtanti. W.

1002. GEUM.

Gen. plant. edit. Schreb. n. 867.

Cal. 10-fidus Petala 5. *Sem.* ariſta geniculata.

1. GEUM *virginianum.*

G. floribus erectis, ariſtis uncinatis nudis, foliis caulinis ternatis, ſuperioribus lanceolatis petalis calyce brevioribus. *Ait. Kew.* 2. *p.* 217.

G. floribus erectis, fructu globoſo: ariſtis uncinatis nudis, foliis ternatis. *Sp. pl.* 716. *Hort. cliff.* 195. *Gron. virg.* 56. *Mill. dict. n.* 6. *Willich. illuſt.* 175.

G. foliis inferioribus quinatis, pinnis rhomboidalibus rariter inciſis, ſuperioribus ternatis, ſtipulis oblongis, amplis, laciniatis, confluentibus, ſeminibus piloſis ariſta nuda. *Murray. comm. nov. Goett. Tom.* 5. *p.* 32. *t.* 32.

Geum canadenſe. *Iacq. hort.* 2. *p.* 82. *t.* 175.

Geum carolinianum. *Walt. carol.* 150. *

Caryophyllata virginiana, albo flore minore, radice inodora. *Herm. par.* 111. *t.* 111.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 170.

Weiſses Geum. W.

Habitat in Virginia, Carolina. ♃ (v. v.)

Differt a G. urbano ſimillimo: Petalis albis calyce brevioribus; Floribus floreſcentibus nutantibus; Pedunculis verſus florem incraſſatis; Seminibus piloſis; Foliis radicalibus ſubbipinnatis.

*2. GEUM *ſtrictum.*

G. floribus erectis, ariſtis uncinatis nudis, foliis caulinis pinnatis, foliolis ſtipulisque inciſo-fiſſis, petalis calyce longioribus. *Ait. Kew.* 2. *p.* 217.

G. (*canadenſe*) floribus erectis, ſeminibus hirſutis, ariſtis nudis, foliis omnibus pinnatis, pinnis cuneiformibus inciſis. *Murr. Comment. goett.* 5. *p.* 33. *t.* 4. *B.*

G. (*aleppicum*) floribus erectis, fructibus oblongis hirſutis, ariſtis uncinatis nudis, foliis pinnatis cauleque hirſutis. *Iacq. ic. rar.* 1. *t.* 93. *Collect.* 1. *p.* 88. *

Steifes Geum. W.

Habitat in America *boreali.* ♃ (v. v.)

3. GEUM *urbanum.*

G. floribus erectis, ariſtis uncinatis nudis, foliis caulinis ternatis, radicalibus lyrato-pinnatis. *Ait. Kew.* 2. *p.* 218.

G. floribus erectis, fructibus globoſis villoſis: ariſtis uncinatis nudis, foliis lyratis. *Hort. cliff.* 195. *Fl. ſuec.* 423, 460. *Mat. med.* 132. *Roy. lugdb.* 276. *Gmel. ſib.* 3. *p.* 188. *Crantz. auſtr. p.* 69. *Pollich. pal. n.* 501. *Blackw. t.* 253. *Ludw. ect. t.* 36. *Kniph. cent.* 9. *n.* 42. *Flor. dan. t.* 672. *Hoffm. germ.* 181. *Roth. germ. I.* 224. *II.* 576.

G. foliis pinnatis, pinna ultima trilobata, floribus patulis, tubis aduncis. *Hall. helv. n.* 1130.

G. foliis inferioribus quinatis, ſuperioribus ternatis, ſupremis ſimplicibus, ſtipulis ſubrotundis diſtinctis trilobis, ſeminibus hiſpidis, ariſta nuda. *Murray. in nov. comm. Goett.* 5. *p.* 33. *t.* 4. *A.*

Caryophyllata urbana. *Scop. carn. n.* 628.

Caryophyllata vulgaris. *Bauh. pin.* 321.

Caryophyllata. *Dod. pempt.* 137.

β. G. (*intermedium*) foliis interrupte-pinnatis, floribus ſubnutantibus, calycibus patentibus (nec conniventibus, nec reflexis) petalis ſubrotundo-obovatis unguiculatis, germinibus piloſis, ariſta nuda uncinata, ſtylo piloſo, fructu pedicellato. *Ehrh. Beitr.* 6. *p.* 143.

Caryophyllata vulgaris majore flore. *Bauh. pin.* 321. *Mapp. alſ.* 58.

Caryophyllata montana ſimilis vulgari. *Camm. hort.* 35.

Caryophyllata ſylveſtris. *Fuchs. hiſt.* 385.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 171.

Gemeines Geum. W.

Habitat in Europae *umbroſis.* ♃ (v. v.)

Folia *radicalia lyrata in hac, et ſtipulae dentatae.*

*4. GEUM *japonicum.*

G. floribus erectis, fructibus hirſutis, ariſtis nudis, foliis ſubtrilobis. *Thunb. jap.* 220.

Japaniſches Geum. W.

Habitat in Japonia. ♃

Caulis *teres flexuoſus erectus hirſutus pedalis et ultra, apice parum et rarius ramoſus.* Rami *alterni breviſſimi ſubtomentoſi.* Folia *alterna petiolata ſubrotunda triloba vel ſubquinqueloba dentata villoſa mollia pollicaria.* Petiolus *tomentoſus folio paullo brevior.* Stipulae *binae foliaceae ovatae inciſo-dentatae petiolo breviores.* Flores *terminales in caule et ramis ſolitarii.* Calyx *decemfidus, laciniis alternis minoribus.* Petala *longitudine calycis.* Semina *ovata pilis griſeis hirta*

ariſtata. Ariſtae *ſubulatae uncinatae reflexae glabrae.* Thunb. l. c.

5. GEUM *rivale.*

G. floribus nutantibus, petalis longitudine calycis, ariſtis plumoſis medio tortis. W.

G. floribus nutantibus, fructu oblongo: ariſtis plumoſis tortis. *Hort. cliff.* 195, *Fl. ſuec* 424, 461. *Roy. lugdb.* 576. *Mat. med.* 132. *Fl. lapp.* 216. *Mill. dict. n.* 2. *Fl. dan. t.* 722. *Kniph. cent.* 1. *n.* 36. *Hoffm. germ.* 181. *Roth. germ. I.* 224. *II.* 576.

G. foliis pinnatis, pinnis acutis, extrema maxima trilobata, floribus conniventibus: nutantibus. *Hall. helv. n.* 1129.

Geum nutans. *Crantz. auſtr. p.* 70.

Caryophyllata rivalis. *Scop. carn. n.* 629.

Caryophyllata aquatica, nutante flore. *Bauh. pin.* 321.

Caryophyllata ſeptentrionalium. *Lob. ic.* 694.

β. G. (*hybridum*) flore nutante, calyce folioſo, corollam polypetalam excedente. *Iacq. ic. rar.* 1. *t.* 94. *Miſc.* 2. *p.* 33. *

Anemone dodecaphylla. *Krock. ſileſ.* 2. *p.* 235. *t.* 20.

Caryophyllata montana flore pleno prolifero, folio hederae terreſtris. *Breyn. cent. p.* 130. *t.* 60.

Caryophyllata prolifera floſculis elegantiſſimis. *Loeſel. pruſſ.* 37. *t.* 6.

Caryophyllata montana tertia. *Cluſ. hiſt.* 2. *p.* 104.

Caryophyllata aqutica floribus plenis magis nutantibus. *Cam. epit.* 726.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 172.

Waſſer-Geum. W.

Habitat in Europae *pratis ſubhumidis.* ♃ (v. v.)

Seminum ariſtae apex in hac piloſus, in antecedentibus nudus.

*6. GEUM *pyrenaicum.* W.

G. floribus nutantibus, petalis calyce longioribus, ariſtis piloſis baſi tortis. W.

Caryophyllata pyrenaica ampliſſimo folio et rotundiore, nutante flore. *Tournef. inſt.* 295.

Pyrenäiſches Geum. W.

Habitat in Pyrenaeis. ♃ (v. ſ.)

Caulis *erectus ſesquipedalis villoſus.* Folia *radicalia pinnata utrinque pilis tecta, pinna terminalis* 3-4-*uncialis ſubrotunda cordata ampliſſima ſubtriloba.* Fo-

lia *caulina cuneiformia inciſo dentata acuminata.* Stipulae *oblongae profunde dentatae.* Petala *ſubrotunda calyce majora.* Semina *villoſo-lanata.* Ariſta *ſupra baſin torta pilis raris tecta, apice nuda. Affine G. rivali, ſed magnitudine foliorum radicalium, et ariſtae fabrica diverſum. An varietas?* W.

*7. GEUM *atlanticum.*

G. villoſum, foliis inferioribus pinnatis, pinnula terminali cordata maxima, caule ſubunifloro, fructibus hirſutis, ariſtis contortis. *Desfont. fl. atl.* 1. *p.* 402.

Atlantiſches Geum. W.

Habitat in monte Atlas *prope* Tlemſen. ♃

Planta *tota villoſa.* Caulis *pedalis ſeu ſesquipedalis et ultra ſimplex uniflorus.* Folia *radicalia petiolata pinnata, pinnulis* 3-5 *quandoque* 7, *remotis ſubrotundis aut ovatis crenato-dentatis, terminalis cordata obtuſa maxima, caulina inferiora obovata, ſuperiora inaequaliter lobata aut laciniata, laciniis acutis.* Stipulae *magnae ovatae inciſae.* Flos *G. montani.* Petala *lutea.* Semina *villoſa.* Ariſta *contorta nunc glabra nunc ſubvilloſa. Affine G. montano, differt* foliolis *paucioribus remotioribus,* caule *altiore,* ariſtis *contortis nec barbatis.* Desfont. l. c.

*8 GEUM *potentilloides.*

G. caule ſubbifloro, ariſtis rectis nudis, calycibus fructus erectis, foliis pinnatis dentatis. *Ait. Kew.* 2. *p.* 219.

Geum Laxmanni. *Gaert. ſem.* 1. *p.* 352. *t.* 74.

Caryophyllata (*potentilloides*) foliis interrupte pinnatis hirſutis ceſpitoſis, ſcapis declinatis ſubtrifloris. *Lamarck. encycl.* 1. *p.* 395.*

Dryas (*geoides*) pentapetala, foliis pinnatis, pinnis extimis confluentibus. *Murr. ſyſt. ed.* 14 *p.* 481. *Pall. it.* 3. *app. n.* 92. *t. Y. f.* 1.* *Iacq. hort.* 3. *p.* 38. *t.* 68. *Falk. ruſſ. t.* 11.

Fingerkrautartiges Geum. W.

Habitat in Sibiriae *rupeſtribus a montibus* Altaicis *usque ultra* Jeniſeam. ♃ (v. v.)

9. GEUM *montanum.*

G. caule unifloro, ariſtis rectis villoſis, foliis pinnatis, piloſis, foliolo extimo maximo ſubrotundo, inferioribus ſenſim minoribus. *Ait. Kew.* 2. *p.* 218.

G. flore inclinato ſolitario, fructu oblongo, ariſtis villo-

ſis rectis. *Sp. pl.* 717. *Mill. dict. n.* 4. *Crantz. auſtr. p.* 70. *n.* 2. *Iacq. auſtr.* 4. *p.* 58. *t.* 373. *Mattuſch. ſil. n.* 372.

G. foliis pinnatis, extrema maxima, rotunde cordata. *Hall. helv n.* 1131.

Caryophyllata montana. *Scop. carn. n.* 630.

Caryophyllata alpina lutea. *Bauh. pin.* 322.

Caryophyllata alpina. *Pon. bald.* 342.

Caryophyllata montana. *Cam. epit.* 727.

β. Caryophyllata alpina minor. *Bauh. pin.* 322. *prodr.* 139.

Caryophyllata alpina minima, flore aureo. *Barr. rar.* 588. *t.* 399.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 173.

Berg-Geum. W.

Habitat in alpibus Helvetiae, Auſtriae, Sileſiae, Delphinatus. ♃ (v. ſ.)

Folia hirſuta. Flos patens. Caulis uniflorus. Petala lutea, emarginata.

Flos *erectus. Crantz. Scop. l. c.*

10. GEUM *reptans.*

G. caulibus unifloris, ariſtis rectis villoſis, foliis pinnatis inciſis piloſis, flagellis reptantibus. *Ait. Kew.* 2. *p.* 219.

G. foliis uniformibus inciſis: alternis minoribus, flagellis reptantibus. *Sp. pl.* 717.

G. foliis pinnatis, trilobatis, inciſis, flagellis folioſis. *Hall. helv. n.* 1132.

Caryophyllata alpina, apii folio. *Bauh. pin.* 322.

Caryophyllata alpina tenuifolia incana, flore luteo longius radicata. *Barr. rar.* 589. *t.* 400. *Bocc. muſ.* 160. *t.* 128.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 174.

Kriechendes Geum. W.

Habitat in Helvetia *et valle* Barſilionenſi. ♃ (v. ſ.)

Caules *uniflori, vix foliis longiores.*

11. GEUM *anemonoides.* W.

G. caule unifloro, ariſtis rectis villoſis, foliis pinnatis glabris, foliolis cuneiformibus apice dentatis. W.

Dryas anemonoides. *Pall. itin.* 3. *app. n.* 92. *nota* 2. *t. E. e. f.* 4.*

Dryas (*pentapetala*) floribus pentapetalis, foliis pinna

tis. *Sp. pl.* 717. *Amoen. acad.* 2. *p.* 353. * *Gmel. ſib.* 3. *p.* 187. *n.* 43.
Caryophyllata (*kamtſchatica*) foliis radicalibus pinnatis glabris, ſcapo ſubnudo unifloro. *Lamarck. encyclp.* 1. *p.* 395. *
Houttuyn Lin. Pfl. Syſt. 7. *p.* 175.
Anemonenartiges Geum. W.
Habitat in Kamtſchatca. ♃ (v. ſ.)

1003. DRYAS.

Gen. pl. ed. Schreb. n. 868.

Cal. ſimplex 8-fidus. *Petala* 8. *Sem.* caudata, piloſa.

1. DRYAS *octopetala.*
D. floribus octopetalis, foliis ſimplicibus. *Fl. lapp.* 215. *Fl. ſuec.* 426, 462. *Hort. cliff.* 195. *Roy. lugdb.* 279. *Segu. ver.* 512. *Gmel. ſib. p.* 188. *Scop. carn. n.* 631. *Moench. haſſ. n.* 428. *Hoffm. germ.* 181. *Roth. germ. I.* 221. *II.* 568.
D. floribus octopetalis, foliis ovatis, ſubtus albiſſimis. *Hall. helv n.* 1133.
Geum (*chamaedrifolium*) flore erecto octopetalo, fructu oblongo, ariſtis lanigeris. *Crantz. auſtr. p.* 70. *n.* 4.
Chamaedrys 3 ſ. montana. *Cluſ. hiſt.* 2. *p.* 351.
Leucas; Chamaedrys alpina. *Oed. dan. t.* 51.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 175.
Alpen-Dryade. W.
Habitat in alpibus Lapponicis, Helveticis, Auſtriacis, Sabaudicis, Hibernicis, Sibiricis, Germanicis. ♃ (v. ſ.)

Petala 10, 6, *plerumque* 8, *raro* 5. Crantz. l. c.

Caulis *proſtratus ramoſus.* Folia *petiolata oblonga profunde dentata, ſuperne viridia glabra rugoſo-venoſa, ſubtus niveo-tomentoſa.* Petioli *villoſi.* Pedunculi *uniflori nudi adſcendentes.* Corolla *alba calyce major. Dryas differt a Geo calyce ſimplici octophyllo, hinc Dryas pentapetala Linnaei et D. geoides Pallaſii ob calycem duplicem Geis adnumeravi.* W.

1004. COMARUM.

Gen. plant. edit. Schreb. n. 869.

Cal. 10-fidus. *Petala* 5, calyce minora. *Recept.* ſeminum ovatum, ſpongioſum, perſiſtens.

1. COMARUM *paluſtre.*

Comarum. *Fl. lapp.* 214. *Fl. ſuec.* 422, 463. *Hort. cliff.* 195. *Roy. lugdb.* 276. *Pollich. pal. n.* 502. *Pall. it. p.* 72. *Fl. dan. t.* 636. *Kniph. cent.* 9. *n.* 24. *Hoffm. germ.* 182. *t.* 7.

C. foliis pinnatis, petalis calyce minoribus. *Roth. germ. I.* 224. *II.* 578.

Fragaria foliis pinnatis, petalis lingulatis minimis. *Hall. helv. n.* 1128.

Fragaria paluſtris. *Crantz. auſtr. p.* 73. *u.* 4.

Potentilla paluſtris. *Scop. carn. ed.* 2. *p.* 617.

Quinquefolium paluſtre rubrum. *Bauh. pin.* 325.

Pentaphyllum paluſtre. *Cord. hiſt.* 95.

Houttuyn Lin. Pſl. Syſt. 7. *p.* 178.

Sumpf Comarum. W.

Habitat in Europae *et* Aſiae *uliginoſis* ♃ (v. v.)

1005. CALYCANTHUS.

Gen. plant. ed. Schreb. n. 870.

Cal. 1-phyllus, urceolatus, ſquarroſus, foliolis, coloratis. *Cor.* 0. *Styli* plurimi Stigmate glanduloſo. *Sem.* plurima, caudata, intra calycem ſucculentum.

1. CALYCANTHUS *floridus.*

C. laciniis calycinis interioribus lanceolatis. W.

C. petalis interioribus longioribus. *Sp. pl.* 718. *Willd. arb.* 51.* *Du Roi. harbk. ed.* 2. 1. *p.* 190.

*. *oblongatus* foliis oblongis. *Ait. Kew.* 2. *p.* 220.

Buttneria anemones flore. *Du Ham. arb.* 1. *p.* 114. *t.* 45.

Baſteria foliis ovatis oppoſitis, floribus lateralibus, caule fruticoſo ramoſo. *Mlli. dict. et ic. t.* 60.

Beureria petalis coriaceis oblongis, calycis foliolis reflexis. *Ehret. pict. t.* 13.

Frutex corni foliis, floribus inſtar anemones ſtellatae, pe-

talis craſſis rigidis colore ſordide rubente, cortice aromatico. *Cateſb. car.* 1. *p.* 46. *t.* 46.

β. ovatus foliis ſubrotundo-ovatis. *Ait. l. c.*
Houttuyn Lin. Pfl. Syſt. 3. *p.* 681.
Caroliniſche Kelchblume. W.
Habitat in Carolina. ♄ (v. v.)
A Cateſbaeo in Europam introductus.

2. CALYCANTHUS *praecox.*

C. laciniis calycinis interioribus ſubrotundo-ovatis. W.
C. petalis interioribus minutis. *Sp. pl.* 718. *Ait. Kew.* 2. *p.* 220. *t.* 10.
Abai ſ. Robai. *Kaempf. amoen.* 878. *t.* 879.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 682.
Japaniſche Kelchblume. W.
Habitat in Japonia. ♄ (v. ſ.)

Ignota mihi, videtur praecedentis generis, ſed diverſae ſpeciei, quod floreat ante folia, quodque petala interiora lutea minora flori adſcribantur.

Diſtincta a praecedente: foliis *multo majoribus ovato-lanceolatis acuminatis*, praecocitate *floris*, laciniis *calycinis interioribus ſubrotundo-ovatis*, ſtaminibus *quinque*. W.

Claſſis XIII.

POLYANDRIA.

MONOGYNIA.

* *Monopetali.*

1054. SWARTZIA. *Cal.* 4-partitus. *Petalum* planum laterale. *Legumen.*
1006. MARCGRAVIA. *Cal.* 6-phyllus, imbricatus. *Cor.* 1-petala, clauſa. *Bacca* multilocularis, multivalvis.
1007. TERNSTROEMIA. *Cal.* 5-partitus. *Cor.* rotata, limbo campanulato, 5-partito. *Bacca* exſucca bilocularis.

* *Tripetali.*

1008. TRILIX. *Cal.* 3-phyllus. *Bacca* 5-locularis, polyſperma.

† *Sterbeckia lateriflora, Tetracerae nonnullae.*

* *Tetrapetali.*

1023. MAMMEA. *Cal.* 2-phyllus. *Bacca* 1-locularis. *Sem.* calloſa.
1015. PAPAVER. *Cal.* 2-phyllus. *Capſ.* 1-locul. coronata.
1014. CHELIDONIUM. *Cal.* 2-phyllus. *Siliqua.*
1027. SPARRMANNIA. *Cal.* 4-phyllus. *Capſ.* pentagona quinquelocularis, loculis diſpermis.
1010. CAPPARIS. *Cal.* 4-phyllus. *Bacca* pedicellata, corticoſa.
1011. ACTAEA. *Cal.* 4-phyllus. *Bacca* 1-locularis. *Sem.* gemino ordine.
1026. CALOPHYLLUM. *Cal.* 4-phyllus. *Drupa* globoſa. *Nucleus* ſubgloboſus.
1025 GRIAS. *Cal.* 4-fidus. *Drupa* 1-ſperma. *Nucleus* 8 ſulcatus.

† *Vallea ſtipularis, Legnotis elliptica, Cleome Chelidonii, felina, Tetracera nitida.*

* *Pentapetali.*

1044. STERBECKIA. *Capſ.* cylindracea corticoſa. *Semina* imbricata in pulpa nidulantia.
1045. LOASA. *Capſ.* ſemi-infera, 1-locularis, ſemi-trivalvis, polyſperma.
1042. MENTZELIA. *Capſ.* infera, 1-locular. 3-valvis, polyſperma.
1049. VALLEA. *Capſ.* ſupera 4-ſ. 5-angularis unilocularis polyſperma.
1050. BONNETIA. *Capſ.* ſupera 3-locularis 3-valvis polyſperma. *Cal.* 5-partitus.
1036. LEGNOTIS. *Capſ.* ſupera 3-locularis 3-valvis. *Sem.* ſolitaria. *Cal.* 5-fidus.
1046. FREZIERA. *Bacca* exſucca 3-locularis. *Cal.* 5-phyllus.
1034. MARILA *Capſula* 4-locularis 4-valvis polyſperma. *Cal.* 5-phyllus.
1048. CISTUS. *Capſ.* ſubrotunda. *Cal.* 5-phylli foliola 2 minora.
1038. LEMNISCIA *Pericarp.* 5-loculare. *Cal.* 5-dentat. *Nectarium* cyathiforme.
1037. MYRODENDRUM. *Pericarp.* 5-loculare. *Cal.* 5-dentat. *Nectarium* o.
1052. CORCHORUS. *Capſ.* ſub 5-locularis. *Cal.* 5-phyllus, longitudine corollae, deciduus.
1018. SARRACENIA. *Capſ.* 5-locularis. *Stigma* clypeatum. *Cal.* exter. 3-phyllus, ſuper. 5-phyllus.
1028. TILIA. *Capſ.* 5-locul. coriacea, 1-ſperma. *Cal.* deciduus.
1022. AUBLETIA. *Capſ.* echinata 10-locularis polyſperma. *Cal.* coloratus.
1024. OCHNA. *Baccae* 5 in receptaculo carnoſo. *Petala* unguibus elongatis.
1039. ASCIUM. *Bacca* unilocularis polyſperma. *Cal.* 5-phyllus.
1031. GREWIA. *Drupa* 4-loba 4-locularis. *Cal.* 5-phyllus.
1017. MUNTINGIA. *Bacca* 5-locularis, umbilicata. *Cal.* partitus.
1035. ELAEOCARPUS. *Drupa* nuce criſpa. *Petala* lacera.

1033. MICROCOS. *Drupa* nuce triloculari. *Petala* linearia.

† *Delphinium Conſolida*, *Aiacis*, *Aconiti*, *Laetia completa*.

* *Hexapetali.*

1016. ARGEMONE. *Cal.* 3-phyllus. *Capſ.* 1-locularis, ſemivalvis.

1045. LAGERSTROEMIA. *Cal.* 6-fidus. *Stam.* 6, exteriora majora. *Capſ.* 6-locul. polyſperma.

1041. ALANGIUM. *Cal* 6-10-dentatus ſuperus. *Petala* 6-ſ. 10. *Bacca* corticoſa 2-3-ſperma.

1047. THEA. *Cal.* 5-ſ. 6-phyllus. *Petala* 6 ſ. 9. *Capſ.* 3-locularis. *Sem.* ſolitaria.

1040. LECYTHIS. *Cal.* 6-phyllus. *Stam.* nectario lingulato connata. *Capſ.* circumſciſſa.

† *Ternſtroemia meridionalis.*

* *Octopetali.*

1012. SANGUINARIA. *Cal* 2-phyllus. *Capſ.* 2-valvis, polyſperma.

* *Enneapetali.*

1013. PODOPHYLLUM. *Cal.* 3-phyllus. *Bacca* 1-locularis, coronata.

† *Thea.*

* *Decapetali.*

1020. BIXA. *Cal.* 5-dentatus. *Cor.* 5-petala, duplex. *Capſ.* 2-valvis.

† *Alangium decapetalum.*

* *Polypetali.*

1019. NYMPHAEA. *Bacca* multilocul. corticoſa. *Cal.* magnus.

* *Apetali.*

1051. PROCKIA. *Cal.* 3-phyllus. *Peric.* 5-loculare.

1032. MAERUA. *Cal.* 4-fidus tubo nectarifero. *Peric.* pedicellatum.

1009. LUDIA. *Cal.* 4-ſ. 9-partitus. *Pericarp.* uniloculare polyſpermum.

1021. SLOANEA. *Cal.* 5-9 fidus. *Peric.* echinatum 5-6 loculare 3-6 valve. *Semina* arillata.

1030. RYANIA *Cal.* 5-phyllus. *Pericarp.* uniloculare polyſpermum. *Semina* arillata.
1029. LAETIA. *Cal.* 5-phyllus. *Peric.* 1-loculare, 3-valve, polyſpermum.
1053. SEGUIERIA. *Cal.* 5-phyllus. *Peric.* 1-ſpermum, alatum.

† *Crataeva Marmelos, Tetracera ſarmentoſa.*

DIGYNIA.

1057. FOTHERGILLA. *Cal.* integerrimus. *Cor.* nulla. *Capſ.* 2-locularis. *Sem.* bina.
1056. CURATELLA. *Cal.* 5-phyllus. *Cor.* 4-petala. *Capſ.* 2-partita, 2-ſperma.
1055. PAEONIA. *Cal.* 5-phyllus. *Cor.* 5-petala. *Capſ.* polyſperma. *Sem.* colorata.
1058. TRICHOCARPUS. *Cal.* 4-ſ. 5-partitus. *Cor.* 0. *Capſ.* ſetoſa polyſperma.
1059. LACIS. *Cal.* 0. *Cor.* 0. *Capſula* 2-valvis polyſperma.

† *Tetracera laevis.*

TRIGYNIA.

1061. DELPHINIUM. *Cal.* nullus. *Cor.* 5-petala: ſupremo petalo cornuto. *Nectar* 2-fidum, ſeſſile.
1062. ACONITUM. *Cal.* nullus. *Cor.* 5-petala: ſupremo galeato. *Nectar.* 2, pedicellata.
1060. HOMALIUM. *Cal.* 6-7-partitus. *Cor.* 6-7-petala. *Stamina* per tria aggregata.

† *Reſeda Luteola. Corchorus aeſtuans.*

TETRAGYNIA.

1063. WINTERA. *Cal.* integer. *Cor.* 6-petala. *Baccae* 4, 4-ſpermae.
1067. CIMICIFUGA. *Cal.* 4-phyllus. *Cor.* nectariis, 4, urceolatis. *Capſ.* 4. *Sem.* ſquamoſa.
1066. WAHLBOMIA. *Cal.* 4-phyllus. *Cor.* 4-petala. *Peric.* 4, roſtrata.
1064. TETRACERA. *Cal.* 6-phyllus. *Capſ.* 4, monoſpermae. *Stamina* dilatata utrinque antherifera.
1065. CARYOCAR. *Cal.* 5-partitus. *Cor.* 5-petala. *Drupa* nucibus 4.

† *Elaeocarpus Dicera.*

PENTAGYNIA.

1068. AQUILEGIA. *Cal.* nullus. *Cor.* 5-petala. *Nectaria* 5, inferne cornuta.
1069. NIGELLA. *Cal.* nullus. *Cor.* 5-petala. *Nectar.* 8, ſuperne 2-labiata.
1070. REAUMURIA. *Cal.* 5-phyllus. *Cor.* 5-petala: nectariis 10, adnatis, ciliatis. *Capſ.* 5-locularis, polyſperma.

† *Aconita et Delphinia nonnulla.*

POLYGYNIA.

1091. HYDRASTIS. *Cal.* nullus. *Cor.* 3-petala. *Bacca* compoſita acinis 1-ſpermis.
1082. ATRAGENE. *Cal.* nullus. *Cor.* 4-petala major: interior polypetala. *Sem.* plurima, criſtata.
1083. CLEMATIS. *Cal.* nullus. *Cor.* 4-petala. *Sem.* plurima, ariſtata.
1084. THALICTRUM. *Cal.* nullus. *Cor.* 4-5-petala. *Sem.* plurima, ſubmutica, nuda.
1088. ISOPYRUM. *Cal.* nullus. *Cor.* 5-petala, decidua. *Nectaria* 5. *Capſ.* polyſpermae.
1089. HELLEBORUS. *Cal.* nullus. *Cor.* 5-petala, perſiſtens. *Nectaria* plura. *Capſ.* polyſpermae.
1090. CALTHA. *Cal.* nullus. *Cor.* 5-petala. *Capſ.* plurimae. *Nectaria* nulla.
1081. ANEMONE. *Cal.* nullus. *Cor.* 6-petala. *Sem.* plurima.
1076. MICHELIA. *Cal.* truncatus. *Cor.* 8-petala. *Baccae* 4-ſpermae, glomeratae.
1087. TROLLIUS. *Cal.* nullus. *Cor.* 14-petala. *Nectaria* linearia. *Capſ.* polyſpermae.
1079. XYLOPIA. *Cal.* 3-phyllus. *Cor.* 6-petala. *Capſulae* 1-ſ. 2-ſpermae in receptaculo hemiſphaerico.
1080. UNONA. *Cal.* 3-phyllus. *Cor.* 6-petala. *Baccae* moniliformes 2-ſpermae in receptaculo hemiſphaerico.
1077. UVARIA. *Cal.* 3-phyllus. *Cor.* 6-petala. *Baccae* polyſpermae recept. longo affixae.
1078. ANNONA. *Cal.* 3-phyllus. *Cor.* 6-petala. *Bacca* cortice imbricato polyſperma.
1075. LIRIODENDRUM. *Cal.* 3-phyllus. *Cor.* 6-petala. *Samarae* plurimae lanceolatae imbricatae.

1074. MAGNOLIA. *Cal.* 3-phyllus. *Cor.* 9-petala. *Capſ.* glomeratae, 2-valves. *Sem.* pendula.

1075. NELUMBIUM. *Cal.* 4-5-phyllus. *Cor.* polypetala. *Nuces* monoſpermae receptaculo immerſae.

1071. DILLENIA. *Cal.* 5-phyllus. *Cor.* 5-petala, *Capſ.* plures, connatae: *Recept.* carnoſo.

1086. RANUNCULUS. *Cal.* 5-phyllus. *Cor.* 5-petala. *Sem.* plurima. *Petala* ungue nectarifero.

1072. ILLICIUM. *Cal.* 6-phyllus. *Pet.* 27. *Capſulae* 1-ſpermae in orbem.

1085. ADONIS. *Cal.* 5-phyllus. *Cor.* 5-ſ. 10-petala. *Sem.* plurima, angulata, corticata.

† *Nigellae nonnullae.*

MONOGYNIA.

1006. MARCGRAVIA.

Gen. pl. ed. Schreb. n. 871.

Cor. 1-petala, calyptriformis. *Cal.* 6-phyllus, imbricatus. *Bacca* multilocularis, polyſperma.

1. MARCGRAVIA *umbellata.*

M. foliis ovato-oblongis acuminatis venoſis. *Vahl. eclog.* 2. *p.* 39.

M. umbellata. *Sp. pl.* 719. *Iacq. amer.* 156. *t.* 96. * *Swartz. obſ.* 205. *

M. ſcandens, fructu radiatim poſito. *Plum. gen.* 7. *ic.* 173.

M. ſcandens, foliis caulinis ſubrotundis ad margines glandulatis, ramorum integris ovatis alternis diſtiche poſitis, floribus umbellatis terminalibus. *Brown. jam.* 244. *t.* 26.

Phyllitidi ſcandenti affinis major, folio craſſo ſubrotundo. *Sloan. jam.* 15. *hiſt.* 1. *p.* 74. *t.* 28. *f.* 1. *mala.*

Houttuyn Lin. Pſl. Syſt. 4. *p.* 1.

Doldenblütige Marcgravie. W.

Habitat in Americae *calidioris montoſis.* ♄ (v. ſ.)

Folia alterna, petiolata, ovato-lanceolata, integerrima. Umbella *ſimplex pedunculis numeroſis, in orbem poſitis, aequalibus, horizontaliter patentibus.* Flores *adſcendentes, nudi, corolla calyceque caducis.* Diſcus *umbellae conſtans glandulis nectariferis, ſeſſilibus, ſaepius quinis, erectis, forma corollae ringentis galeae ſ. ſupremi petali aconiti, ore ambitum reſpicientibus.* Receptaculum *poſt caſum floris integrum, nec quinquelobum.*

*2. MARCGRAVIA *coriacea.*

M. foliis ellipticis obtuſis coriaceis aveniis. *Vahl. eclog.* 2. *p.* 39. *

Lederartige Marcgravie. W.

Habitat in Cajenna. ♄

Rami *glabri, cortice cinereo, punctis raris adſperſi, inferne teretes, ſuperne ſubtetragoni.* Folia *breviter petiolata* 4-*ſ.* 5-*pollicaria utrinque glabra, laevia, margine parum reflexa. Pedicelli verſus ſummitatem pedunculi communis verticillati* 17 *vel* 18

patentiſſimi bipollicares aequales teretes verſus florem incraſſati, e punctis frequentiſſimis ſparſis tuberculati. Flores *adſcendentes.* Vahl l. c.

1007. TERNSTROEMIA.

Gen. pl. ed. Schreb. n. 872.

Cor. 1-petala rotata, limbo campanulato 5-ſ. 6-partito. *Cal.* quinquepartitus. *Antherae* apice craſſae. *Bacca* exſucca bilocularis.

1. TERNSTROEMIA *meridionalis.*

T. foliis obovatis emarginatis integerrimis, pedunculis axillaribus. *Vahl. ſymb.* 2. *p.* 60.

T. foliis obovatis integerrimis, pedunculis terminalibus. *Swartz. prodr.* 81. *obſ. bot.* 207. *

Ternſtroemia meridionalis. *Lin. ſuppl.* 264.

Mittägige Ternſtrömie. W.

Habitat in Jamaica, Nevis, Dominica, Nova Granada. ♄

Arbor *ramis determinatis, ramulis ſimplicioribus rigidulis, cortice cinereo.* Folia *alterna ſubpetiolata coriacea perennia ovalia obtuſa ſubemarginata, margine ſubrecurvata integerrima avenia.* Stipulae *nullae.* Pedunculi *axillares groſſiores compreſſo-ancipites foliis dimidio breviores cernui.* Calyx *rigidus fere cartilagineus.* Corolla *ſaepius quinquepartita rarius ſexpartita alba, primo globoſa, dein campanulata.* Bacca *ſicca, quia matura caſu absque valvulis rumpitur, atque adeo conſtantem nullam valvularum ſtructuram retinet.* Semina *ſanguinea ſericea.*

*2. TERNSTROEMIA *elliptica.*

T. foliis ellipticis acutis integerrimis pedunculis lateralibus. *Vahl. ſymb.* 2. *p.* 61. * *Swartz. prodr.* 81.

Elliptiſche Ternſtroemie. W.

Habitat in India *occidentali.* ♄ (v. ſ.)

Folia *potius obtuſa quam acuta dicenda.* W.

*3. TERNSTROEMIA *punctata.*

T. foliis oblongis integerrimis ſubemarginatis margine punctatis, pedunculis axillaribus. *Swartz. prodr.* 81.

Taonabo (*punctata*) foliis ovato-oblongis emarginatis, fructu majore. *Aubl. guj.* 1. *p.* 571. *t.* 228. *

Punctirte Ternſtrömie. W.

Habitat in Gujanae *ſylvis montis* Serpent *dicti.* ♄

*4. TERNSTROEMIA *japonica.*
T. foliis ovato - lanceolatis apice ſerrulatis, pedunculis lateralibus.
T. (*japonica*) foliis acuminatis exciſis. *Thunb. Act. ſoc. Lin. Lond.* 2. *p.* 335.
Cleyera japonica. *Thunb. jap.* 224.*
Mokokf. *Kaempf. amoen.* 873. *t.* 774.
Japaniſche Ternſtrömie. W.
Habitat in Japonia. ♄

*5. TERNSTROEMIA *dentata.*
T. foliis oblongis acuminatis dentato - ſerratis, pedunculis axillaribus lateralibusque. *Swartz. prodr.* 81.
Taonabo (*dentata*) foliis ovato - oblongis denticulatis, floribus flaveſcentibus, fructu minore. *Aubl. guj.* 1. *p.* 569. *t.* 227. *
Gezähnte Ternſtrömie. W.
Habitat in Gujanae *ſylvis.* ♄ (v. f.)

1008. TRILIX.

Gen. plant. edit. Schreb. n. 873.

Cor. 3 - petala. *Cal.* 5 - phyllus. *Bacca* 5 - locularis, polyſperma.

1. TRILIX *lutea.*
T. lutea. *Mant.* 247.
Iacquinia. *Mutis mſſ.*
Houttuyn Lin. Pfl. Syſt. 4. *p.* 3.
Gelbe Trilix. W.
Habitat Carthagenae *Americes.* ♄
Frutex *biorgyalis, ramoſiſſimus.* Rami *teretes, ſcabriuſculi.* Folia *alterna, petiolata, ſubpeltata, cordato - ovata, ſerrata, acuminata, venoſa, pubeſcentia.* Petioli *teretes, glabri.* Flores *lutei* (*non a corolla, ſed ab antheris*). Pedunculi *plerumque terminales: Pedicellis alternis, unifloris, teretibus, pubeſcentibus.*

*1009. LUDIA.

Juſſ. gen. pl. ed. Uſteri 379.

Cal. 4 - 7 - partitus. *Cor.* o. *Styl.* 3 - 4 - fidus. *Bacca* ſicca 1 - locularis polyſperma.

*1. LUDIA *heterophylla.*
L. foliis obovatis nitidis venoſis, junioris fruticis mini-

mis dentato-ſpinoſis, fruticis adulti majoribus integerrimis. *Lamarck. encycl.* 3. *p.* 586.*
α. adulta foliis obovatis integerrimis. *Lamarck. illuſt. gen. t.* 466. *f.* 1.
β. junior foliis orbiculatis acuminato-dentatis Malpighiae cocciferae. *Lamarck. illuſt. gen. t.* 466. *f.* 2.
Verſchiedenblättriger Aenderling. W.
Habitat in inſula Mauritii. ♄

*2. LUDIA *myrtifolia.*
L. foliis ovatis integerrimis venoſis, ſtylo incurvo. *Lamarck. encycl.* 3. *p.* 586.* *illuſt. gen. t.* 466. *f.* 3.
Myrtenblättriger Aenderling. W.
Habitat in inſula Borbonia. ♄

*3. LUDIA *ſeſſiliflora.*
L. foliis ovato-oblongis acutiuſculis, floribus ſubſeſſilibus, ſtylo ſuperne trifido. *Lamarck. encycl.* 3. *p.* 586.*
Ludia tuberculata. *Iacq. hort. Schoenb.* 1. *p.* 59. *t.* 112.*
Stiellosblütiger Aenderling. W.
Habitat in inſula Mauritii. ♄
In hac ſpecie eſt calyx 4-*fidus et* bacca *trilocularis, ſeminibus ſolitariis.* W.

1010. CAPPARIS.

Gen. pl. ed. Schreb. n. 876.

Cal. 4-phyllus, coriaceus. *Petala* 4. *Stam.* longa. *Bacca* corticoſa, 1-locularis, pedunculata.

* *Stipulis ſpineſcentibus.*

1. CAPPARIS *ſpinoſa.*
C. pedunculis unifloris ſolitariis, ſtipulis ſpinoſis, foliis ſubrotundis obtuſis glabris, capſulis ovalibus. W.
C. pedunculis unifloris ſolitariis, ſtipulis ſpinoſis, foliis annuis, capſulis ovalibus. *Mat. med.* 133. *Hall. helv. n.* 1077. *Blackw. t.* 417. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 179. *Scop. carn. ed.* 2. *n.* 622. *Ludw. ect. t.* 107. *Desfont. fl. atlant. I. p.* 403.*
C. aculeata. *Hort. cliff.* 203. *Roy. lugdb.* 476. *Gron. orient.* 154.
C. retuſo folio. *Lobel. ic.* 635.
C. ſpinoſa. *Bauh. hiſt.* 2. *p.* 63.

C. ſpinoſa, fructu minore, folio rotundo. *Bauh. pin.* 480.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 4.
Gemeine Capper. W.
Habitat in Europae *auſtralis et* Orientis *ruderatis, muris.* ♄ (v. ſ.)

*2. CAPPARIS *ovata.*
C. pedunculis unifloris ſolitariis, ſtipulis ſpinoſis, foliis ſubrotundo-ovatis acutis glabris, capſulis ovalibus. W.
Capparis ovata. *Desfont. fl. atlant.* 1. *p.* 404.
Capparis ſpinoſa. *β. Sp. pl.* 720.
C. ſicula duplicata ſpina, folio acuto. *Bocc. ſic.* 79. *t.* 42. *f.* 3.
Capparis. *Dod pempt.* 745.
C. folio acuto. *Bauh. pin.* 460. *Lob. ic.* 634.
Eyförmige Capper. W.
Habitat in Sicilia, Hiſpania *et* Africa *boreali.* ♄

*3. CAPPARIS *aegyptia.*
C. pedunculis ſolitariis unifloris, ſtipulis ſpinoſis, foliis rotundo-cuneiformibus apice mucronatis. *Lamarck. encycl.* 1. *p.* 598. *
Aegyptiſche Capper. W.
Habitat in Aegypto. ♄
Rami *teretes.* Spinae *recurvae aureae.* Folia *parva ſemiuncialia glauca petiolata ſubrotundo-cuneiformia evidenter mucronata.* Corolla *albida.* Fructus *triuncialis.* Lamarck. l. c.

*4. CAPPARIS *tomentoſa.*
C. ſpinoſa, floribus axillaribus ſolitariis pedunculatis, foliis ovato-oblongis tomentoſis, ſiliquis ſphaericis. *Lamarck. encycl.* 1. *p.* 599. *
Filzige Capper. W.
Habitat ad Senegal. ♄
Rami *griſei tomentoſi.* Spinae *ſtipulares geminae incurvae.* Folia *uncialia vel ſesquiuncialia ovato-oblonga obtuſa utrinque tenui pube tecta.* Fructus *ſphaericus.* Lamarck. l. c.

*5. CAPPARIS *acuminata.* W.
C. pedunculis unifloris geminatis, ſtipulis ſpinoſis, foliis ovatis acuminatis ſubtus ferrugineo-tomentoſis. W.
Langſpitzige Capper. W.
Habitat in India *orientali.* ♄ (v. ſ.)
Rami *teretes flexuoſi, juniores ferrugineo tomento*

levi tecti. Spinae *ſtipulares geminae rectae.* Folia *petiolata alterna ſesquiuncialia et unciam lata, ſubrotundo-ovata valde acuminata, ſuperne glabra, ſubtus tomento ferrugineo obducta.* Pedunculi *petiolo longiores* 2 *ſ.* 3 *axillares.* Calyx *tomentoſus.* Petala *ciliata.* Stamina *corolla longiora.* W.

Capparis pyrifolia. Lamarck. encycl. 1. *p.* 599. *forte non ab hac diverſa, tamen folia juniora albo-tomentoſa, ſeniora glabra, et formam eorum ovato-lanceolatam dixit. An varietas noſtrae?* W.

6. CAPPARIS *zeylanica.*

C. pedunculis ſolitariis unifloris, ſtipulis ſpinoſis, foliis ovatis utrinque acutis. *Fl. zeyl.* 210. *Loureiro cochin.* 403.

C. zeylanica, duplicatis ſpinis, folio acuto. *Pet. muſ.* 625.

C. ſpinoſa, foliis oblongis. *Burm. zeyl.* 53. *ind.* 118. *Houttuyn Lin. Pfl. Syſt.* 4. *p.* 5.

Zeylaniſche Capper. W.

Habitat in Zeylona. ♄

7. CAPPARIS *horrida.*

C. arborea, ſtipulis aculeatis, ramis flexuoſis, foliis ovato-lanceolatis mucronatis glabris, floribus axillaribus binis. *Lin. ſuppl.* 264.

Steifſtachliche Capper. W.

Habitat in Coromandelia. ♄ (v. ſ.)

Arbor *rigida, ramis flexuoſis, aculeis binis ſtipularibus, ab exortu foliorum rubicundis rigidis.* Folia *petiolata baſi ovata ſenſim attenuata ſpinula terminata.* Pedunculi *bini axillares uniflori.*

Pedunculi 2 *ſ.* 3, *in hac ſpecie ſupra-axillares.* W.

*8. CAPPARIS *erythrocarpos.*

C. pedunculis unifloris ſolitariis, ſtipulis ſpinoſis, foliis oblongis acuminatis mucronatis, pericarpiis hexagonis. W.

C. (*erythrocarpos*) pedunculis ſolitariis unifloris, ſtipulis ſpinoſis, foliis ovatis obtuſis, pericarpiis hexagonis. *Gmel. ſyſt. nat. p.* 806. *Iſert. Act. Soc. Nat. Scrut. Berol.* 9. *p.* 339. *t.* 9. *mala.*

Eckige Capper. W.

Habitat in Guinea. ♄ (v. ſ. ſ. fl.)

Rami *divaricati teretes.* Spinae *ſtipulares geminae*

reflexae. Folia *alterna petiolata oblonga acuminata mucronata glabra, pilis raris ſtellatis praeſertim ad coſtam mediam ſubtus tecta.* Pedunculi *ſolitarii axillares.* Fructus *oblongus hexagonus.* W.

9. CAPPARIS *ſepiaria.*

C. pedunculis umbellatis, ſtipulis ſpinoſis, foliis annuis ovatis emarginatis. *Sp. pl.* 720.

Arbuſcula baccifera ſcandens ſpinoſa, foliis buxi pallidioribus. *Pluk. alm.* 27. *t.* 338. *f.* 3.

Houttuyn Lin. Pſl. Syſt. 4. *p.* 6.

Hecken-Capper. W.

Habitat in India. ♄

Rami *teretes, lignoſi.* Spinae *ſtipulares, geminae, recurvae, breves, nigrae.* Folia *ſubpetiolata, ovata, nuda, integerrima, emarginata.* Umbellae *terminales, ſimplices, multiflorae: Pedunculo communi, brevi.*

*10. CAPPARIS *citrifolia.*

C. ſpinoſa, floribus ad apices ramorum umbellatis, foliis ovato-oblongis glabris coriaceis. *Lamarck. encycl.* 1. *p.* 599. *

Citronenblättrige Capper. W.

Habitat ad Cap. b. ſpei. ♄ (v. ſ.)

*11. CAPPARIS *corymboſa.*

C. floribus corymboſis terminalibus, ſtipulis ſpinoſis, foliis ovalibus ſubtus pubeſcentibus. *Lamarck. encycl.* 1. *p.* 599. *

Doldentraubige Capper. W.

Habitat ad Senegal. ♄

** *Inermes.*

*12. CAPPARIS *mariana.*

C. pedunculis unifloris ſolitariis, ſtipulis nullis, foliis cordato-ovatis. *Iacq. hort. ſchoenb.* 1. *p.* 57. *t.* 109. *

C. (*cordifolia*) pedunculis ſolitariis unifloris, foliis cordatis petiolatis, junioribus tomentoſo-farinoſis. *Lamarck. encycl.* 1. *p.* 602. *

Latroniſche Capper. W.

Habitat in inſulis Marianis. ♄

Habitu ſimilis C. ſpinoſae, ſed diverſa: caule *inermi,* foliis *ovatis baſi cordatis.* Floris *figura eadem ac in C. ſpinoſa.* W.

*13. CAPPARIS *panduriformis.*
C. pedunculis unifloris ad apices ramorum aggregatis, foliis oblongis panduriformibus. *Lamarck. encycl.* 1. *p.* 603. *
Geigenblättrige Capper. W.
Habitat in inſula Mauritii. ♄
Rami *teretes glabri punctati.* Folia *alterna breve petiolata quadriuncialia, et ſesquiunciam lata panduriformia ovata glabra.* Pedunculi *uniflori breves in apice ramorum fere racemoſi.* Lamarck. l. c.

14. CAPPARIS *Baducca.*
C. pedunculis unifloris, foliis perennantibus ovato-oblongis determinate confertis nudis. *Sp. pl.* 720. *Mill. dict. n.* 2.
C. inermis foliis ovato-oblongis per ſpatia confertis perennantibus. *Hort. cliff.* 204. *
Baducca. *Rheed. mal.* 6. *p.* 105. *t.* 57. *Raj. dendr.* 1630. *Burm. ind.* 118.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 8.
Malabariſche Capper. W.
Habitat in India *orientali.* ♄
Folia *utrinque glabra, venis obſolete reticulata, imprimis ſubtus.*
Pedunculi *uniflori terminales et laterales.* Flores *magnitudine C. ſpinoſae.* W.

*15. CAPPARIS *toruloſa.*
C. pedunculis ſubbifloris teretibus terminalibus, foliis lanceolato-ovalibus, ſubtus albido-punctatis, ſiliquis teretibus linearibus toruloſis. *Swartz. prodr.* 81.
Breynia arboreſcens, foliis ovatis utrinque acuminatis, ſiliqua toroſa longiſſima. *Brown. jam.* 246.
Knotige Capper. W.
Habitat in Jamaica. ♄

*16. CAPPARIS *longifolia.* †
C. pedunculis - - , foliis lineari-lanceolatis, ſubtus farinoſo-punctatis. *Swartz. prodr.* 81. *Pluk. phytog. t.* 327. *f.* 6.
Langblättrige Capper. W.
Habitat in Jamaica. ♄

17. CAPPARIS *frondoſa.*
C. pedunculis umbellatis, foliis paſſim confertis. *Sp. pl.* 1674.
C. pedunculis ſubumbellatis, foliis ad intervalla frondoſis. *Jacq. amer.* 162. *t.* 104.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 6.
Buſchige Capper. W.
Habitat Carthagenae *et in* Domingo *in ſylvaticis.* ♄ (v. ſ. ſ. fl.)
Folia *lato-lanceolata.*

18. CAPPARIS *ferruginea.*

C. pedunculis umbellatis, foliis perſiſtentibus lanceolatis ſubtus tomentoſis, floribus octandris. *Amoen. acad.* 5. *p.* 398. * *Swartz. obſ.* 208. *
C. octandra. *Iacq. amer.* 160. *t.* 100. *mala.*
Crataeva fruticoſa, foliis ſingularibus oblongis utrinque acutis ſubtus quaſi villoſis, floribus octandris, racemis comoſis alaribus. *Brown. jam.* 247. *t.* 28. *f.* 1. *bona.*
Houttuyn Lin. Pfl. Syſt. 4. *p.* 7.
Roſtfarbene Capper. W.
Habitat in fruticetis littoribus maris proximis Jamaicae. ♄ (v. ſ.)

19. CAPPARIS *grandis.*

C. arborea mitis, foliis ovatis acutis glabris, corymbis terminalibus, fructu globoſo. *Lin. ſuppl.* 265.
Groſse Capper. W.
Habitat in Zeylona. *Koenig* ♄
Arbor *magna cana.* Folia *alterna ovata integerrima acuta venoſa glabra.* Stipulae *nullae.* Corymbi *terminales, fructiferi in racemum elongati.* Flores *luteo-albidi.* Fructus *majores globoſi.*

*20. CAPPARIS *jamaicenſis.*

C. pedunculis multifloris, foliis oblongis obtuſis emarginatis ſubtus tomentoſis, corollis campanulatis. W.
C. pedunculis multifloris, foliis oblongis emarginatis ſubtus tomentoſis, corollis ſemierectis. *Iacq. amer.* 160. *t.* 101. *
Weſtindiſche Capper. W.
Habitat in Jamaica. ♄

*21. CAPPARIS *odoratiſſima.*

C. pedunculis multifloris, foliis oblongo-lanceolatis acutis ſubtus ſquamoſo-punctatis. W.
C. pedunculis racemoſis, foliis ellipticis perennantibus ſubtomentoſe punctatis. *Iacq. hort. ſchoenb.* 1. *p.* 57. *t.* 110. *
Wohlriechende Capper. W.

Habitat ad Caracas. ♄ (v. ſ. ſ. fl.)

Folia *oblongo-lanceolata baſi attenuata rotundata, apice acuta.* Flores *magnitudine Myrti communis.* Petala *violacea.* Stamina *corolla parum longiora.* Tomentum *foliorum in Capparide hac ut in C. ferruginea et aliis, non eſt tomentum, ſed e ſquamis parvis adpreſſis ut in Hippophaë, nonnullis Ciſti ſpeciebus, et paucis aliis vegetabilibus eſt compoſitum; hinc folia harum plantarum potius ſquamata nuncupanda.* W.

*22. CAPPARIS *verrucoſa.*

C. pedunculis multifloris, foliis oblongis acutis utrinque nudis, fructu verrucoſo. *Iacq. amer.* 159. *t.* 99.*

Warzige Capper. W.

Habitat in ſylvis et fruticetis Carthagenae. ♄

Habitu *convenit cum* C. *Cynophallophora, at folia ſemper oblonga et acuta.* Glandulas *in axillis habet nullas.* Flos *ſtructurae ejusdem, at vix odorus.* Petala *habet candidiſſima et patentiſſima,* Glandulas *nectareas planas puncto umbilicatas,* Stamina *pollicaria corollam haud admodum ſuperantia.* Fructus *eſt ſiliqua oblonga teres nitida tota verrucoſa viridis ſesquipollicaris bivalvis; pulpa alba.* Iacq. l. c.

*23. CAPPARIS *ampliſſima.*

C. pedunculis unifloris terminalibus axillaribusque, foliis ellipticis glabris obtuſinſculis, fructibus obovato-ellipticis teretibus. W.

C. (*ampliſſima*) inermis, foliis ovalibus glabris venoſis, floribus ſolitariis axillaribus et terminalibus, ſtaminibus corolla longioribus, fructu obovato. *Lamarck. encycl.* 1. *p.* 600. *

C. alia arboreſcens lauri foliis, fructu oblongo-ovato. *Plum. ſpec.* 7. *ic.* 75. *f.* 2.

Anſehnlichſte Capper. W.

Habitat in Hiſpaniola. ♄.

24. CAPPARIS *Cynophallophora.*

C. pedunculis multifloris terminalibus, foliis ellipticis obtuſis glabris, glandulis axillaribus, fructibus cylindraceis toruloſis. W.

C. pedunculis multifloris terminalibus, foliis ovalibus obtuſis perennantibus, glandulis axillaribus. *Syſt. nat. ed.* 13. *p.* 358. *Mill. dict. n.* 3. *Swartz. obſ.* 209. *

C. pedunculis multifloris, axillis glanduloſis, foliis oblongis utrinque nitidis. *Iacq. amer.* 158. *t.* 98. *

C. (*flexuoſa*) pedunculis congeſtis terminalibus, foliis perſiſtentibus oblongis obtuſis glabris, ramis flexuoſis. *Sp. pl.* 722. *Swartz. obſ.* 211.
C. arboreſcens, lauri foliis, fructu longiſſimo. *Plum. ſpec.* 7. *ic.* 73. *f.* 1.
Moriſonia flexuoſa. *Amoen. acad.* 5. *p.* 398.
Breynia fruticoſa, foliis oblongis obtuſis. *Brown. jam.* 246. *t.* 27. *f.* 1.
Cynophallophorus ſ. Penis caninus caribaeorum arbor, foliis ſubrotundis clavos pulpiferos phalloides pro ſiliquis gerens. *Pluk.* 126. *t.* 172. *f.* 4.
Acaciae affinis arbor ſiliquoſa, folio ſubrotundo ſingulari, flore ſtamineo albido, ſiliqua tereti ventricoſa interiore tunica mucoſa miniata. *Sloan. jam.* 158. *hiſt.* 2. *p.* 59. *Raj. dendr.* 102.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 9.
Langſchotige Capper. W.
Habitat in America *meridionali.* ♄ (v. ſ. c. fr.)
Fructus *ſiliquoſus, bivalvis.*

*25. CAPPARIS *ſaligna.*

C. foliis lineari-lanceolatis inferne dilatatis utrinque obtuſis glabris, fructu tereti toruloſo. *Vahl. ſymb.* 3. *p.* 66.
Weidenartige Capper. W.
Habitat in inſula St. Crucis. ♄

Rami *teretes glabri, cortice cineraſcente, flexuoſi.* Folia *petiolata alterna lineari-lanceolata tripollicaria, ſuperne ſemiunguem lata, baſin verſus latiora, rarius undique ejusdem latitudinis, integerrima mucronata, baſi rotundata vel ſubcordata venoſa glabra.* Petiolus *brevis.* Glandulae *ſaepe in axiliis foliorum.* Pedunculi *terminales ſolitarii* 2-*ſ.* 3-*flori.* Pedicelli *unguiculares.* Calycis *foliola rotundata coriacea.* Filamenta *numeroſa longiſſima capillaria glabra.* Germen *lineari-oblongum truncatum.* Bacca 2-3-*pollicaris glabra craſſitie pennae cygneae obtuſa, inter ſingula ſemina coarctata.* Pedicellus *fructu longior.* Semina 2 *vel* 5. Vahl. l. c.

26. CAPPARIS *pulcherrima.*

C. racemis erectis terminalibus, foliis ovato-lanceolatis obtuſis, baccis ſubrotundis. W.
C. pedunculis racemoſis, foliis oblongis obtuſis, fructibus baccatis. *Sp. pl.* 1674. *Iacq. amer.* 163. *t.* 106.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 11.
Schönſte Capper. W.
Habitat Carthagenae *Americes in declivibus montium.* ♄ Baccae *ſubrotundae.*

*27. CAPPARIS *tenuiſiliqua.*
C. racemis erectis terminalibus, foliis ovato-lanceolatis acutis, fructibus cylindricis teretibus. W.
C. (*tenuiſiliqua*) racemis ſimplicibus erectis, fructibus ſiliquoſis, foliis diciduis. *Iacq. amer.* 162. *t.* 105. *
Dünnfrüchtige Capper. W.
Habitat in Carthagena Americes. ♄

28. CAPPARIS *linearis.*
C. pedunculis ſubracemoſis, foliis linearibus. *Sp. pl.* 1674. *Iacq. amer.* 161. *t.* 102.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 12. *t.* 9.
Linienblättrige Capper. W.
Habitat Carthagenae *Americes in arenoſis collibus.* ♄

29. CAPPARIS *Breynia.*
C. pedunculis racemoſis, foliis perennantibus oblongis, calycibus pedunculisque tomentoſis, floribus octandris. *Syſt. nat.* 13. *p.* 359. *Mill. dict. n.* 9. *Swartz. obſ.* 210. *
C. (*Breynia*) pedunculis ſubumbellatis, foliis ovali-oblongis acutis ſubtus ſcabris, fructibus tomentoſis. *Iacq. amer.* 161. *t.* 103.
Capparis Cynophallophora. *Sp. pl. ed.* 1. *p.* 503.
C. inermis, foliis ovalibus alternis perennantibus. *Hort. cliff.* 204.
Breynia foliis oblongo-ovalibus. *Roy. lugdb.* 476. *Sp. pl.* 1. *p.* 503.
Breynia elaeagni foliis *Plum. gen.* 40. *Breyn. ic.* 13. *c. fig.*
Breyniae affinis arbor octandra. *Loefl. it.* 207. *
β. C. (*ſiliquoſa*) pedunculis unifloris compreſſis, foliis perſiſtentibus lanceolato-oblongis acuminatis ſubtus punctatis. *Sp. pl.* 721.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 12.
Breyniſche Capper.
Habitat in fruticetis ſiccis Jamaicae *verſus mare.* ♄ Flores *polyandri.* R.

30. CAPPARIS *haſtata.*
C. pedunculis multifloris, foliis haſtato-lanceolatis nitidis. *Sp. pl.* 722. *Iacq. amer.* 159. *t.* 174. *f.* 56.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 13.
Spontonblättrige Capper. W.
Habitat Carthagenae *in ſylvis.* ♄

1011. ACTAEA.

Gen. plant. edit. Schreb. n. 877.

Cor. 4-petala. *Cal.* 4-phyllus. *Bacca* 1-locularis. *Sem.* ſemiorbiculata.

1. ACTAEA *ſpicata.*

A. racemo ovato, fructibus baccatis. *Sp. pl.* 722.

α. Actaea *ſpicata nigra.*

Actaea. *Hort. cliff.* 209. *Fl. ſuec.* 431, 464. *Hort. upſ.* 138. *Roy. lugdb.* 480. *Dalib. pariſ.* 152. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 181. *Crantz. auſtr. p.* 142. *Gunn. norv. n.* 27. *Scop. carn. n.* 653. *Pall. it.* 1. *p.* 50. *Pollich. pal. n.* 503. *Flor. dan. t.* 589. *Blackw. t.* 565. *Hoffm. germ.* 182. *Roth. germ. I.* 228. *II.* 584.

A. foliis circumſerratis, duplicato-pinnatis, pinnula extrema trilobata. *Hall. helv. n.* 1076.

A. caule inermi. *Fl. lapp.* 217.

Aconitum bacciferum. *Bauh. pin.* 183.

Chriſtophoriana. *Cluſ. hiſt.* 2. *p.* 86.

β. Actaea *ſpicata alba.*

A. racemo ovato, fructibus niveis, radice tuberoſa. *Mill. dict. n.* 2.

Aconitum baccis niveis. *Corn. canad.* 76. *t.* 77.

Chriſtophoriana americana racemoſa baccis niveis et rubris. *Moriſ. hiſt.* 2. *p.* 8. *ſ.* 1. *t.* 2. *f.* 7.

γ. Actaea *ſpicata rubra.*

A. baccis rubris. *Ait. Kew.* 2. *p.* 221

Houttuyn Lin. Pfl. Syſt. 7. *p.* 181.

Aehrentragendes Chriſtophskraut. W.

Habitat in nemoribus Europae, β. γ. Americae. ♃ (v. v.)
Petala *rhombea, plano-membranacea.*

2. ACTAEA *racemoſa.*

A. racemis longiſſimis, fructibus ſiccis. *Mat. med.* 133. *Willich. in Nov. act. A. N. C. p.* 107. *Mill. dict. n.* 3.

Actaea racemis longiſſimis. *Gron. virg.* 59.

Chriſtophoriana americana procerior et longius ſpicata. *Dill. elth.* 79. *t.* 67. *f.* 78.

Chriſtophorianae facie herba ſpicata. *Pluk. amalth.* 54. *t.* 383. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 184.
Traubentragendes Chriſtophskraut. W.
Habitat in Florida, Virginia, Canada. ♃ (v. v.)
Petala craſſa, gibba, minima, pedicellata, acumine ſetaceo-acuminata, calycis foliolis plura.

*3. ACTAEA *japonica.*
A. ſpicis longiſſimis, foliolis inciſis palmatis diviſis. *Thunb. jap.* 221.*
Japaniſches Chriſtophskraut. W.
Habitat in Japonia. ♄
Planta *herbacea tota glabra.* Folia *petiolata ternata.* Foliola *ſimplicia cordata inciſa quinqueloba vel ſeptemloba, lobis inciſis, ſerrata, ſubtus pallida, palmam lata et longa.* Petiolus *et petioluli ſtriati longiores.* Flores *ſpicati.* Spica *longiſſima palmaris usque ſpithamaea.* Calyx *et* Corolla *caducae.* Germen *oblongum glabrum.* Stylus *nullus.* Thunb. l. c.

1012. SANGUINARIA.

Gen. pl. ed. Schreb. n. 878.

Cor. 8-petala. *Cal.* 2-phyllus. *Siliqua* ovata, 1-locularis.

1. SANGUINARIA *canadenſis.*
Sanguinaria. *Hort. cliff.* 202. *Gron. virg.* 57. *Mill. dict. n.* 1. *Giſeck. ic. faſc.* 1. *n.* 13.
S. minor, flore ſimplici. *Dill. elth.* 335. *t.* 252. *f.* 326.
Chelidonium majus canadenſe acaulon. *Corn. canad.* 212. *Moriſ. hiſt.* 2. *p.* 257. *ſ.* 3. *t.* 11. *f.* 1. *Raj. hiſt.* 1887.
Ranunculus virginienſis albus. *Park. theat.* 327. *Raj. ſuppl.* 314.
β. Sanguinaria major, flore ſimplici. *Dill. elth.* 335. *t.* 252. *f.* 325.
γ. Sanguinaria major, flore pleno. *Dill. elth.* 335. *t.* 252. *f.* 326.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 185.
Canadiſches Blutkraut. W.
Habitat in America *ſeptentrionali.* ♃ (v. v.)
Folium *radicale tenerrime in ſinu fovet et amplecti-*

tur infantiam floris, more Oſmundae Lunariae, et eſt Folium unicum cucullatum et Scapus *uniflorus e ſingula gemma radicis bivalvi.* Stamina *meae vix habuere* Antheras *polliniferas; an dioica?* Lacteſcens *ſucco fulvo Chelidonii.*

1013. PODOPHYLLUM.

Gen. plant. ed. Schreb. n. 879.

Cor. 9-petala. *Cal.* 3-phyllus. *Bacca* 1-locularis, coronata ſtigmate.

1. PODOPHYLLUM *peltatum.*
P. foliis peltatis lobatis *Sp. pl.* 723.
Podophyllum. *Hort. cliff.* 202. *Hort. upſ.* 137. *Gron. virg.* 57. *Roy. lugdb.* 480. *Trew. ehret. t.* 29. *Mill. dict.*
Anapodophyllum canadenſe. *Cateſb. car.* 1. *p.* 24. *t.* 24.
Aconitifolia humilis, flore albo unico campanulato fructu cynosbati. *Mentz. pug. t.* 11.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 187.
Schildblättriger Entenfuſs. W.
Habitat in America *ſeptentrionali.* ♃ (v. v.)

2. PODOPHYLLUM *diphyllum.*
P. foliis binatis ſemicordatis. *Sp. pl.* 723. *Barton. Act. Soc. Amer.* 3 *p.* 334. *
Houttuyn Lin. Pfl. Syſt. 7. *p.* 188.
Zweyblättriger Entenfuſs. W.
Habitat in Virginia. *Collinſon.* ♃

An Sanguinariae *ſpecies? cum* Folium *unicum binatum et* Scapus *aphyllus radicalis et* Capſula *oblonga.* Flores *non vidi, ſub Podophylli ſpecie miſſum.*

1014. CHELIDONIUM.

Gen. plant. edit. Schreb. n. 880.

Cor. 4-petala. *Cal.* 2-phyllus. *Siliqua* 1-locularis, linearis.

1. CHELIDONIUM *majus.*
C. pedunculis umbellatis. *Mat. med.* 134. *Mill. dict. n.* 1. *Pollich. pal. n.* 504. *Regn. bot. Scop. carn. n.* 634. *Hoffm. germ.* 183. *Blackw. t.* 91. *Ludw. ect. t.*

132. *Kniph. cent.* 8. *n.* 27. *Knorr. del.* 1. *t. S.* 23. *Roth. germ. I.* 228. *II.* 582.

C. lobis foliorum ſubrotundis. *Hall. helv. n.* 1059.

C. pedunculis multifloris. *Hort. cliff.* 202. *Fl. ſuec.* 430, 465. *Hort. upſ.* 137. *Roy. lugdb.* 478. *Dalib. pariſ.* 152.

C. floribus ex ſiliquis breviſſimis, ſ. ch. pedunculis umbellatis, caule piloſo. *Crantz. auſtr. p.* 141.

C. majus vulgare. *Bauh. pin.* 144.

C. majus. *Fuchſ. hiſt.* 865.

β. C. majus, foliis quernis, *Bauh. pin.* 144. *Crantz. l. c.* β. *Kniph. cent.* 8. *n.* 26.

C. (*laciniatum*) foliis quinquelobatis, lobis anguſtis, acute laciniatis. *Mill. dict. n.* 2. *ic. t.* 92.

C. majus, folio laciniato. *Cluſ. hiſt.* 2. *p.* 203.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 189.

Gemeines Schoellkraut. W.

Habitat in Europae *ruderatis.* ♃ (v. v.)

*2. CHELIDONIUM *japonicum.*

C. pedunculis unifloris, foliis petiolatis pinnatis ovatis. *Thunb. jap.* 221.

Japaniſches Schoellkraut. W.

Habitat in Japonia.

Caulis *herbaceus ſtriatus glaber debilis erectus.* Folia *alterna petiolata pinnata cum impari.* Foliola 3 *vel* 5 *oppoſita, breviſſime petiolata ovato-oblonga acuta inciſa inaequaliter ſerrata, ſerraturis acutis ciliatis, glabra, ſubtus pallida, pollicaria et ſesquipollicaria, pinna ultima ſemper majori.* Petioli *cauli ſimilis digitalis.* Flores *axillares ſolitarii pedunculati.* Pedunculus *uniflorus capillaris longitudine fere foliorum.* Calyx *diphyllus glaber.* Corolla *calyce paullo longior lutea.* Stamina *numeroſa corolla quadruplo breviora.* Thunb. l. c.

3. CHELIDONIUM *Glaucium.*

C. pedunculis unifloris, foliis amplexicaulibus ſinuatis, caule glabro. *Oed. dan. t.* 585. *Mill. dict. n.* 3. *Kniph. cent.* 4. *n.* 17. *Knorr. del.* 1. *t. R.* 19. *Regn. bot.* *Hoffm. germ.* 183. *Roth. germ. I.* 228. *II* 583.

Glaucium foliis radicalibus ſemipinnatis, caulinis amplexicaulibus. *Hall. helv. n.* 1060.

Glaucium (*flavum*) caule glabro, ſiliquis cornutis deflexis. *Crantz. auſtr. p.* 141.

Glaucium luteum. *Scop. carn.* 2. *n.* 63.
Chelidonium glabrum, pedunculis unifloris. *Hort. upſ.* 137. *Roy. lugdb.* 473. *Dalib. pariſ.* 152.
C. pedunculis unifloris. *Hort. cliff.* 201. *Gron. virg.* 57.
Papaver corniculatum luteum. *Bauh. pin.* 171.
Papaver corniculatum, flavo flore. *Cluſ. hiſt.* 2. *p.* 91.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 191.
Groſsblütiges Schoellkraut. W.
Habitat in Angliae, Helvetiae, Galliae, Italiae, Virginiae *arenoſis.* ♃ (v. v.)
Planta *glauca.* Caulis *laevis.* Folia *ſcabriuſcula: radicalibus pinnatifidis; caulinis ſolum lobatis.* Flores *flavi.* Siliquae *ſcabrae, diſſepimento in loculamenta, uti ſequentes, diſtinctae.*
Variat corolla rubra. W.

4. CHELIDONIUM *corniculatum.*

C. pedunculis unifloris, foliis ſeſſilibus pinnatifidis, caule hiſpido. *Sp. pl.* 724. *Mill. dict. n.* 4.
C. hiſpidum, pedunculis unifloris. *Hort. upſ.* 137. *Sauv. monſp.* 286.
Glaucium (*phoeniceum*) caule hiſpido, ſiliquis cornutis rectis. *Crantz. auſtr. p.* 141.
Glaucium hirſutum, flore phoeniceo. *Tournef. inſt.* 254.
Papaver corniculatum phoeniceum hirſutum. *Bauh. pin.* 171.
Papaver cornutum, phoeniceo flore. *Cluſ. hiſt.* 2. *p.* 91.
β. Papaver corniculatum phoeniceum glabrum. *Bauh. pin.* 171.
Chelidonium glabrum. *Mill. dict n.* 5.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 193.
Rothes Schoellkraut. W.
Habitat in Hungaria, Bohemia, Auſtria, Monſpelii. ☉ (v. v.)
Planta *viridis.* Folia *omnia pinnatifida.* Flores *rubro-lutei.* Siliquae *hiſpidae.*

5. CHELIDONIUM *hybridum.*

C. pedunculis unifloris, foliis pinnatifidis linearibus, caule laevi, ſiliquis trivalvibus. *Sp. pl.* 724. *Mill dict. n.* 6. *Kniph. cent.* 10. *n.* 19.
Papaver caule ramoſo, foliis linearibus pinnatifidis, capſulis ſulcatis. *Hort. upſ.* 136. *Sauv. monſp.* 252.
Papaver corniculatum violaceum. *Bauh. pin.* 172.
Papaver corniculatum, violaceo flore. *Cluſ. hiſt.* 2. *p.* 92.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 194.

Feinblättriges Schoellkraut. W.
Habitat in Europa *auſtraliore.* ☉ (v. v.)
Stigmata. 3.

1015. PAPAVER.

Gen. plant. edit. Schreb. n. 881.

Cor. 4-petala. *Cal.* 2-phyllus. *Capſ.* 1-locularis, ſub ſtigmate perſiſtente poris dehiſcens.

* *Capſulis hiſpidis.*

1. PAPAVER *hybridum.*

P. capſulis ſubgloboſis toroſis hiſpidis, caule folioſo multifloro. *Sp. pl.* 725. *Mill. dict. n.* 2. *Scholl. barb. n.* 418. *Hoffm. germ.* 183. *Roth. germ. I.* 226. *II.* 580.
P. foliis ternato-pinnatifidis, capſulis ſulcatis hiſpidis. *Hort. cliff.* 210.
P. caule ramoſo, foliis lineari-pinnatifidis, capſulis ſulcatis hiſpidis. *Hort. upſ.* 136.
Argemone capitulo breviore. *Bauh. pin.* 172.
Argemone capitulo hirſuto rotundo torulis canaliculato. *Moriſ. hiſt.* 2. p. 278. *ſ.* 3. *t.* 14. *f.* 9. *Lob. ic.* 276.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 195.
Kleiner Mohn. W.
Habitat in Europa *auſtraliore.* ☉ (v. v.)

2. PAPAVER *Argemone.*

P. capſulis clavatis hiſpidis, caule folioſo multifloro. *Sp. pl.* 725. *Mill. dict. n.* 3. *Scop. carn. n.* 636. *Pollich. pal. n.* 405. *Hoffm. germ.* 183. *Roth. germ. I.* 227. *II.* 580.
P. capſulis hiſpidis oblongo-turbinatis ſulcatis. *Fl. ſuec.* 439, 466. *Dalib. pariſ.* 152. *Gron. orient.* 158.
P. foliis pinnatifidis, fructu oblongo turbinato hiſpido. *Neck. gallob. p.* 229.
P. foliis hiſpidis pinnatis, pinnis lobatis, capſulis ellipticis hiſpidis. *Hall. helv. n.* 1061.
P. foliis ternato-pinnatis, fructu angulato. *Hort. cliff.* 201. *Roy. lugdb.* 419. *Sauv. monſp.* 252.
Argemone capitulo longiore. *Bauh. pin.* 172. *Lob. ic.* 276.

Argemone capitulo tenuiore longiore hirſuto. *Moriſ. hiſt. 2. p. 278. ſ. 3. t. 14. f. 10.*

Houttuyn Lin. Pfl. Syſt. 7 p. 196.

Acker-Mohn. W.

Habitat in Europae *campis arenoſis* ⊙ (v. v.)

Folia tripartita pinnatifida. Pedunculi ſcabri. Calyces *capſulaeque piloſae.*

3. PAPAVER *alpinum.*

P. capſula hiſpida, ſcapo unifloro nudo hiſpido, foliis bipinnatis. *Mill. dict. n. 4. Iacq. auſtr. t. 83. Scop. carn. n. 637.*

P. caule nudo, unifloro, foliis pinnatis, pinnis integris et diviſis, capitulis ſubrotundis hiſpidis. *Hall. helv. n. 1062.*

P. alpinum ſaxatile, coriandri folio. *Segu. ver. 1. p. 416. t. 4. f. 4.*

P. Burſeri. *Crantz. auſtr. p. 138. t. 6. f. 4. ſec. Hall. et Iacq.* R.

Argemone alpina, foliis ſcandicis, lutea. *Bauh. pin. 173. prodr. 93. Burſ. IX. 58.*

Argemone alpina, coriandri folio. *Bauh. pin. 172.*

Argemone minor Penae. *Dalech. hiſt. 1714.*

Houttuyn Lin. Pfl. Syſt. 7. p. 197.

Alpen-Mohn. W.

Habitat in Helvetia, *Schneeberg* Auſtriae, Pyrenaeis. ♃ (v. f.)

Calyx *hirſutus.* Petala *alba baſi flava.* Capſ. *ovata ſtigmatibus* 5.

4. PAPAVER *nudicaule.*

P. capſulis hiſpidis, ſcapo unifloro nudo hiſpido, foliis ſimplicibus pinnato-ſinuatis. *Hort. cliff. 136. Gmel. ſib. 4. p. 180. Gunn. norv. n. 578. Kniph. cent. 10. n. 68. Flor. dan. t. 41. Ait. Kew. 2. p. 224.*

P. erraticum, nudicaule flore flavo odorato. *Dill. elth. 202. t. 224. f. 291.*

P. erraticum, luteo flore, capite oblongo hiſpido, et laevi. *Amm. ruth. 61. n. 81.*

Houttuyn Lin. Pfl. Syſt. 7. p. 198.

Nacktſtieliger Mohn. W.

Habitat in Sibiria. ♂ (v. v.)

Pedunculi *et* Calyces *piloſi. Nimis affine* P. alpino

et variat corolla alba et lutea; an cultura ſolum diverſum?

Variat flore albo. Ait. l. c.

** * Capſulis glabris.*

5. PAPAVER *Rhoeas.*

P. capſulis glabris globoſis, caule piloſo multifloro, foliis pinnatifidis inciſis. *Mat. med.* 134. *Pollich. pal. n.* 507. *Crantz. auſtr. p.* 137. *n.* 1. *Pallas itin.* 3. *p.* 546. *Scop. carn. n.* 638. *Ludw. ect. t.* 17. *Blackw. t.* 2. *et* 560. *Kniph. cent.* 5. *n.* 63. *et* 64. *Knorr. del.* 1. *t. R.* 14. *Hoffm. germ.* 183. *Roth. germ. I.* 227. *II.* 581.

P. foliis ſemi-pinnatis, hiſpidis, fructu ovato glabro. *Hall. helv. n.* 1064.

P. foliis bipinnatifidis alternis, fructu glabro ovato. *Neck. gallob. p.* 229.

P. foliis pinnatifidis hiſpidis, fructu ovato. *Hort. cliff.* 201. *Fl. ſuec.* 428, 468. *Roy. lugdb.* 478. *Dalib. pariſ.* 151.

P. caule multifloro folioſo hiſpido, foliis pinnatifidis, capſulis laevibus. *Hort. upſ.* 136. *Sauv. monſp.* 252. *Gron. orient* 157.

P. erraticum majus. *Bauh. pin.* 171.

P. rhoeas prius. *Dalech. hiſt.* 439.

β. P. erraticum, pleno flore. *Bauh. pin.* 171. *Knorr. del. t. R.* 12. 13.

γ. P. erraticum minus. *Bauh. pin.* 171.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 198.

Wilder Mohn. W.

Habitat in Europae *arvis, agris.* ⊙ (v. v.)

Pili in caule patentes ſunt.

Varietas *habetur caule ſimpliciſſimo, capſula oblonga, flore albo.* Hall. Crantz. R.

6. PAPAVER *dubium.*

P. capſulis oblongis glabris, caule multifloro ſetis adpreſſis, foliis pinnatifidis inciſis. *Fl. ſuec.* 2. *n.* 467. *Pollich. pal. n.* 506. *Hoffm. germ.* 184. *Roth. germ. I.* 227. *II.* 581.

P. foliis hiſpidis pinnatis, pinnis lobatis, fructu ovato laevi. *Hall. helv. n.* 1063.

P. capſulis oblongis, caule multifloro, ſetis adpreſſis, foliis pinnatifidis inciſis. *Crantz. auſtr. p.* 140. *cum* P. argemone *conjungit.* R.

P. erraticum, capite longiſſimo glabro. *Tournef. inſt.* 238. *Rupp. jen.* 2. *p.* 71. *ed. Hall. p.* 79.
Argemone capitulo longiore glabro. *Moriſ. bleſ.* 233. *hiſt.* 2. *p.* 279 *ſ.* 3. *t.* 14. *f.* 11.
β. Papaver dubium. *Iacq. auſtr.* 1. *p.* 17. *t.* 25.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 200.
Saat-Mohn. W.
Habitat inter ſegetes Europae *ſeptentrionalioris.* ⊙ (v. v.)
Papaver *Iacquini in flor. auſtr. t.* 25. *pictum*, Petala *crenata eaque tantum non ſemper alba habet; apud nos vero* P. dubium *petala conſtanter rubra, et ſemper integra gerit.* De la Chenal in *Act. Helv.* VIII. *p.* 145. *n.* 25.

7. PAPAVER *ſomniferum.*
P. calycibus capſulisque glabris, foliis amplexicaulibus inciſis. *Mat. med. p.* 154. *Mill. dict. n.* 8. *et* 9. *Hall. helv. n.* 1065.
P. caule multifloro, foliis ſimplicibus glabris inciſis caulinis. *Hort. upſ.* 135.
P. foliis ſimplicibus glabris inciſis. *Hort. cliff.* 200. *Roy. lugdb.* 879.
P. nigrum et album offic. *Crantz. auſtr. p.* 138. *Blackw. t.* 482. 483. *Regn. bot.*
P. hortenſe, ſemine albo. *Bauh. pin.* 170.
β. P. hortenſe, ſemino nigro. *Bauh. pin.* 170.
γ. P. criſtatum, floribus et ſemine album. *Bauh. pin.* 171.
δ. P. criſtatum, floribus rubris, ſemine nigro. *Bauh. pin.* 171.
ε. P. flore pleno, album. *Bauh. pin.* 171.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 200.
Garten-Mohn. W.
Habitat in Europae *auſtralioris ruderatis.* ⊙ (v. v.)
Caulis, Folia, Calyces, Capſulae glabra. Stigmata 10.

8. PAPAVER *cambricum.*
P. capſulis glabris oblongis, caule multifloro laevi, foliis pinnatis inciſis. *Hudſ. angl.* 203. *Mill. dict. n.* 5
P. foliis pinnatis, fructu acuminato. *Hort. cliff.* 201. *Roy. lugdb.* 479.
P. cambricum perenne, flore ſulphureo. *Dill. elth.* 300. *t.* 223. *f.* 290.

Argemone cambro-britannica lutea, capite longiore glabro. *Moriſ. hiſt.* 2. *p.* 297. *ſ.* 3. *t.* 14. *f.* 12.
P. erraticum pyrenaicum, flore flavo. *Bauh. pin.* 171. *prodr.* 92. *Burſ. IP.* 45.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 204.
Engliſcher Mohn. W.
Habitat in Cambriae *ſeptentrionalis nemoroſis,* Pyrenaeis. ♃ (v. ſ.)
Stigmata 5 *ſ.* 6 *diſtincta. Capſula oblonga, inferne anguſtior, apice ſubroſtrata.*

9. PAPAVER *orientale.*
P. capſulis glabris, caulibus unifloris ſcabris folioſis, foliis pinnatis ſerratis. *Hort. upſ.* 136. *Mill. dict. n.* 7. *Knorr. del. t. R.* 14. *a.*
P. foliis pinnatis, fructu globoſo. *Roy. lugdb.* 279.
P. orientale hirſutiſſimum, flore magno. *Tournef. cor.* 17. *itin.* 3. *p.* 127. *t.* 127. *Comm. rar.* 34. *t.* 34.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 205.
Orientaliſcher Mohn W.
Habitat in Oriente. ♃ (v. v.)
Stigmata 16. Setae *in caule ſparſae, cauli adpreſſae, baſi prominula aſperae.*

1016. ARGEMONE.

Gen. pl. ed. Schreb. n. 882.

Cor. 6-petala. *Cal.* 3-phyllus. *Capſ.* ſemivalvis.

1. ARGEMONE *mexicana.*
A. capſulis ſexvalvibus, foliis ſpinoſis. *Sp. pl.* 727.
A. capſulis quinquevalvibus. *Sp. pl.* 2. *p.* 727. *Mill. dict. Kniph. cent.* 5. *n.* 16.
Argemone. *Hort. cliff.* 201. *Hort. upſ* 135. *Roy. lugdb.* 479. *Brown. jam.* 244. *Merian. ſurin.* 24. *t.* 24.
Papaver ſpinoſum. *Bauh. pin.* 171. *prodr.* 93. *t.* 92. *Cluſ. hiſt.* 2. *p.* 93.
Papaver ſpinoſum luteum, foliis albis venis notatis. *Moriſ. hiſt.* 2. *p.* 277. *ſ.* 3. *t.* 14. *f.* 5.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 207.
Mexicaniſche Argemone. W.
Habitat in Mexico, Iamaica, Caribaeis, *nunc in* Europa *auſtrali.* ⊙ (v. v.)

2. ARGEMONE *armeniaca.*

A. capſulis trivalvibus. *Sp. pl.* 727.
Papaver orientale, hypecoi folio, fructu minimo. *Tournef. cor.* 17.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 208.
Armeniſche Argemone. W.
Habitat in Armenia.

3. ARGEMONE *pyrenaica.*

A. capſulis quadrivalvibus, caule nudo. *Sp. pl.* 728.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 208.
Pyrenaiſche Argemone. W.
Habitat in Pyrenaeis. *Tournef.*

Haec *ſemiquadrivalvis, uti ſpecies ſecunda ſemitrivalvis, et prima ſemiquinquevalvis.* Hae *ſpecies vel ſub diſtincto genere ſimul tradendae, vel etiam ad genus papaverinum amandandae omnes. Hujus habitus plurimum accedit Papaveri alpino.*

1017. MUNTINGIA.

Gen. pl. ed. Schreb. n. 884.

Cor. 5-petala. *Cal.* 5-partitus. *Bacca* 5-locularis, 1-polyſperma.

1. MUNTINGIA *Calabura.*

M. Calabura. *Sp. pl.* 728. *Iacq. amer.* 166. *t.* 107. *Mill. dict. Swartz. obſ.* 211. *
M. pedunculis unifloris. *Hort. cliff.* 203. *
M. fruticoſa villoſa, foliis ſerratis oblongis uno latere brevioribus. *Brown. jam.* 245.
M. folio ſericeo molli, fructu majori. *Plum. gen.* 41. *ic.* 205.
Calabura alba. *Plum. alm.* 75. *mant* 34. *t.* 152. *f.* 4.
Loti arboris folio anguſtiore, rubi flore, fructu polyſpermo umbilicato. *Sloan. jam.* 162. *hiſt.* 2. *p.* 80. *t.* 194. *Raj. dendr.* 32.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 15.
Weſtindiſche Muntingie. W.
Habitat in Jamaicae *collibus calcareis ſubalpinis.* ♄
Flores *axillares, bini, albi.*

1018. SARRACENIA.

Gen. pl. ed. Schreb. n. 885.

Cor. 5-petala. *Cal.* duplex, 3-phyllus et 5-phyllus. *Capſ.* 5-locularis. *Stylus* ſtigmate clypeatus.

1. SARRACENIA *flava.*

S. foliis erectis tubulatis, valva collo contracto, apice plana erecta. *Walt. carol.* 153.

S. foliis ſtrictis. *Sp. pl.* 729. *Mill. dict. n.* 2.

Sarracena foliis rectis. *Hort. cliff.* 497. *Gron. virg.* 164.

Sarracena foliis longioribus et anguſtioribus. *Catesb. car.* 2. *p.* 69. *t.* 69.

Coilophyllum virginianum, longiore folio erecto, flore luteo. *Moriſ. hiſt.* 3. *p.* 533.

Bucanephyllum elatius virginianum ſ. Limonio congeneris altera ſpecies elatior, foliis triplo longioribus. *Pluk. alm.* 72. *amalth.* 46. *t.* 152. *f.* 3. *(mala) t.* 376. *f.* 5. *bona.*

Thuris limpidi folium. *Bauh. hiſt.* 1. *p.* 307. *Dalech. hiſt.* 1754. *Lob. adv.* 430.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 210.

Gelbe Sarracene. W.

Habitat in Americae *ſeptentrionalis udis.* ♃

*2. SARRACENIA *minor.* †

S. foliis minoribus tubulatis erectis, valva concava nutante. *Walt. carol.* 153.

Kleine Sarracene. W.

Habitat in Carolinae *udis.* ♃

*3. SARRACENIA *rubra.* †

S. foliis erectis tubulatis, valva plana erecta. *Walt. carol.* 152.

Rothe Sarracene. W.

Habitat in Carolinae *udis.* ♃

4. SARRACENIA *purpurea.*

S. foliis cucullatis ventricoſis patulis arcuatis. *Walt. carol.* 152.

S. foliis gibbis. *Sp. pl.* 728. *Hort. cliff.* 427. *Gron. virg.* 164. *Mill. dict. n.* 1. *et ic.* 161. *t.* 241.

Sarracena foliis brevioribus latioribus. *Catesb. car.* 2. *p.* 70. *t.* 70.

Coilophyllum virginianum, breviore folio, flore purpuraſcente. *Moriſ hiſt. 3. p. 533.*
Bucanephyllum americanum, limonio congener dictum. *Pluk. alm. 71. amalth. 46 t. 376. f. 6.*
Limonium peregrinum, foliis forma floris ariſtolochiae. *Bauh. pin. 192.*
Houttuyn Lin. Pfl. Syſt. 7. p. 209.
Purpurfarbene Sarracene. W.
Habitat in Americae *ſeptentrionalis udis.* ♃ *P. Collinſon.* (v. ſ.)

Folia *ſeſſilia, cucullata: tubo ventricoſo, ſenſim ampliato: fauce contractiuſcula, margine antice dilatata.* Limbus *reniformis, patulus, emarginatus.* Venter *antice carina membranacea, ſemiellipiica, binervi. Sic Metamorphoſis folii Nymphaeae in folium* Sarraceniae, *ut ipſa aquam pluvialem excipiens et retinens extra aquas creſcat; mira Naturae providentia!*

1019. NYMPHAEA.

Gen. plant. edit. Schreb. n. 886.

Cor. polypetala. *Cal.* 4-ſ. 5-phyllus. *Bacca* multilocularis, loculis polyſpermis.

1. NYMPHAEA *lutea.*

N. foliis cordatis integerrimis, lobis approximatis, calyce pentaphyllo petalis longiore. *Ait. Kew. 2. p. 227.*
N. foliis cordatis integerrimis, calyce petalis majore pentaphyllo. *Fl. lapp. 218. Fl. ſuec. 426, 469. Hort. cliff. 203. Roy. lugdb. 480. Dalib. pariſ. 150. Mill. dict. n. 1. Gmel. ſib. 4. p. 184. Crantz. auſtr. p. 142. Hoffm. germ. 184. Scop. carn. n. 639. Gunn. norv. n. 451. Pollich. pal. n. 508. Flor. dan. t. 603. Blackw. t. 497. 499. et 500. Kniph. cent. 3. n. 66. Knorr. del. 2. tab. N. 1. Roth. germ. I. 229. II. 588.*
N. calyce pentaphyllo, petalis majori, fructu conico. *Hall. helv. n. 1066.*
N. foliis cordato-ovatis, calycibus pentaphyllis orbiculatis: petalis ſubcuneatis. *Neck. gallob. p. 228.*
N. lutea major. *Bauh. pin. 183.*
N. lutea. *Cam. epit. 635.*
β. N. *minima* petiolis baſi ſemiteretibus, ſuperne ancipitibus, ſtigmate dentato. W.

Nymphaea lutea β. pumila. *Timm. in Magazin für die Naturkunde Mecklenburgs. 3. p. 256.* *
Houttuyn Lin. Pfl. Syſt. 7. p. 211.
Gelbe Seeroſe. W.
Habitat ad Europae *littora, ſub dulci aqua. β. in ducatu* Megapolitano. ♃ (v. v.)

Varietas β, eſt forte diſtincta ſpecies, differt a vulgari, flore *magnitudine Ranunculi acris,* ſtigmate *dentato, et* petioli *forma.* W.

*2. NYMPHAEA *advena.*

N. foliis cordatis integerrimis, lobis divaricatis, calyce hexaphyllo petalis longiore. *Ait. Kew. 2. p. 226.*
N. floribus flavis. *Gronov. virg. ed. 1. p. 164.*
Dreyfarbige Seeroſe. W.
Habitat in America *boreali.* ♃

Petioli *ſemiteretes plerumque ſupra aquam eriguntur* Folia *cordato-oblonga verſus apicem magis anguſtata, quam in N. lutea.* Pedunculus *teres.* Calyx *hexaphyllus;* foliola *ſubrotunda obtuſa concava:* tria exteriora *patentia extus viridia, intus ſordide purpuraſcentia, diametro unciali;* tria interiora *duplo majora ſuberecta, extus flava, intus ſordide purpuraſcentia.* Petala *circiter 13, lateribus receptaculi ſimplici ſerie inſerta, ſubcuneiformia, obtuſe rotundata patenti-reflexa, luteo-flavicantia, tres lineas lata.* Filamenta *numeroſiſſima, multiplici ſerie receptaculo inſerta linearia, lineam lata, vix ſemiunciam longa, primum erecta, dein reflexa flava, medio rubra.* Antherae *lineares, binae in ſingulo filamento, marginibus adnatae.* Germen *columnare, profunde ſulcatum, angulis* (13) *extantibus, flavum.* Stigma *peltatum leviter umbilicatum flavum, radiis 13 rotatum, margine profunde 13-dentatum, dentibus obtuſis ſuperne viridibus.* Ait. l. c.

3. NYMPHAEA *alba.*

N. foliis cordatis integerrimis, lobis imbricatis rotundis, calyce tetraphyllo. *Ait. Kew. 2. p. 227.*
N. foliis cordatis integerrimis, calyce quadrifido. *Mat. med. p. 135. Mill. dict. n. 2. Crantz. auſtr. p. 142. Pollich. pal. n. 509. Hoffm. germ. 184. Scop. carn. n. 640. Fl. dan. t. 602. Blackw. t. 498, 499. Knorr. del. 2. t. N. Roth. germ. I. 230. II. 589.*
N. calyce tetraphyllo, corolla multiplici. *Fl. lapp. 219.*

Fl. ſuec. 427. 470. *Hort. cliff.* 203. *Roy. lugdb.* 480. *Dalib. pariſ.* 150.

N. calyce quadrifido, petalis minori, fructu ſphaerico. *Hall. helv n.* 1067.

N. foliis cordato - orbiculatis, petalis ovatis calycem aequantibus: fructu truncato. *Neck. gallob. p.* 227.

N. alba major. *Bauh. pin.* 193.

N. alba. *Cam. epit.* 634.

β. N. alba minor. *Besl. eyſt. vern. ord.* 7. *t.* 3. *f.* 1. *Moriſ. hiſt.* 3. *p.* 513.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 212.

Weiſse Seeroſe. W.

Habitat in Europae *aquis dulcibus.* ♃ (v. v.)

*4. NYMPHAEA *odorata.*

N. foliis cordatis integerrimis emarginatis, lobis divaricatis, acumine obtuſo, calyce tetraphyllo. *Ait Kew.* 2. *p.* 227.

N. alba flore pleno odorato. *Gron. virg. ed.* 1. *p.* 57. *ed.* 2. *p.* 81.

N. alba minor. *Gmel. ſib.* 4. *p.* 184. *t.* 72. *(excluſo ſynonymo Moriſoni.)*

Wohlriechende Seeroſe. W.

Habitat in America *boreali inque* Sibiria *orientali.* ♃

Omnino diverſa a N. alba minore Moriſoni et aliorum, quae a N. alba, nil niſi magnitudine differt. Ait. l. c.

*5. NYMPHAEA *ſtellata.* W. †

N. foliis cordatis integerrimis, lobis divaricatis, acutis, calyce acuto tetraphyllo petalis acutis longiore. W.

Citambel. *Rheed. mal.* 11. *p.* 53. *t.* 27.

Sternförmige Seeroſe. W.

Habitat in Malabariae *aquis.* ♃

6. NYMPHAEA *Lotus.*

N. foliis cordatis dentatis glaberrimis, lobis approximatis, calyce tetraphyllo. W.

N. foliis cordatis dentatis. *Fl. zeyl.* 194. *Haſſelq. it.* 471.

N. foliis amplioribus profunde crenatis ſubtus areolatis. *Brown. jam.* 243?

N. indica, flore candido, folio in ambitu ſerrato. *Sloan. jam* 120. *hiſt.* 1. *p.* 252? *Raj. ſuppl.* 630?

Lotus aegyptia. *Alp. aegypt.* 103. *exot.* 214. *t.* 213. 216. 218. 220. 222. 226.

Ambel. *Rheed. mal* 11. *p.* 51. *t.* 26.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 213.
Aegytipſche Seeroſe. W.
Habitat in Indiae, Aegypti, Hungariae *aquis.* ♃ (v. v.)
Dentes *foliorum acuminati: Planta americana ſub eodem nomine a Botanicis indicata, videtur differre.* W.

*7. NYMPHAEA *pubeſcens.* W.
N. foliis reniformibus dentatis ſubtus pubeſcentibus, lobis rotundis, calyce tetraphyllo. W.
N. indica foliis amplis eleganter ad ambitum crenatis, ſubtus fulva lanugine tecta, flore albo ſimplici. *Pluk. alm.* 267.
Rauhe Seeroſe. W.
Habitat in India *orientali.* ♃ (v. ſ.)
Folia *ſex uncias et ultra longa, octo vel decem uncias lata reniformia, nec cordata, lobis rotundata, margine profunde et acute dentata, ſuperne ſcabra, ſubtus pilis denſis breviſſimis fuſceſcentibus tecta.* Petiolus *pubeſcens, videtur in ſicco ſpecimine ſemi-teres.* Flos *ut in Nymphaea alba.* Calyx *tetraphyllus viridis obtuſus.* Petala *numeroſa.* W.

1020. B I X A.

Gen. plant. éd. Schreb. n. 887.

Cor. 10-petala. *Cal.* 5-dentatus. *Capſ.* hiſpida, bivalvis.

1. BIXA *Orellana.*
Bixa. *Hort. cliff.* 211. *Mat. med.* 135. *Roy. lugdb.* 477. *Brown. jam.* 254. *Mill. dict.*
Orleana ſ. Orellana folliculis lappaceis. *Pluk. alm.* 272. *t.* 209. *f.* 4. *Comm. hort.* 1. *p.* 65. *f.* 33.
Arbor mexicana, fructu caſtaneae, coccifera. *Bauh. pin.* 419.
Urucu. *Sloan. jam.* 150. *hiſt.* 2. *p.* 52. *t.* 181. *f.* 1. *Piſ. braſ.* 133.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 9.
Gemeiner Orleanbaum. W.
Habitat in America *calidiore, ad foſſas et fontes.* ♄ (v. ſ. ſ. fl.)

1021. SLOANEA.

Gen. plant. edit. Schreb. n. 888.

Cor. 0. *Cal.* 1-phyllus 5-9-fidus. *Antherae* filamentis infra apicem adnatae. *Capſ.* echinata 3-6-locularis, 3-6-valvis. *Semina* 2 arillo baccato.

1. SLOANEA *dentata*.

S. foliis ovatis, ſtipulis cordato-triangularibus ſerratis. *Swartz. prodr.* 82. *obſerv.* 213.*
S. foliis cordato-ovatis denticulatis, ſtipulis ſerratis. *Sp. pl.* 730.
Sloanea Plumierii. *Aubl. guj.* 1. *p.* 536.*
Sloanea. *Hort. cliff.* 210?
S. amplis caſtaneae foliis, fructu echinato. *Plum. gen.* 49. *ic.* 244.
Caſtanea foliis oblongo-ovatis ſerratis, fructu rotundo maximo echinato. *Mill. dict.*
Houttuyn Lin. Pfl. Syſt. 2. *p.* 12.
Gezähnte Sloanie. W.
Habitat in America *meridionali.* ♄

*2. SLOANEA *Maſſoni*.

S. foliis cordato-ellipticis, ſtipulis linearibus, calyce quinquepartito, ſetis capſulae longiſſimis. *Swartz. prodr.* 82.
Maſſoniſche Sloanie. W.
Habitat in inſulis Caribaeis. ♄

*3. SLOANEA *ſinemarienſis*.

S. foliis ſubrotundo-ovatis integerrimis, capſulis ovatis ſetoſis ex apice dehiſcentibus. *Swartz prodr.* 82.
S. (*ſinemarienſis*) foliis ſubrotundis integerrimis, fructu parvo aculeis hiſpido. *Aubl. guj. p.* 534. *t.* 212.*
Sinemarienſiſche Sloanie. W.
Habitat in inſulis Caribaeis *et ſylvis* Gujanae. ♄

Sloanea (emarginata) foliis oblongis integerrimis emarginatis Sp. pl. 731. *eſt planta valde dubia, forte nullibi obvia, cujus deſcriptionem Clariſſ. Linnaeus ex icone Cateſbaei concinnavit, ſed planta hujus Auctoris eſt Achras Sapota.* W.

*1022. AUBLETIA.

Gen. pl. ed. Schreb. n. 889.

Cal. 5-phyllus. *Cor.* 5 petala. *Capſula* echinata multilocularis.

*1. AUBLETIA *Tibourbou.*

A. foliis cordato-lanceolatis ſerratis ſubtus hirſutis, capſulis ſetoſis. W.

Apeiba (*Tibourbou*) foliis acute-ſerratis hirſutis. *Swartz. prodr.* 82. *Ait. Kew.* 2. *p.* 228.

Apeiba (*Tibourbou*) foliis cordatis ovato-oblongis acutis ſerrulatis, ſubtus hirſutis ruſeſcentibus, fructu magno compreſſo hiſpido. *Aubl. guj.* 1. *p.* 538. *t.* 213. *

Sloanea. *Loefl. itin.* 511. *

Apeiba. *Marcg. braſ.* 123. *t.* 124. *Raj. hiſt.* 1643.

Braunhaarige Aubletie. W.

Habitat in collibus Gujanae *et* Cajennae *inque inſula* Tabago. ♄ (v. ſ.)

*2. AUBLETIA *Petoumo.*

A. foliis oblongis ſubcordatis ſerrulatis ſubtus incanis, capſulis ſetoſis. W.

Apeiba (*Petoumo*) foliis ellipticis acutis ſerrulatis ſubtus incanis. *Swartz. prodr.* 82.

Apeiba (*Petoumo*) foliis ovatis ſubtus incanis, floribus racemoſis luteis. *Aubl. guj.* 1. *p.* 543. *t.* 215. *

Graue Aubletie. W.

Habitat in ſylvis Gujanae. ♄

*3. AUBLETIA *aſpera.*

A. foliis oblongis ſubcordatis integerrimis ſubtus pubeſcentibus, capſulis muricatis. W.

Apeiba (*aſpera*) foliis integerrimis ſubtus pubeſcentibus, fructu compreſſo. *Swartz. prodr.* 82.

Apeiba (*aſpera*) fructu rotundo compreſſo leviter echinato. *Aubl. guj.* 1. *p.* 545. *t.* 216. *

Scharffrüchtige Aubletie. W.

Habitat in Cajenna. ♄

*4. AUBLETIA *laevis.*

A. foliis oblongo-obovatis acuminatis integerrimis glabris, petalis obtuſis, capſulis ſcabris. W.

Apeiba (*laevis*) foliis integerrimis utrinque glabris, fructu aſpero depreſſo. *Swartz. prodr.* 83.

Apeiba (*glabra*) floribus vireſcentibus, fructu aſpero. *Aubl. guj.* 1. *p.* 541. *t.* 214.*
Glatte Aubletie. W.
Habitat in Cajenna. ♄

1023. M A M M E A.

Gen. plant. edit. Schreb. n. 1586.

Cor. 4-petala. *Cal.* 2-phyllus. *Bacca* maxima, 4-ſperma.

1. MAMMEA *americana.*

M. foliis obtuſiſſimis ſtriatis, pedunculis brevibus, baccis tetraſpermis. *Vahl. eclog.* 2. *p.* 40.
M. ſtaminibus flore brevioribus. *Sp. pl.* 731. *Mill. dict. n.* 1. *Iacq. amer.* 268. *t.* 181. *f.* 82.
M. foliis ovalibus nitidis, fructu ſubrotundo ſcabro. *Brown. jam.* 248.
Mammei magno fructu, perſicae ſapore. *Plum. gen.* 44. *ic.* 170.
Mamay. *Bauh. hiſt.* 1. *p.* 172. *Dalech. hiſt.* 1836. *Laet. amer.* 356.
Malus perſica maxima, foliis rotundis ſplendentibus glabris, fructu maximo ſcabro rugoſo. *Sloan. jam.* 179. *hiſt.* 2. *p.* 123. *t.* 217. *f.* 3.
Arbor indica mamei dicta. *Bauh. pin.* 417. *Raj. hiſt.* 1665.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 16.
Genieſsbare Mammei. W.
Habitat in Hiſpaniola, Jamaica. ♄

Arbor *alta, pyramidalis, ramis ſcabris.* Folia *oppoſita, oblonga, obtuſa.* Fructus *fulvus, magnitudine capitis infantis, gratiſſimus, ſed cortex et pulpa ſeminum propria amariſſima. Ad polygamiam monoeciam.* Iacq.

*2. MAMMEA *humilis.*

M. foliis acutis, pedunculis petiolo longioribus, baccis triſpermis. *Vahl. eclog.* 2. *p.* 40.*
Niedrige Mammei. W.
Habitat in inſula Montſerrat. ♄

Eandem forte eſſe plantam ac Rheedia lateriflora Linnaei, credit Clariſſ. Vahl. l. c. W.

1024. OCHNA.

Gen. plant. ed. Schreb. n. 891.

Cor. 5-petala. *Cal.* 5-phyllus. *Baccae* 1-ſpermae, receptaculo ſubrotundo magno affixae.

1. OCHNA *ſquarroſa.*

O. pedunculis multifloris, corollis octopetalis. W.
O. racemis lateralibus. *Sp. pl.* 731. *Fl. zeyl.* 209. * *Roxb. corom.* 1. *p.* 62. *t.* 89. *

Japotapita cinnamomi folio, floribus ſpicatis. *Burm. zeyl.* 123. *t.* 56.

β. Arbor africana, ſubrotundo folio margine denticulis acutis aſpero, floribus pentapetalis atropurpureis. *Pluk. alm.* 41. *t.* 263. *f.* 1. 2.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 21.
Sparrige Ochna. W.
Habitat in India, β. *in* Africa. ♄ (v. ſ.)

Racemi *perſiſtunt ſquarroſi infra folia a caſu pedicellorum et fructificationum.*

Si planta Plukenetii ad hoc genus pertineat, erit ſpecies diſtincta; differt enim, quoad iconem foliis obtuſis, et corollis pentapetalis, ut et colore petalorum rubro, qui flavus eſt in O. ſquarroſa. W.

*2. OCHNA *parvifolia.*

O. pedunculis unifloris. *Vahl. ſymb.* 1. *p.* 33.
Evonymus (*inermis*) foliis alternis ovatis ſerratis. *Forſk. deſcr.* 204. *
Kleinblättrige Ochna. W.
Habitat in Arabia *felici.* ♄

Refert O ſquarroſam, demta magnitudine foliorum et inflореſcentia. Frutex *ramis teretibus glabris, cortice cinereo punctato.* Folia *tantummodo in ramulis petiolata oblonga ſerrulata baſi integerrima venoſa nitida unguicularia.* Pedunculi *ad latera ramulorum ſolitarii uniflori, ſuperne ſenſim incraſſati purpurei folio longiores.* Vahl. l. c.

1025. G R I A S.

Gen. plant. ed. Schreb. n. 892.

Cor. 4-petala. *Cal.* 4-fidus. *Stigma* ſeſſile, cruciatum. *Drupa* nucleo octoſulcato.

1 GRIAS *cauliflora.*

G. cauliflora. *Sp. pl.* 732. *Swartz. obſ.* 215.*

Calophyllum foliis tripedalibus obovatis, floribus per caulem et ramos ſparſis. *Brown. jam.* 245.

Palmis affinis Malus Perſica maxima, candice non ramoſo, foliis longiſſimis, flore tetrapetalo pallide luteo, fructu ex arboris trunco. *Sloan. jam.* 179. *hiſt.* 2. *p.* 123. *t.* 217. *f.* 1. 2.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 24.

Stammblütige Grias. W.

Habitat in Jamaicae *uliginoſis ſubalpinis.* ♄

Arbor *ſaepe* 50-*pedalis.* Rami *terminales ſimplices breves vel nulli.* Folia *brevius petiolata pendula longiſſima* 2-3-*pedalia, baſi cuneata oblongo-attenuata integerrima venoſa rugoſa glabra conſiſtentia.* Flores *caulini pedunculati.* Pedunculi *breviſſimi ſquamoſi multiflori.* Pedicelli *breves conferti uniflori.* Corollae *magnae albidae.* Swartz. l. c.

1026. C A L O P H Y L L U M.

Gen. plant. ed. Schreb. n. 1587.

Cor. 4-petala. *Cal.* 4-phyllus, coloratus *Drupa* globoſa.

1. CALOPHYLLUM *Inophyllum.*

C. foliis ovalibus. *Sp. pl.* 732. *Fl. zeyl.* 101.*

Arbor indica, mali medicae amplioribus foliis. *Pluk. alm.* 41. *t.* 147. *f.* 3.

Bintangor maritima. *Rumph. amb.* 2. *p.* 211. *t.* 71.

Ponna ſ. Ponna-maram. *Rheed. mal.* 4. *p.* 76. *t.* 38. *Raj. hiſt.* 1525.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 25.

Groſses Schönblatt. W.

Habitat in India. ♄ (v. ſ.)

Ex hac arbore venit Reſina Tacamahaca dicta; quae in inſulis Borboniae, Madagaſcariae colligitur. Lamarck. encycl. 1. p. 457.

2. CALOPHYLLUM *Calaba.*
C. foliis ovatis obtuſis. *Sp. pl.* 732. *Fl. zeyl.* 202.* *Iacq. amer.* 269. *t.* 165.* *Swartz. obſ.* 216.*
C. foliis ovatis obtuſis: ſtriis parallelis transverſis. *Hort. cliff.* 206. *Roy. lugdb.* 476.
Inophyllum flore quadrifido. *Burm. zeyl.* 130. *t.* 60.
Arbor altiſſima foliis oblongis nitidiſſimis nervoſis. *Brown. jam.* 372.
Mali perſicae Mammeae dictae folio arbor. *Sloan. cat.* 180.
Tſierou-ponna. *Rheed. mal.* 4. *p.* 81. *t.* 39.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 28.
Kleines Schönblatt. W.
Habitat in Indiis. ♄ (v. ſ.)
Ad Polygamiam Monoeciam. Iacq.
Poſſideo tantum americanam varietatem, aſiatica eſt forte diverſa. W.

1027. SPARRMANNIA.

Gen. plant. ed. Schreb. n. 893.

Cal. 4-phyllus. *Cor.* 4-petala reflexa. *Nect.* plura toruloſa. *Capſ.* angulata 5-locularis echinata.

1. SPARRMANNIA *africana.*
S. africana. *Linn. ſuppl.* 265. *Thunb. Nov. pl. gen.* 5. *p.* 89.* *Retz. obſ.* 5. *p.* 25. *n.* 65. *t.* 3. (*ſynonymum, patria et deſcriptio vero ad Lageſtroemiam pertinent.*)
Africaniſche Sparrmannie. W.
Habitat in Cap. b. ſpei *ſylvis* Eſſeboſch, Houtniquas, *inque lateribus montium ad* Lange Kloof. ♄

Frutex *ramis erectis alternis piloſis teretibus.* Folia *alterna longe petiolata dependentia cordata ovata ſublobata ſerrata acuminata novemnervia utrinque hiſpida.* Petioli *teretes piloſi.* Stipulae *laterales ſubulatae piloſae erectae.* Pedunculi *oppoſitifolii petiolis longiores erecti piloſi terminati in umbellam more Geraniorum, involucro polyphyllo ſubulato breviſſimo.* Pedicelli *piloſi.* Calyces *piloſi ſimpliciſſimi non ariſtati.* Petala *arcte reflexa flava.* Nectaria *lutea.* Stamina *atropurpurea.* Piſtillium *ſtaminibus multo longius luteum.* Capſula *echinata fuſca.* Semina *nigra.*

Flores ante floreſcentiam pedunculis deflexis, ſub flo-

reſcentia eriguntur pedunculi, at flores nutant, peracta floreſcentia eriguntur flores pro germine maturando, eodem modo ac in Dodecatheo.

Herba ſimilis Triumfettis ſ. Commerſoniae exactius, pedunculus et infloreſcentia Geraniorum, fructus Hibiſci, at aculeatus ut in Geis, inde quaſi a pluribus generibus compoſita.

1028. TILIA.

Gen. plant. edit. Schreb. n. 894.

Cor. 5-petala. *Cal.* 5-partitus. *Capſula* coriacea globoſa, 5-locularis, 5-valvis, baſi dehiſcens, monoſperma.

1. TILIA *europaea.*

T. floribus nectario deſtitutis. *Mat. med. p.* 136. *Gmel. ſib.* 4. *p.* 179. *Crantz. auſtr. p.* 96. *Pollich. pal. n.* 510. *Ludw. ect. t.* 48. *Fl. dan. t.* 553. *Kniph. cent.* 10. *n.* 87. *Knorr. del.* 1. *t. L.* 11. *Roth. germ. I.* 229. *II.* 587. *Willd. arb.* 387.

Tilia. *Hort. cliff.* 204. *Fl. ſuec.* 432, 471. *Phil. bot.* 259. * *Roy. lugdb.* 477. *Dalib. pariſ.* 153.

T. foliis cordato-lanceolatis, ſpongiolis ad nervorum angulos ſeſſilibus. *Hall. helv. n.* 1030.

α. T. (*grandifolia*) foliis cordatis acuminatis ſerratis ſubhirſutis, fructu quadriloculari. *Hoffm. germ.* 184.

T. foliis acuminatis ſerratis ſubhirſutis, fructibus quadrangularibus ſubpiloſis. *Mill. dict. n.* 2. *Du Roi. harbk.* 2. *p.* 461.

T. platyphyllos. *Scop. carn. n.* 641.

T. femina, folio majore. *Bauh. pin.* 426. *Blackw. t.* 469.

T. urbana. *Geſn. tab. III.*

β. T. montana, maximo folio. *Bauh. pin.* 426.

γ. T. femina, folio minore. *Bauh. pin.* 326.

T. (*parvifolia*) foliis cordatis acuminatis inaequaliter ſerratis, fructu quinqueloculari. *Hoffm. germ.* 185.

T. (*cordata*) foliis cordatis acuminatis inaequaliter ſerratis, fructibus quinquelocularibus tomentoſis. *Mill. dict. n.* 1. *Du Roi. harbk.* 2. *p.* 466.

T. ulmi folia. *Scop. carn. n.* 642.

δ. T. ulmi folio, ſemine hexagono. *Raj. angl.* 3. *p.* 473.

ε. T. foliis molliter hirſutis, viminibus rubris, fructu tetragono. *Raj. angl.* 3. *p.* 473. *Du Ham. arb.* 4.

ζ. T. bohemica, foliis minoribus glabris, fructu oblongo utrinque acuminato minimo coſtulato. *Till. piſ.* 165. *t.* 49 *f.* 9.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 31.
Gemeine Linde. W.
Habitat in Europae *pratis.* ♄ (v. v.)

2. TILIA *americana.*

T. floribus nectario inſtructis, foliis profunde cordatis argute ſerratis glabris. *Ait. Kew.* 2. *p.* 229.
T. floribus nectario inſtructis. *Sp. pl.* 733. *Mill. dict. n.* 3.
Tilia caroliniana. *Wangenh. amer.* 56. *Du Roi. harbk.* 2. *p.* 469.
T. foliis majoribus mucronatis. *Gron. virg.* 58.
T. ampliſſimis glabris foliis, noſtrati ſimilis. *Pluk. mant.* 181.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 35.
Schwarze Linde. W.
Habitat in Virginia, Canada. ♄ (v. v.)

*3. TILIA *pubeſcens.*

T. floribus nectario inſtructis, foliis baſi truncatis obliquis denticulato - ſerratis ſubtus pubeſcentibus. *Ait. Kew.* 2. *p.* 229.
Tilia caroliniana. *Mill. dict. n.* 4.
Tilia americana. *Wangenh. amer.* 55.
Feinbehaarte Linde. W.
Habitat in Carolina. ♄ (v. ſ.)

*4. TILIA *alba.*

T. foliis profunde cordatis ſubſinuatis dentatis, ſubtus tomentoſis. *Ait. Kew.* 2. *p.* 230. *Willd. arb.* 389.* *Waldſtein et Kitaibel pl. hung. t.* 3.
T. (*tomentoſa*) foliis cordatis acutis ſerratis, ſubtus tomentoſis. *Moench. weiſſenſt.* 136.
Tilia americana. *Du Roi. harbk.* 2. *p.* 467.
Weiſse Linde. W.
Habitat in Hungária. ♄ (v. v.)

Folia *ſubtus nivea.* Flores *ut in T. americana ſed minores. Clariſſ. Aiton, patriam hujus Americam eſſe, dixit, ſed a nullo peregrinatore indicata eſt. Amicus meus Clariſſ.* Kitaibel *ſylvas hujus ſpeciei in Hun-*

garia nuper detexit, et mecum ſpecimina communicavit, iis examuſſim ſimillima, quae ex hortis noſtris accepi; hinc de origine americana hujus valde dubito. W.

1029. LAETIA.

Gen. plant. edit. Schreb. n. 895.

Cor. 5-petala ſ. o. *Cal.* 5-phyllus. *Fructus* 1-locularis, trigonus. *Semina* arillo pulpoſo.

1. LAETIA *apetala.*

L. floribus apetalis, pedunculis ſubtrifloris axillaribus, foliis oblongis obtuſis ſerrulatis nitidis. W.

L. floribus apetalis. *Sp. pl.* 733. *Iacq. amer.* 167. *t.* 108. * *Swartz. obſ.* 219.

Guidonia Laetia. *Loefl. it.* 190.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 37.

Dreyblütige Lätie. W.

Habitat in America. ♄

*2. LAETIA *Guidonia.*

L. floribus apetalis, pedunculis unifloris terminalibus, foliis oblongis acuminatis ſerratis pubeſcentibus. *Swartz. prodr.* 83.

Guidonia foliis ovatis utrinque porrectis alternis quandoque crenatis, racemis laxis alaribus. *Brown. jam.* 249.

Einblütige Lätie. W.

Habitat in Jamaica. ♄

*3. LAETIA *Thamnia.*

L. floribus apetalis, pedunculis multifloris ſubdiviſis axillaribus, foliis oblongis acutis ſubcrenatis nitidis. *Swartz prodr.* 83.

Thamnia foliis ovatis leviſſime crenatis laete virentibus nitidis alternis, petiolis brevibus, pedunculis geniculatis. *Brown. jam.* 245.

Vielblütige Lätie. W.

Habitat in Jamaica. ♄

4. LAETIA *completa.*

L. floribus corollatis, foliis ovatis obtuſis ſerratis rugoſis glabris. W.

L. floribus petaloideis completis. *Sp. pl.* 733. *Iacq. amer.* 167. *t.* 183. *f.* 60.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 38.
Vollkommene Lätie. W.
Habitat in America. ♄

Laetiae apetalae Ludiis ſunt affines, ſed differunt fructus ſtructura. W.

*1030. RYANIA.

Vahl. eclog. 1. *p.* 51.

Cal. 5-phyllus, perſiſtens, coloratus. *Cor.* o. *Stigmata* 4. *Bacca* ſuberoſa unilocularis polyſperma.

*1. RYANIA *ſpecioſa.*

R. ſpecioſa. *Vahl. eclog.* 1. *p.* 51. *t.* 9. *
Prächtige Ryanie. W.
Habitat in inſula Trinitatis. ♄

Arbor *ramis ſubcineraſcentibus.* Folia *alterna petiolata oblonga acuminata integerrima venoſa glabra, coſta media ſubtus pulverulenta.* Pedunculi *uniflori axillares.* Flores *ſpecioſi.* W.

1031. GREWIA.

Gen. pl. ed. Schreb. n. 896.

Cal. 5-phyllus. *Cor.* 5-petala. *Nectar.* ſquamae 5. *Drupa* quadriloba 4-locularis. *Nuce* 1-2-ſperma.

1. GREWIA *occidentalis.*

G. foliis ſubrotundo-ovatis obtuſis dentatis glabris, pedunculis ſolitariis unifloris. W.
G. foliis ſubovatis, floribus ſolitariis. *Sp. pl.* 1367. *Mill. dict. n.* 1. *Willich. obſ. n.* 94. *Kniph. cent.* 9. *n.* 43.
G. corollis acutis. *Hort. cliff.* 433. *Du Ham. arb.* 1. *p.* 276. *t.* 108.
Ulmifolia arbor africana baccifera, floribus purpureis. *Pluk. alm.* 393. *t.* 237. 1.
Ulmi facie arbuſcula aethiopica, ramulis alatis, floribus purpuraſcentibus. *Comm. hort.* 1. *p.* 165. *t.* 85. *Seb. theſ.* 1. *p.* 46. *t.* 26. *f.* 3. *Raj. dendr.* 13.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 261.

Africaniſche Grewie. W.
Habitat ad Cap. b. ſpei. ♄ (v. v.)

*2. GREWIA *populifolia.*

G. foliis orbiculatis, pedunculis ſolitariis unifloris. *Vahl. ſymb.* 1. *p.* 33. *
Chadara tenax. *Forſk. deſcr. p.* 105. *n.* 23. *
Pappelblättrige Grewie. W.
Habitat in Arabia *felici.* ♄

Affinis quam maxime G. occidentali, habitus *vero laxior*, folia *orbiculata Populi tremulae dependentia*, pili *ſtellati ſubtus*. Vahl. l. c.

3. GREWIA *orientalis.*

G. foliis ovatis crenatis utrinque ſcaberrimis, pedunculis axillaribus trifloris. *Vahl. ſymb.* 1. *p.* 34.
G. foliis ſublanceolatis, floribus ſolitariis. *Sp. pl.* 1367. *Fl. zeyl.* 324. *
G. corollis obtuſis. *Roy. lugdb.* 467. *Hort. cliff.* 433.
Pai-paraea. *Rheed. mal.* 5. *p.* 91. *t.* 46. *Raj. hiſt.* 1624.
Frutex baccifer malabaricus, fructu plano rotundo piloſo tetragono. *Pluk. alm.* 275. *t.* 50. *f.* 4.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 262.
Oſtindiſche Grewie. W.
Habitat in India *orientali.* ♄ (v. ſ.)

Folia *obtuſiuſcula baſi utrinque ſaepe triglanduloſa, crenata, ſubtus punctis numeroſiſſimis, quibus inſident pili ſtellati.* Petioli, Pedunculi, Rami *ſuperne*, Calycesque *intenſe piloſi.* Vahl. l. c.

Poſſideo varietatem vel ſpeciem ex India orientali foliis *duplo longioribus quinque uncialibus, quae praeter magnitudinem a vulgari differt:* foliis *baſi cordatis*, ſtipulis *lineari-ſubulatis petiolo multo longioribus, denique* petalis *linearibus fere longitudine calycis.* W.

4. GREWIA *Mallococca.*

G. foliis cordatis ovato-oblongis crenatis ſcabris, pedicellis axillaribus trifloris, fructu tetracocco. *Lin. ſuppl.* 409. *Forſt. prod. n.* 327.
Mallococca crenata. *Forſt. gen. n.* 39. *Forſt. Nov. Act. Upſ.* 3. *p.* 179. *
Rauhfrüchtige Grewie. W.
Habitat in lucis inſulae Tongatabu *atque in montoſis inſulae* Huaheine. ♄

Difficile differentia eruitur, qua a praecedente dignoſcatur.

*5. GREWIA *laevigata.*

G. foliis ellipticis acuminatis utrinque laevibus baſi integerrimis, pedunculis trifloris. *Vahl. ſymb.* 1. *p.* 34.*

Glatte Grewie. W.

Habitat in India *orientali.* ♄ (v. ſ.)

Differt ab inſequente: foliis *longioribus baſi integerrimis eglanduloſis,* pedunculis *trifloris longis.* Vahl. l. c.

*6. GREWIA *glanduloſa.*

G. foliis ovato-lanceolatis acuminatis utrinque laevibus, baſi glanduloſis, floribus ſolitariis ſubſeſſilibus. *Vahl. ſymb.* 1. *p.* 34.

Drüſige Grewie. W.

Habitat in inſula Mauritii. ♄

Rami *ſcabriuſculi.* Folia *ovato-lanceolata acuminata acuta glabra utrinque laevia crenata, crenis* 3-5, *baſi proximis glanduloſis.* Petiolus *brevis.* Flores *axillares ſolitarii breviſſimi pedunculati.* Fructificatio *ut in reliquis.* Vahl. l. c.

*7. GREWIA *hirſuta.*

G. foliis lanceolato-ovatis mollibus, calycibus piloſiſſimis, pedunculis trifloris. *Vahl. ſymb.* 3. *p.* 34.*

Rauhe Grewie. W.

Habitat in ſummitate montium Indiae *orientalis.* ♄ (v. ſ.)

Facili negotio diſtinguitur a reliquis, foliis *molliſſimis lanceolato-ovatis.* Vahl. l. c.

*8. GREWIA *excelſa.*

G. foliis oblongis ſubtus tomentoſis incanis. *Vahl. ſymb.* 1. *p.* 35.

Chadara (*arborea*) foliis ovatis ſerratis glabris. *Forſk. deſcr.* 105.*

Hohe Grewie. W.

Habitat in Arabia *felici.* ♄

Folia *inaequaliter obtuſe ſerrata, ſupra laevia nervoſo-venoſa, ſubtus incana.* Pedunculi *axillares ſaepe ſolitarii triflori pedicellati angulati.* Vahl. l. c.

9. GREWIA *aſiatica.*

G. foliis cordato-ſubrotundis ſubtus canis, pedunculis axillaribus ſubquaternis petiolo longioribus. *Vahl. ſymb.* 1. *p.* 35.*

G. foliis cordatis. *Mant.* 122. *Sonnerat. itin.* 2. *p.* 191 *t.* 138. *
Houttuyn Lin. Pfl. Syſt. 2. *p.* 263.
Aſiatiſche Grewie. W.
Habitat in Suratte. ♄

Arbor *ramis vix tomentoſis.* Stipulae *lanceolatae.* Folia *alterna petiolata cordata ſubrotunda acutiuſcula quinquenervia vix lobata ſubtus tomentoſa magnitudine Mali.* Petioli *teretes tomentoſi foliis quintuplo breviores.* Pedunculi *axillares ſaepius quaterni trifidi triflori foliis dimidio breviores.* Petala *calyce non majora.* Baccae *parvae acidae.*

10. GREWIA *tiliaefolia.*

G. foliis cordato-ſubrotundis utrinque glabris, pedunculis petiolo brevioribus. *Vahl. ſymb.* 1. *p.* 35. *
Microcos (*lateriflora*) pedunculis axillaribus confertis dichotomis. *Sp. pl.* 734. *Fl. zeyl.* 208. *
Hamdamanias. *Herm. zeyl.* 3.
Lindenblättrige Grewie. W.
Habitat in India *orientali.* ♄

Similis G. aſiaticae, diverſa vero: foliis *duplo majoribus obtuſe ſerratis,* petiolis *vix incraſſatis,* ſtipulis *ſemicordatis nec ſubulatis,* pedunculis *petiolo dimidio brevioribus, minime vero duplo longioribus,* floribus *duplo minoribus.* Vahl. l. c.

*11. GREWIA *velutina.*

G. foliis ovalibus utrinque molliſſimis ſubtus incanis, pedunculis axillaribus ſubternis trifloris. *Vahl. ſymb.* 1. *p.* 35. *
Chadara (*velutina*) foliis alternis ovato-acutis ſerratis velutinis baſi obliquis. *Forſk. deſcr.* 106. *
Weichblättrige Grewie. W.
Habitat in Arabia *felici.* ♄

A G. hirſuta differt: ramis foliis*que minime piloſis, ſed tomento tenuiſſimo albidis, imprimis ſubtus, demum* ſerraturis *minutioribus. A G. Mallocca differt:* foliis *molliſſimis.* Vahl. l. c.

*1032. MAERUA.

Vahl. ſymb. 1. *p.* 36.

Cal. 4-fidus tubo nectarifero. *Cor.* o. *Stigma* ſeſſile. *Drupa*?

*1. MAERUA *uniflora.*

M. pedunculis axillaribus terminalibusque unifloris, nectariis diviſis. *Vahl. ſymb.* 1. *p.* 36. *
Maerua craſſifolia. *Forſk. deſcr.* 104. *
Einblumige Märua. W.
Habitat in Arabia *felici.*

*2. MAERUA *racemoſa.*

M. floribus racemoſis, nectariis integris, foliis retuſis. *Vahl. ſymb.* 1. *p.* 36. *
Traubentragende Märua. W.
Habitat in Arabia *felici.* ♄

Folia *minus craſſa quam in priori.* Habitus, Calyx, Germen, Stylus *et* Stigma *ut in praecedente ideoque hujus generis, quamvis nectario integro gaudet.* Vahl l. c.

1033. MICROCOS.

Gen. pl. ed. Schreb. n. 897.

Cal. 5-phyllus. *Cor.* 5-petala. *Nectar.* o. *Drupa* nuce triloculari.

1. MICROCOS *paniculata.*

M. panicula terminali. *Sp. pl.* 733. *Fl. zeyl.* 207. *
Grewia (*Microcos*) foliis ovato-oblongis, floribus paniculatis. *Syſt. veg.* 689.
Microcos foliis alternis oblongis acuminatis. *Burm. zeyl.* 159. *t.* 74.
Schageri Cottam. *Rheed. mal.* 1. *p.* 105. *t.* 56.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 264.
Riſpentragende Microcos. W.
Habitat in India *orientali.* ♄ (v. ſ.)

Rami *teretes glabri, juniores tomentoſi.* Folia *alterna oblonga acuminata, apice ſerrata, baſi obliqua, trinervia venoſa, venis partialibus horizontalibus, ſubtus tomentoſa.* Panicula *terminalis bracteata.* W.

*1034. MARILA.

Gen. plant. edit. Schreb. n. 1756.

Cor. 5-phyllus. *Cor.* 5-petala. *Stigma* ſimplex. *Capſ.* 4-locularis polyſperma.

*1. MARILA *racemoſa.*
M. racemoſa. *Swartz. prodr.* 84.
Traubige Marile. W.
Habitat in inſulis Montſerrat *et* Chriſtopher Americes. ♄

1035. ELAEOCARPUS.

Gen. pl. ed. Schreb. n. 898.

Cor. 5-petala, lacera. *Anthera* apice 2-valvi. *Cal.* 5-phyllus. *Drupa* nucleo criſpo.

1. ELAEOCARPUS *ſerratus.*
E. foliis alternis lanceolato-ellipticis ſerratis, racemis axillaribus. *Vahl. ſymb.* 3. *p.* 67.
E. ſerrata. *Sp. pl.* 734. *Retz. obſ.* 4. *p.* 27. *
Elaeocarpus. *Fl. zeyl.* 206.
Elaeocarpus foliis lauri ſerratis, floribus ſpicatis. *Burm. zeyl.* 39. *t.* 40.
Ganitrus. *Rumph. amb.* 3. *p.* 60. *t.* 101.
Perin-Kara. *Rheed. mal.* 4. *p.* 51. *t.* 24. *Raj. hiſt.* 1546. *dendr.* 42.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 40.
Geſägter Ganiterbaum. W.
Habitat in India. ♄ (v. ſ.)

Rumphius *plures ſpecies commemorat.* Stamina *in Zeylanica* 20, *in Javanica* 8, *in Burmanniana* 16, 20 *a me viſa. In nupero ſpecimine vidi vel aliam ſpeciem vel varietatem, cui*

Calyx 5-*partitus, acutus.* Petala 3, *unguiculata, calyce longiora, lacera ramentis capillaribus.* Filamenta 8, *brevia.* Antherae *columnares, dehiſcentes apice bivalvi.* Germen *ſuperum, ſubgloboſum.* Stylus *filiformis, longitudine ſtaminum.* Stigma *ſimplex.*

Rami *teretes cicatriſati glabri.* Folia *petiolata tripollicaria obtuſe ſerrata glabra nervoſo-venoſa.* Racemi *axillares.* Flores *diſtantes.* Vahl l. c.

2. ELAEOCARPUS *dentatus.*
E. foliis alternis oblongis ſuperne dentato-ſerratis, race-

mis axillaribus, floribus monogynis. *Vahl ſymb.* 3. *p.* 67.

E. (*ſerratus*) monogynus, foliis alternis oblongis ſerratis, racemis axillaribus. *Lin. ſuppl.* 266. (*excluſo ſynonymo Syſt. veg.*)

Dicera (*dentata*) monogyna, foliis oblongis acuminatis dentatis, racemis ſimplicibus axillaribus laxis. *Forſt. prod. n.* 226. *Forſt. gen.* 40. *n.* 1.

Gezähnter Ganiterbaum. W.

Habitat in Nova Zeelandia. ♄

Rami *teretes punctati glabri.* Ramuli *pubeſcentes.* Folia *petiolata alterna bipollicaria, ſuperne remote dentato-ſerrata baſi acuta, utrinque glabra, excepto in coſta, ſupra nervoſa ſubtus ſubvenoſa.* Petioli *villoſi.* Vahl l. c.

3. ELAEOCARPUS *Dicera.*

E. foliis oppoſitis ovatis duplicato-ſerratis, racemis compoſitis, floribus tetragynis. *Vahl. ſymb.* 3. *p.* 67. *

E. tetragynus, foliis oppoſitis ovatis duplicato-ſerratis, racemis lateralibus oppoſitis. *Lin. ſuppl.* 266.

Dicera (*ſerrata*) foliis oppoſitis cordato-ovatis inaequaliter ſerratis, racemis lateralibus compoſitis. *Forſt. prod. n.* 227. *Forſt. gen.* 40. *n.* 2.

Viergriffliger Ganiterbaum. W.

Habitat in Nova Zeelandia. ♄

Structura *floris convenit cum E. dentato, at differt:* ſtylis *quatuor,* fructu *baccato quadriloculari: loculis diſpermis.*

*4. ELAEOCARPUS *integrifolius.*†

E. foliis alternis ovato-oblongis obtuſis integerrimis, racemis axillaribus ſimplicibus, floribus quadrifidis. *Lamarck. encycl.* 2. *p.* 597. *

Ganitrum oblongum. *Rumph. amb.* 3. *p.* 163. *t.* 102. *

Ungezähnter Ganiterbaum. W.

Habitat in inſula Mauritii. ♄

5. ELAEOCARPUS *copalliferus.*

E. foliis integerrimis, panicula terminali. *Vahl ſymb.* 3. *p.* 67. *

E. (*copalliferus*) foliis integris. *Retz. obſ.* 4. *p.* 27. *

Vateria indica. *Sp. pl.* 734. *Fl. zeyl.* 204.

Paenoe. *Rheed. mal.* 4. *p.* 33. *t.* 15.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 48.

Copaltragender Ganiterbaum. W.
Habitat in Zeylona. ♄

Calycem *non ſuperum invenio, ut habet Celeb. Retzius, tamen eandem eſſe cum noſtra, omnino perſuaſus ſum.* Vahl l. c.

*1036. LEGNOTIS.

Gen. plant. edit. Schreb. n. 900.

Cal. 5-fidus. *Cor.* 5-petala receptaculo inſerta lacera. *Capſ.* 3-locularis.

*1. LEGNOTIS *elliptica.*

L. foliis ellipticis, floribus pedicellatis. *Swartz. prodr.* 84.
Elliptiſche Saumblume. W.
Habitat in Jamaica. ♄ (v. ſ.)

Folia *oblonga acuminata integerrima*, Flores *bini vel ſolitarii axillares pedunculati.* Pedunculi *uniflori.* W.

*2. LEGNOTIS *Caſſipourea.*

L. foliis ovatis, floribus ſeſſilibus. *Swartz. prod.* 84.
Caſſipourea gujanenſis. *Aubl. guj.* 1. *p.* 529. *t.* 211. *
Sitzende Saumblume. W.
Habitat in locis paludoſis Gujanae. ♄

*1037. MYRODENDRUM.

Gen. plant. edit. Schreb. n. 901.

Cal. 5-dentatus. *Cor.* 5-petala. *Stigma* 5-lobum. *Pericarp.* 5-loculare, loculis monoſpermis.

*1. MYRODENDRUM *amplexicaule.* W.

Houmiri balſamifera. *Aubl. guj. p.* 564. *t.* 225. *
Stielumfaſſender Balſambaum. W.
Habitat in Cajenna, Gujana. ♄

Arbor *procera.* Folia *alterna lanceolata acuta, ſeſſilia amplexicaulia crenata venoſa utrinque glabra.* Flores *terminales albi parvi corymboſi.* W.

*1038. LEMNESCIA.

Gen. pl. ed. Schreb. n. 902.

Cal. 5-dentatus. *Cor.* 5-petala. *Nectarium* cyathiforme ſtaminiferum. *Capſula?* 5-locularis, loculis 1-ſpermis.

*1. LEMNESCIA *floribunda.* W.
Vantanea gujanenſis. *Aubl. guj.* 1. *p.* 572. *t.* 229. *
Vielblütige Lemneſcie. W.
Habitat in Gujana. ♄
Folia *alterna oblonga acuminata integerrima venoſa.* Flores *coccinei ſpecioſi, in corymbo multifloro terminali.* W.

*1039. ASCIUM.

Gen. plant. ed. Schreb. n. 903.

Cal. 5-phyllus. *Cor.* 5-petala. *Bacca* 4-locularis, loculis diſpermis.

*1. ASCIUM *violaceum.*
A. violaceum. *Vahl eclog. p.* 41. *
Norantea gujanenſis. *Aubl. guj.* 1. *p.* 554. *t.* 220. *
Violette Becherblume. W.
Habitat in ſylvis paludoſis Gujanae *et in inſula* Trinitatis. ♄
Arbor *altiſſima* 80-*pedalis.* Folia *alterna obovata, breve petiolata, coriacea, reticulato-venoſa, integerrima, emarginata mucronata glabra.* Racemus *terminalis ſecundus.* Flores *parvi violacei,* Bractea *pedicellata aſcidiformi coccinea magna, ſuffulti.* W.

1040. LECYTHIS.

Gen. plant. ed. Schreb. n. 904.

Cor. 6-petala. *Cal.* 6-phyllus. *Nectarium* lingulatum, ſtaminiferum. *Pericarp.* circumſciſſum, polyſpermum.

1. LECYTHIS *Ollaria,*
L. foliis ſeſſilibus cordato-ovatis ſubintegerrimis. *Syſt. veg.* 410. *Loefl. it.* 189.
Jacapucaya. *Margr. braſ.* 128. *Piſ. braſ.* 135.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 43.
Herzblättriger Topfbaum. W.
Habitat in Braſilia, Cumana. ♄
Folia *alterna, ſeſſilia. Flos Fructusque ſingulares.*
Natura *fructu ollas producit, ut in Cucurbita et Creſcentia lagenas.*

2. LECYTHIS *minor.*
L. foliis petiolatis lanceolatis ſerratis. *Syſt. veg.* 410. *Iacq. amer.* 168. *t.* 109.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 45.
Sägeblättriger Topfbaum. W.
Habitat Carthagenae Americes *in ſylvis.* ♄

*3. LECYTHIS *grandiflora.*
L. foliis petiolatis oblongis acutis integerrimis, racemis axillaribus terminalibusque petiolo multoties longioribus petalis obtuſis. W.
L. (*grandiflora*) foliis ovatis, florum pedunculis craſſis. *Aubl. guj.* 2. *p.* 712. *t.* 283. *
Groſsblütiger Topfbaum. W.
Habitat in ſylvis Gujanae. ♄

*4. LECYTHIS *amara.*
L. foliis petiolatis oblongis acuminatis integerrimis, racemis axillaribus terminalibusque petiolo duplo longioribus, petalis acutis. W.
L. (*amara*) foliis ovato-lanceolatis acuminatis, fructu parvo, nucleo amaro. *Aubl. guj.* 2. *p.* 716. *t.* 286. *
Bitterer Topfbaum. W.
Habitat in ſylvis Gujanae. ♄

*5. LECYTHIS *Zabucajo.*
L. foliis petiolatis lanceolatis acuminatis integerrimis, racemo terminali, petalis acutis. W.
L. (*Zabucajo*) foliis lanceolato-oblongis acuminatis, fructu magno, nucleo eduli. *Aubl. guj.* 2. *p.* 718. *t.* 288. (*excluſis ſynonymis.*)
Eſsbarer Topfbaum. W.
Habitat in ſylvis Gujanae. ♄

*6. LECYTHIS *Idatimon.*
L. foliis petiolatis oblongis acutis integerrimis, racemis axillaribus, pedunculis glanduloſis, petalis obtuſis. W.
L. (*Idatimon*) foliis ovato-lanceolatis acuminatis, fructu parvo quadriloculari. *Aubl. guj.* 2. *p.* 721. *t.* 289. *

β. L. (*lutea*) foliis lanceolatis oblongis acuminatis, flore parvo. *Aubl. guj.* 2. *p.* 721.
Drüſiger Topfbaum. W.
Habitat in ſylvis Gujanae. ♄

*7. LECYTHIS *parviflora.*
L. foliis petiolatis oblongis acuminatis integerrimis, racemis paniculatis terminalibus, petalis acutis. W.
L. (*parviflora*) foliis ovato-lanceolatis acuminatis, fructu parvo biloculari, operculo cum appendice lignoſa intus producta. *Aubl. guj.* 2. *p.* 717. *t.* 287. *
Kleinblütiger Topfbaum. W.
Habitat ad fluviorum ripas Gujanae. ♄

*8. LECYTHIS *bracteata.* W. †
L. foliis petiolatis elliptico-obovatis acuminatis integerrimis, racemis lateralibus, calyce bracteato, cortice fructus duplici. W.
Couroupita gujanenſis. *Aubl. guj.* 2. *p.* 708. *t.* 282. *
Beblätterter Topfbaum. W.
Habitat in Cajenna. ♄
Diſtincta a reliquis: Bracteis *binis ad baſin calycis, et* cortice *fructus duplici.* W.

*1041. ALANGIUM.
Juſſ. gen. pl. ed. Uſteri 358.

Cal. 6-f. 10-dentatus ſuperus. *Cor.* 6-f. 10-petala. *Bacca* corticoſa 2-3-ſperma.

1. ALANGIUM *decapetalum.*
A. floribus decapetalis, ramis ſpineſcentibus. *Vahl ſymb.* 2. *p.* 61.
A. (*decapetalum*) ſpinoſum, foliis alternis oblongis, baccis glabris ſubtriſpermis. *Lamarck. encycl.* 1. *p.* 173. *
Grewia (*ſalvifolia*) foliis oblongis integerrimis, floribus axillaribus pluribus pedicellatis, petalis recurvis linearibus. *Lin. ſuppl.* 409. *
Baccifera indica, fructu rotundo umbilicato, ceraſi magnitudine, dicocco. *Raj. hiſt.* 1497.
Angolam. *Rheed. mal.* 4. *p.* 39. *t.* 17.
Zehnblättriges Alangium. W.
Habitat in India *orientali.* ♄
Rami *teretes glabri ſparſi, ſuperne vix villoſi, apice in ſpinam deſinentes.* Folia *petiolata tripollicaria la-*

to-lanceolata membranacea venoſa, ſubtus pallidiora ſubtrinervia, ſecundum nervos rare piloſa, utrinque acuta integerrima. Pedunculi *axillares plerumque terni ſemiunguiculares tomentoſi, ſuperne ſub germine annulati articulati.* Calyces *et* Petala *extus tomentoſa.* Bacca *magnitudine nucis Coryli, lineis aliquot elevatis, farinoſo-ſubvilloſa.* Vahl l. c.

*2. ALANGIUM *hexapetalum.*

A. floribus hexapetalis, ramis inermibus. *Vahl ſymb.* 2. *p.* 62.

A. (*hexapetalum*) inerme, foliis ovato-lanceolatis alternis, baccis tomentoſis monoſpermis. *Lamarck encycl.* 1. *p.* 173.*

Prunifera indica, arbor fructu umbilicato corticoſo perſici ſimili. *Raj. hiſt.* 1483.

Arbor baccifera maderaſpatana. *Pluk. amalth.* 24. *t.* 370. *f.* 1?

Kara-Angolam. *Rheed. mal.* 4. *p.* 55. *t.* 26.

Sechsblättriges Alangium. W.

Habitat in India *orientali.* ♄

Differt a praecedente: foliis *latioribus in acumine longiori productis et notis indicatis, caeterum ſatis ſimile.* Vahl l. c.

1042. MENTZELIA.

Gen. pl. ed. Schreb. n. 907.

Cor. 5-petala. *Cal.* 5-phyllus. *Capſ.* infera, cylindrica, polyſperma.

1. MENTZELIA *aſpera.*

M. caule ramoſo, pedunculis axillaribus, petalis crenatis obtuſis. W.

Mentzelia. *Hort. cliff.* 492. *Amoen. acad.* 5. *p.* 398.* *Loefl. it.* 223. *Mill. dict. n.* 1.

M. foliis et fructibus aſperis. *Plum. gen.* 41. *ic.* 174. *f.* 1.

M. ſetis uncinatis munita, foliis lobatis, fructibus ſeſſilibus ſingularibus ad divaricationes ramorum. *Brown. jam.* 249.

Onagra americana, folio betonicae, fructu hiſpido. *Tournef. inſt.* 302.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 216.

Scharfe Mentzelie. W.
Habitat in America.

*2. MENTZELIA *hiſpida.* W.
M. caule dichotomo, pedunculis unifloris in caulis dichotomia, petalis integerrimis cuſpidatis. W.
Mentzelia (*aſpera*) foliis ovato-lanceolatis biſerratis inciſis, floribus ſolitariis ſeſſilibus. *Cavan. ic.* 1. *p.* 51. *t.* 70.* (*excluſis ſynonymis.*)
Rauhe Mentzelie. W.
Habitat in Mexico. ♃
Diverſiſſima a praecedente. W.

1043. LOASA.

Gen. plant. ed. Schreb. n. 908.

Cor. 5-petala. *Cal.* 5-phyllus. *Capſ.* ſemiinfera, 1-locularis, ſemitrivalvis, polyſperma.

1. LOASA *hiſpida.*
L. hiſpidiſſima, foliis alternis bipinnatifidis, laciniis calycinis margine revolutis, corolla reflexa. *Lamarck. encycl.* 3. *p.* 554. *
L. hiſpida. *Syſt. nat. ed.* 13. *p.* 364.
L. urens. *Iacq. obſ.* 2. *p.* 15. *t.* 38.
Loaſa. *Adanſ. fam.* 2. *p.* 501?
Houttuyn Lin. Pfl. Syſt. 7 *p.* 218. *t.* 53. *B.*
Brennende Loaſe. W.
Habitat in Peru. ☉
Petala *flava.* Nectaria *alba punctis rubris viridibusque.* Filamenta *in* 5 *faſciculos interſtitia nectariorum occupantia.* Stamina *primum erecta, demum reflexa. Affinitate* Mentzeliae *accedit.* Iacq.

*2. LOASA *contorta.*
L. ſcandens, foliis oppoſitis petiolatis ſubruncinatis, capſula oblonga contorta nutante. *Lamarck. encycl.* 3. *p.* 554. * *illuſt. gen. t.* 426. *f.* 2.
Gedrehte Loaſe. W.
Habitat in Peru.

*3. LOASA *acanthifolia.*
L. foliis oppoſitis pinnatifidis, ſuperioribus ſeſſilibus, calyce reflexo, petalis apice bidentatis. *Lamarck. encycl.* 3. *p.* 555. *

Ortiga chilienſis urens Acanthi folio. *Feuill. peruv.* 2. *p.* 757. *t.* 43.
Bärenklaublättrige Loaſe. W.
Habitat in Chili.

*4. LOASA *grandiflora.*

L. hiſpida, foliis oppoſitis alternisque cordato-ovatis lobatis, petalis planiuſculis, flore maximo. *Lamarck. encycl.* 3. *p.* 555. *
Groſsblütige Loaſe. W.
Habitat in Peru.

*5. LOASA *chenopodifolia.*

L. foliis ſubovatis inciſo-dentatis, racemis laxis folioſis terminalibus, floribus cernuis. *Lamarck. encycl.* 3. *p.* 556. *
Gänſefuſsblättrige Loaſe. W.
Habitat in Peru.

*6. LOASA *nitida.*

L. proſtrata, foliis cordato-lobatis dentatis, ſupra nitidis, ſuperioribus ſeſſilibus, pedunculis axillaribus. *Lamarck. encycl.* 3. *p.* 556. *
Glänzende Loaſe. W.
Habitat in rupibus Limae. ⊙

*1044. STERBECKIA.

Gen. plant. edit. Schreb. n. 909.

Cal. 3-ſ. 5-phyllus. *Cor.* 3-ſ. 5-petala. *Capſula* corticoſa non dehiſcens leguminiformis polyſperma. *Semina* imbricata in pulpa nidulantia.

*1. STERBECKIA *lateriflora.* W.

Singana gujanenſis. *Aubl. guj.* 1. *p.* 574. *t.* 230. *
Seitenblütige Sterbeckie. W.
Habitat in ſylvis Gujanae. ♄

Frutex *ſcandens.* Folia *ſuboppoſita petiolata elliptica acuminata integerrima venoſa glabra.* Pedunculi *multiflori breviſſimi laterales.* Flores *albi parvi.* W.

1045. LAGERSTROEMIA.

Gen. plant. edit. Schreb. n. 910.

Cor. 6-petala. *Cal.* 6-fidus, campanulatus. *Stam.* multa: horum 6 exteriora craſſiora. *Capſ.* 4-ſ. 6-locularis polyſperma.

1. LAGERSTROEMIA *indica.*

L. petalis criſpis, panicula terminali, foliis ſubrotundo-ovatis acutis glabris. W.

L. foliis ovatis, calycibus nudis laevibus. *Retz. obſ.* 5. *p.* 25. *et p.* 20. *

L. indica. *Sp. pl.* 734. *Thunb. jap.* 224. * *Loureiro cochin.* 415. * *Curt. mag.* 405.

Tſiin-kin. *Rumph. amb.* 7. *p.* 61. *t.* 28.

Sibi. *Kaempf. amoen.* 855.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 18.

Chineſiſche Lagerſtrömie. W.

Habitat in China, Cochinchina, Japonia. ♄ (v. ſ.)

Arbor *Punicae magnitudinis.* Folia *oppoſita. interdum alterna, ſubſeſſilia, oblonga, integerrima, glabra: floralia ſubrotunda.* Thyrſus *terminalis, laxus, pedicellis trifidis ſ. trifloris.* Flores *incarnati: Petalis longe unguiculatis,* 6, *meſenterii inſtar criſpis.*

*2. LAGERSTROEMIA *Reginae.*

L. petalis undulatis, panicula terminali, foliis oblongis glabris. W.

Lagerſtroemia Reginae. *Roxb. corom.* 1. *p.* 46. *t.* 65. *

L. (*Flos Reginae*) foliis ovato-oblongis, calycibus rugoſis tomentoſis. *Retz. obſ.* 5. *p.* 25. * *et* 1. *p.* 20. *

Adambea glabra. *Lamarck. encycl.* 1. *p.* 39.

Alcea indica arborea, pericarpio carnoſo in plura loculamenta partito. *Raj. hiſt.* 1902.

Adamboe. *Rheed. mal.* 4. *p.* 45. *t.* 20. 21.

Prächtige Lagerſtrömie. W.

Habitat in ſylvis Calcuttae, Javae. ♄

*3. LAGERSTROEMIA *hirſuta.* W.

L. petalis planis, panicula corymboſa terminali, foliis oblongis pubeſcentibus. W.

Adambea hirſuta. *Lamarck. encycl.* 1. *p.* 39.

Alcea indica arborea elatior, pericarpio craſſo ſubaſpero. *Raj. hiſt.* 1902.

Katou-Adamboe. *Rheed. mal.* 4. *p.* 47. *t.* 22.
Behaarte Lagerſtrömie. W.
Habitat in montoſis Malae *et* Poigae *provinciis* Malabariae. ♄

4. LAGERSTROEMIA *Münchhauſia.* W.

L. petalis planis, racemo ſubcompoſito terminali, foliis ovatis glabris. W.
Münchhauſia ſpecioſa. *Mant.* 243. *Murr. prod. in praef. c. ic. Münchh. hausv.* 5. *p.* 357. *t.* 356.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 303.
Schöne Lagerſtrömie. W.
Habitat in China. ♄

Arbuſcula *ramis teretibus laevibus.* Folia *alterna petiolata ovata ſeu ovato-oblonga acuminata integerrima glabra ſubtus pallidiora.* Racemus *terminalis compoſitus erectus.*

*5. LAGERSTROEMIA *parviflora.*

L. petalis planis, pedunculis axillaribus trifloris, foliis oblongis obtuſis ſcabris. W.
Lagerſtroemia parviflora. *Roxb. corom.* 1. *p.* 48. *t.* 66.*
Kleinblütige Lagerſtrömie. W.
Habitat in montoſis Indiae. ♄

*1046. FREZIERA.

Eroteum. Gen. pl. ed. Schreb. n. 1757.

Cal. 5-phyllus. *Cor.* 5-petala. *Stylus* trifidus. *Bacca* exſucca 3-locularis polyſperma.

*1. FREZIERA *Theaeoides.*

F. foliis ovato-lanceolatis ſerrato-dentatis, floribus axillaribus ſolitariis.
Eroteum Theaeoides. *Swartz. prod.* 85.
Theeartige Freziere. W.
Habitat in Jamaica. ♄

*2. FREZIERA *undulata.*

F. foliis elliptico-lanceolatis acuminatis ſerratis, floribus confertis axillaribus.
Eroteum undulatum. *Swartz. prod.* 85. *Vahl ſymb.* 2. *p.* 61.*
Wellenblättrige Freziere. W.

Habitat in India *occidentali.* ♄ (v. ſ.)

Ob conformem denominationem generis Erodii in claſſe Monadelphia, mutavit amiciſſimus mihi Clariſſ. Swartz nomen, in litteris ad me datis. W.

1047. THEA.

Gen. pl. ed. Schreb. n. 911.

Cor. 6-ſ. 9-petala. *Cal.* 5-ſ. 6-phyllus. *Capſ.* 3-cocca.

1. THEA *Bohea.*

T. floribus hexapetalis. *Sp. pl.* 734. *Hort. cliff.* 204. *Mat. med.* 136. *Amoen. acad.* 7. *p.* 239. *t.* 4. *Hill. exot. t.* 22. *Blackw. t.* 352. *Thunb. jap.* 225.

α. *laxa*, foliis elliptico-oblongis rugoſis. *Ait. Kew.* 2. *p.* 230.

Thee. *Kaempf. jap.* 605. *t.* 606.

Thee frutex. *Bart. act.* 4. *p.* 1. *t.* 1. *Bont. jav.* 87 *t.* 88. *Barr. rar.* 128. *t.* 904. *Letſom. diſſ. lugdb.* 1769. *t.* 1. *f.* 1. 2.

Thee ſinenſium. *Breyn. cent.* 111. *t.* 112. *ic.* 17. *t.* 3. *Bocc. muſ.* 114. *t.* 94.

Chaa. *Bauh. pin.* 147.

Evonymo affinis arbor orientalis nucifera, flore roſeo. *Pluk. alm.* 139. *t.* 88. *f.* 6.

β. *ſtricta*, foliis lanceolatis planis. *Ait. Kew.* 2. *p.* 231.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 19.

Brauner Thee. W.

Habitat in Japonia, China. ♄ (v. v.)

Folia *alterna, elliptica, conſiſtentia, laevia, obtuſiuſcula, obtuſe ſerrata: Petiolis brevibus, ſubtus teretibus, gibbis.* Stipulae *nullae.*

Nobis attulit vivam primus Car. Guſt. Ekeberg, navis ad Chinenſes gubernator, d. 3 *Octob.* 1763. *Obtinetur, ſi ſemina, ad navis receſſum e China, in nave terrae mandentur, ut in itinere ſerius germinent.*

Perfidi Chinenſes ſaepe pro Theae frutice offerunt Camelliam, *qua leviter verſatus absque floribus facile ſeducitur, non obſervatis attente petiolis. Multi oſtenderunt alium fruticem pro Thea, ſed, ni fallor, nullus vivam habuit in Europa ante me.*

2. THEA *viridis.*

T. floribus enneapetalis. *Sp. pl.* 735. *Hill. exot. t.* 22.

T. cantonienſis. *Loureiro cochin.* 414. *
T. ſinenſis. *Blackw. t.* 351. R.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 22.
Grüner Thee W.
Habitat in China. ♄
Folia hujus longiora, illius breviora.

Florum *libram integram aliquando accepi; eſſe Theam viridem ex Hillio didici.* Stylos 3 *conglutinatos eſſe, nec ſimplicem. Io. Coakly Letſom.*

1048. CISTUS.

Gen. pl. ed. Schreb. n. 913.

Cor. 5-petala. *Cal.* 5-phyllus: foliolis duobus minoribus. *Capſula.*

* *Exſtipulati fruticoſi.*

1. CISTUS *capenſis.*

C. arboreſcens exſtipulatus, foliis ovato-lanceolatis petiolatis trinerviis denticulatis utrinque nudis. *Sp. pl.* 736. *Vahl. ſymb.* 3. *p.* 68. *
Houttuyn Lin. Pfl. Syſt. 4. *p.* 23.
Capiſche Ciſtenroſe. W.
Habitat ad Cap. b. Spei. ♄
Differt a ſequentibus foliorum denticulis.

Rami *teretes, pilis rarioribus adſperſi purpuraſcentes.* Folia *tripollicaria remota, ſuperiora ſeſſilia, baſi non connata; inferiora baſi in petiolum breviſſimum attenuata acuta ſubtriplinervia, nervis verſus medium evaneſcentibus, utrinque glabra, margine denticulata, pilis longis ciliata, praeſertim ſuprema. Petioli diſtincti* Stipulae *nullae.* Pedunculi *terminales trifidi, pedunculis partialibus trifloris.* Calycis *foliola cordata acuminata hirſuta denticulata ciliata, juniora hirſutiſſima.* Corolla *flava.* Vahl. l. c.

2. CISTUS *villoſus.*

C. arboreſcens exſtipulatus, foliis ovatis petiolatis hirtis. *Syſt. veg.* 412. *Desfont. fl. atl.* 1. *p.* 408. *
C. piloſus. *Lin. ſpec. pl.* 2. *p.* 737. *Mill. dict. n.* 1.
C. mas, folio rotundo hirſutiſſimo. *Bauh. pin.* 464.
C. mas major, folio rotundiore. *I. B. Du Ham. arb.* 1. *p.* 167. *t.* 64.

C. mas matthioli. *Dalech. hiſt.* 222.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 23.
Rauhe Ciſtenroſe. W.
Habitat in Italia, Hiſpania *et* Africa *boreali* ♄.

Folia *petiolata, rugoſa, utrinque viridia et pubeſcentia.* Pedunculi *uniflori cum geniculo.*

3. CISTUS *populifolius.*

C. arboreſcens exſtipulatus, foliis cordatis laevibus acuminatis petiolatis. *Hort. cliff.* 205. *Roy. lugdb.* 474. *Mill. dict. n.* 15. *Cavan. ic.* 3. *p.* 8. *t.* 215. *
C. Ledon, foliis populi nigrae, major. *Bauh. pin.* 467.
Ledum latifolium 2. majus. *Cluſ. hiſt.* 1. *p.* 78.
β. Ciſtus Ledon, foliis populi nigrae, minor. *Bauh. pin.* 467.
Ledum latifolium 2. minus. *Cluſ. hiſt.* 1. *p.* 78.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 24.
Pappelblättrige Ciſtenroſe. W.
Habitat in Luſitania, Hiſpania. ♄ (v. v.)

Planta *ramis et foliis piloſa, etiam piloſiſſima calycibus; adultior ramis et foliis nudus eſt.* Petala *alba absque macula margine purpuraſcentia.* Stam. *lutea.* Calyces *ante explicationem triquetri.*

4. CISTUS *laurifolius.*

C. fruteſcens exſtipulatus, foliis ovato-lanceolatis petiolatis trinerviis ſupra glabris, ſubtus tomentoſis, petiolis baſi connatis, pedunculis nudis multifloris. *Lamarck. encycl.* 2. *p.* 16. *
C. arboreſcens exſtipulatus, foliis oblongo-ovatis petiolatis trinerviis ſupra glabris: petiolis baſi connatis. *Mill. dict. n.* 11.
Ciſtus Ledon, foliis laurinis. *Bauh. pin.* 467.
Ciſtus Ledon latifolium creticum. *Bauh. hiſt.* 2. *p.* 9.
Ledon latifolium. *Cluſ. hiſt.* 1. *p.* 77. *hiſp.* 156.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 25.
Lorbeerblättrige Ciſtenroſe. W.
Habitat in Hiſpania. ♄ (v. v.)

Caulis *arboreſcens ramis piloſis.* Folia *lanceolata trinervia, acuta, rugoſa, utrinque viridia, vix manifeſte piloſa.* Petioli *baſi purpuraſcentes, ſaepe diſtincti.* Flores *albi.*

Folia *ſubtus tomentoſa.* W.

*5. CISTUS *vaginatus.*

C. arboreſcens exſtipulatus, foliis oblongis piloſis ſubtus reticulato-rugoſis, petiolis baſi coalitis vaginantibus ſulcatis. *Ait. Kew.* 2. *p.* 232. *Iacq. hort. ſchoenb.* 3. *p.* 17. *t.* 282. *

C. (*ſymphytifolius*) fruticoſus exſtipulatus, foliis petiolatis oblongo-lanceolatis ſupra villoſis, petiolis baſi vaginantibus connatis. *Lamarck. encycl.* 2. *p.* 15. *

Scheidentragende Ciſtenroſe. W.

Habitat in inſula Teneriffa. ♄ (v. v.)

Folia *ſubtus trinervia reticulato-venoſa.* Petioli *connati.* Corolla *purpurea.* W.

*6. CISTUS *Ledon.*

C. fruticoſus exſtipulatus, foliis ſubſeſſilibus lanceolatis nervoſis connatis ſupra glabris, floribus corymboſis erectis, pedunculis calycibusque villoſo-ſericeis. *Lamarck. encyl.* 1. *p.* 17. *

C. ladanifera Monſpelienſium. *Bauh. pin.* 467. *Du Ham. arb.* 1. *p.* 168. *t.* 66.

Ledum Matthioli. *Dalech. hiſt.* 230.

Ladanum. *Tabern. ic.* 1064

Mittländiſche Ciſtenroſe. W.

Habitat in Gallia *auſtrali.* ♄ (v. v.)

7. CISTUS *ladaniferus.*

C. fruticoſus exſtipulatus, foliis ſubſeſſilibus connatis lanceolato-linearibus ſupra glabris ſubtus tomentoſis, pedunculis bracteatis unifloris, capſulis decemlocularibus. *Lamarck. encycl.* 2. *p.* 16. *

C. arboreſcens exſtipulatus, foliis lanceolatis ſupra laevibus: petiolis baſi coalitis vaginantibus. *Hort. cliff.* 205. *Sauv. monſp.* 147, 150. *Mill. dict. n.* 6.

α. *undulatus*, foliis undulatis. *Ait. Kew.* 2. *p.* 282.

C. ladanifera hiſpanica incana. *Bauh. pin.* 467.

Ciſtus Ledon 1. anguſtifolius. *Cluſ. hiſt.* 1. *p.* 77.

β. *planifolius* foliis planis. *Ait. l. c.*

Ciſtus Ledon flore macula nigricante notato. *I. B. Comm. hort.* 1. *p.* 39. *t.* 20.

Ciſtus ladanifera hiſpanica ſalicis folio, flore candido. *Tournef. inſt.* 260.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 26.

Ladanum Ciſtenroſe. W.

Habitat in Hiſpaniae, Luſitaniae *collibus.* ♄ (v. v.)

Corolla *alba*, *magnitudine Papaveris ſomniferi.* Germen *torulis* 10. Stigma *ſeſſile absque ſtylo.*

8. CISTUS *monſpelienſis.*
C. arboreſcens exſtipulatus, foliis lineari-lanceolatis ſeſſilibus utrinque villoſis trinerviis. *Hort. cliff.* 205. *Hort. upſ.* 144. *Roy. lugdb.* 475. *Sauv. monſp.* 147.
C. ladanifera ſ. Ledon monſpelienſium anguſto folio nigricans. *Bauh. hiſt.* 2. *p.* 10.
Ciſtus Ledon foliis oleae, ſed anguſtioribus. *Bauh. pin.* 467.
Ciſtus oleaefolius. *Mill. dict. n.* 10.
Ledon narbonenſe. *Lob. ic. p.* 119.
Ledon 5. *Cluſ. hiſt.* 1. *p.* 79.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 27.
Franzöſiſche Ciſtenroſe. W.
Habitat in G. Narbonenſi *et* Regno Valentino. ♄ (v. ſ.)
Folia *ſuprema baſi latiora, reliqua linearia, ſubtus reticulata trinervia.* Racemus *longe pedunculatus, ſaepe bifidus.* Corollae *albae.*

*9. CISTUS *laxus.*
C. arboreſcens exſtipulatus, foliis ovato-lanceolatis undulatis denticulatis glabris, ſummis hirtis, foliolis calycinis ſubrotundo-cordatis. *Ait. Kew.* 2. *p.* 233.
C. (*hirſutus*) fruticoſus exſtipulatus, foliis ſeſſilibus oblongis obtuſis hirſutis, pedunculis multifloris, capſulis parvis calyce maximo et pyramidali tectis. *Lamarck. encycl.* 2. *p.* 17.*
Ledon IV. *Cluſ. hiſt.* 1. *p.* 78.
Schlaffäſtige Ciſtenroſe. W.
Habitat in Hiſpania, Luſitania. ♄
Affinis videtur C. capenſi, at Flos *albus.* W.

10. CISTUS *ſalvifolius.*
C. arboreſcens exſtipulatus, foliis ovatis petiolatis utrinque hirſutis. *Hort. cliff.* 205. *Roy. lugdb.* 475. *Sauv. monſp.* 150. *Mill. dict. n.* 8. *Scop. carn. n.* 643. *Willd. arb.* 62. *Cavan. ic.* 2. *p.* 31. *t.* 137.*
C. ſubarboreſcens exſtipulatus, foliis oppoſitis petiolatis oblongo-ovatis rugoſis ſubcrenulatis utrinque hirſutis, pedunculis florum longiſſimis ante anthelin nutantibus. *Iacq. Collect.* 2. *p.* 120. *t.* 8.*
C. fruticoſus, foliis petiolatis, ovatis, rugoſis, ſerratis. *Hall. helv. n.* 1021.

C. femina, folio ſalviae. *Bauh. pin.* 464.
C. femina. *Cluſ. hiſt.* 1. *p.* 70.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 28.
Salbeyblättrige Ciſtenroſe. W.
Habitat in Italia, Sicilia, Narbona, Helvetia, Carniolia. ♄ (v. v.)

Folia *ovata, obtuſa, avenia, minus mollia.* Pedunculi *laterales, ſolitarii, uniflori, foliis longiores.* Flores *albi.*

*11. CISTUS *heterophyllus.*

C. exſtipulatus, foliis ovato-lanceolatis baſi vaginantibus, margine revolutis, calycibus pedunculisque hirſutis ſubunifloris. *Desfont. fl. atl.* 1. *p.* 411. *t.* 104.
Verſchiedenblättrige Ciſtenroſe. W.
Habitat in collibus incultis Algeriae. ♄

Frutex *bipedalis erectus ramoſiſſimus.* Rami *juniores villoſi incani teretes.* Stipulae *nullae.* Folia *oppoſita ovato-oblonga ſeu elliptica, ſuperne laeviuſcula, inferne pallidiora, margine revoluta nervoſa, nervis transverſis villoſis, villis breviſſimis.* Petioli *breves connati baſi vaginantes.* Flores *terminales,* 1, 2, 3 *aut* 4 *pedunculati.* Pedunculi *hirſuti folioſi plerumque uniflori, nodulo interſecti foliola* 2 *lanceolata emittente.* Calyx *villoſus* 5-*phyllus, foliolis ovato-oblongis ſubaequalibus, duobus interioribus acutis.* Corolla *magna roſea.* Capſula *ſubrotunda villoſa,* 5-*locularis. Varietatem poſſideo diſtinctam, foliis rotundatis. Eadem planta ſaepe folia inferiora rotunda aut ſubrotunda, ſuperiora lanceolata profert, unde nomen C. heterophylli. Affinis C. incano.* Desfont. l. c.

12. CISTUS *incanus.*

C. arboreſcens exſtipulatus, foliis ſpathulatis tomentoſis rugoſis: inferioribus baſi vaginantibus connatis. *Hort. cliff.* 205. *Hort. upſ.* 143. *Roy. lugdb.* 475. *Mill. dict. n.* 2.
C. mas anguſtifolius. *Bauh. pin.* 464.
C. mas 2. *Cluſ. hiſt.* 1. *p.* 69. *hiſp.* 137.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 29.
Beſtäubte Ciſtenroſe. W.
Habitat in Hiſpania, G. Narbonenſi. ♄ (v. v.)

Rami *villoſi.* Folia *minime nervoſa in mucronem deſinentia, diſco parum flexuoſa, deſinunt baſi in Petiolos coalitos vaginantes, vel potius obovato-ſpathula-*

ta, inferioribus magis connatis et fere vaginantibus. Calyces *piloſi foliolis ſubcordatis.* Petala *purpurea. emarginata ſ. obcordata, integerrima, concava.*

13. CISTUS *creticus.*
C. arboreſcens exſtipulatus, foliis ſpathulato-ovatis rugoſo-venoſis ſcabris, petiolatis, petiolis diſtinctis, pedunculis unifloris, calycinis foliolis acuminatis. W.
C. arboreſcens exſtipulatus foliis ſpathulato-ovatis petiolatis enerviis ſcabris, calycinis lanceolatis. *Mat. med. p.* 137. *Mill. dict. n.* 9. *Iacq. ic. rar.* 1. *t.* 95. *Collect.* 1. *p.* 80. *
C. fruticoſus exſtipulatus, foliis ſpathulato-ovatis petiolatis rugoſis hirtis margine undulatis, pedunculis brevibus unifloris, foliis calycinis mucronatis villoſis. *Lamarck encycl.* 2. *p.* 13. *
C. ladanifera cretica. *Tournef. cor.* 19. *it.* 1. *p.* 30. *Buxb. cent.* 3. *p.* 34. *t.* 64. *f.* 1.
Ciſtus Ledon cretenſe. *Bauh. pin.* 467. *non ſynonyma.*
Ladanum creticum. *Alp. exot.* 89. *t.* 88.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 29.
Cretiſche Ciſtenroſe. W.
Habitat in Creta, Syria. ♄ (v. v.)
Flores *purpurei.*

*14. CISTUS *parviflorus.*
C. arboreſcens exſtipulatus, foliis ovatis acutis tomentoſis rugoſo-venoſis petiolatis, petiolis baſi vaginantibus, calycibus villoſis acuminatis. W.
C. (*parviflorus*) fruticoſus exſtipulatus, foliis petiolatis ovatis acutis tomentoſis, pedunculis villoſis ſubunifloris. *Lamarck. encycl.* 2. *p.* 14. *
Ciſtus mas creticus breviori folio, parvo flore. *Tournef. cor.* 19.
Kleinblumige Ciſtenroſe. W.
Habitat in Creta. ♄ (v. ſ.)
Similis praecedenti, ſed petiolis *baſi in vaginam breviſſimam connatis,* foliis *tomentoſis acutis,* calycibus *villoſis,* corollisque *minoribus diverſus.* W.

15. CISTUS *albidus.*
C. arboreſcens exſtipulatus, foliis ovato-lanceolatis tomentoſis incanis ſeſſilibus ſubtrinerviis. *Sauv. monſp.* 150. *Mill. dict. n.* 7. *Gouan. monſp.* 225. *Ger. prov.* 398. *Garid. aix.* 114.

C. mas, folio oblongo incano. *Bauh. pin.* 464.
C. mas 1. *Cluſ. hiſt.* 1. *p.* 68.
C. mas 4. monſpelienſis, folio oblongo albido. *Bauh. hiſt.* 2. *p.* 3.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 31.
Weiſsliche Ciſtenroſe. W.
Habitat in G. Narbonenſi, Hiſpania. ♄ (v. v.)

Differt a C. incano, *cui ſimillimus*, Ramis *tomentoſis nec piloſis*; Foliis *pallidioribus, mollibus, horizontalibus, ſeſſilibus, minime petiolatis aut vaginantibus, lanceolatis, ſubtrinerviis.* Calyces *non hirti foliolis exterioribus longioribus, lateribus reflexis.* Pedunculi *ex apice ramorum, plures, foliis vix longiores.* Petala *purpurea, non emarginata, margine ſubcrenata, planiuſcula.*

*16. CISTUS *ſericeus.*

C. arboreſcens exſtipulatus, foliis ovatis tomentoſis trinerviis, inferioribus petiolatis, ſupremis ſeſſilibus, pedunculis hirtis. *Vahl. ſymb.* 1. *p.* 37. *
C. latifolius magno flore. *Barrel. ic.* 1315.
Seidenartige Ciſtenroſe. W.
Habitat in Hiſpania. ♄

Caulis *bi-ſ. tripedalis.* Rami *teretes denſe tomentoſi incano-albi.* Folia *molliſſima obtuſa plana ſubpollicaria, quatuor ſuprema ſeſſilia, baſi ſubcordata, reliqua, in petiolum brevem deſinentia.* Pedunculus *ex apice ramorum ſolitarius erectus ſpithamaeus incanus, totus obſitus uti pedicelli pilis longis purpuraſcentibus.* Pedicelli *verſus ſummitatem patentes, inferiores triflori, ſuperiores uniflori.* Calyx *pilis mollibus ſericeis hirſutiſſimus.* Foliola interiora 3 *ovata intus glaberrima*; exteriora *lanceolata ad baſin calycis.* Petala *purpurea baſi macula flava.* Filamenta *purpurea.* Antherae *flavae.* Vahl l. c.

*17. CISTUS *hybridus.*

C. arboreſcens exſtipulatus, foliis ovatis petiolatis incanis, ramis ſquamoſis, pedunculis elongatis hirtis. *Vahl. ſymb.* 1. *p.* 37. *
Baſtard Ciſtenroſe. W.
Habitat in Hiſpania. ♄

Frutex *altitudine C. halimifolii.* Rami *teretes apice angulati incani, ſquamis flaveſcentibus tecti fere ut in*

C. ſquamato et halimifolio. Folia *omnia petiolata ovalia plana, juniora undulata utrinque incana, ſupra oculo armato minutiſſime punctata, nervoſa fragilia ſemipollicaria.* Racemus *terminalis.* Calycis *foliola exteriora caduca.* Corolla *purpurea.* Germen *villoſum.* Rami *et* Folia *C. halimifolii,* Pedunculus *et* Calyx *antecedentis. A praecedente dignoſcitur,* foliis *omnibus petiolatis albidioribus,* ramis *flavo-ſquamoſis.* Vahl l. c.

18. CISTUS *criſpus.*

C. arboreſcens exſtipulatus, foliis lanceolatis pubeſcentibus trinerviis undulatis. *Hort. cliff.* 206. *Roy. lugdb.* 475. *Sauv. monſp.* 147. *Mill. dict. n.* 16. *Kniph. cent.* 5. *n.* 23.

C. fruticoſus exſtipulatus, foliis lanceolatis undulatis trinerviis pubeſcentibus, floribus ſubſeſſilibus. *Cavan. ic.* 2. *p.* 57. *t.* 174. *

C. mas, foliis chamaedryos. *Bauh. pin.* 464.

C. ladanifera. *Blackw. t.* 197.

C. mas 5. *Cluſ. hiſt.* 1. *p.* 69.

C. mas foliis undulatis et criſpis. *Tournef. inſt.* 259. R. *Houttuyn Lin. Pfl. Syſt.* 4. *p.* 32.

Krauſe Ciſtenroſe. W.

Habitat in Luſitania. ♄ (v. v.)

Ramuli, pedunculi et calyces tenui lana adſperſi. Flores *purpurei.* Calyces *lanceolati.*

*19. CISTUS *formoſus.*

C. arboreſcens exſtipulatus, foliis lanceolato-obovatis trinerviis hirtis, pedunculis calycibusque piloſis, petalis emarginatis. W.

Ciſtus formoſus. *Curt. mag.* 264.

Schöne Ciſtenroſe. W.

Habitat in Luſitania. ♄

Petala *flava baſi macula atroſanguinea notata. Planta ceterum ſequenti ſimillima.* W.

20. CISTUS *halimifolius.*

C. arboreſcens exſtipulatus, foliis lanceolato-obovatis trinerviis incanis pedunculis calycibusque glabris, petalis ſubretuſis, foliolis duobus calycinis linearibus. W.

C. arboreſcens exſtipulatus, foliolis duobus calycinis linearibus. *Hort. cliff.* 205. *Roy. lugdb.* 475. *Sauv. monſp.* 147. *Mill. dict. n.* 17. *Cavan. ic. rar.* 2. *p.* 31. *t.* 138?

C. arboreſcens exſtipulatus, foliis lanceolatis incanis, pedunculis virgatis. *Vahl. ſymb.* 1. *p.* 38. *
C. femina, portulacae marinae folio latiore obtuſo. *Bauh. pin.* 465.
C. folio halimi 1. *Cluſ. hiſt.* 1. *p.* 71.
C. halimi minoris folio. *Barrel. ic.* 287.
β. C. femina portulacae marinae folio anguſtiore mucronato. *Bauh. pin.* 465.
C. halimi folio 2. *Cluſ. hiſt.* 1. *p.* 71.
γ. Ciſtus fruticoſus erectus, foliis oppoſitis ſeſſilibus utrinque incanis. *Mill. ic. t.* 290.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 32.
Meldenartige Ciſtenroſe. W.
Habitat in Luſitaniae, Hiſpaniae *maritimis.* ♄ (v. ſ.) Flores *lutei.*

*21. CISTUS *elongatus.*

C. arboreſcens exſtipulatus: foliis lanceolatis incanis, pedunculo elongato diphyllo, calycibusque racemoſis hirſutis. *Vahl. ſymb.* 1. *p.* 38. *
Verlängerte Ciſtenroſe. W.
Habitat in Hiſpania. ♄ (v. ſ. ſ. fl.)

Dignoſcitur a reliquis, pedunculis *diphyllis absque tomento, ſubtus modo piloſis, a C. halimifolio*, foliolis *floralibus alternis.* Vahl l. c.

22. CISTUS *libanotis.*

C. arboreſcens exſtipulatus, foliis linearibus revolutis, floribus umbellatis. *Sp. pl.* 739.
C. fruticoſus exſtipulatus, foliis linearibus margine revolutis, floribus ſubumbellatis, calyce triphyllo. *Lamarck. encycl.* 2. *p.* 17.
C. anguſto libanotidis folio, flore ſingulari. *Barr. ic.* 294.
Ciſtus Ledon anguſtis foliis. *Bauh. pin.* 467.
Ciſtus Ledon, foliis Rorismarini ſubtus incanis. *Bauh. pin.* 467.
Ledon VIII. *Cluſ. hiſt.* 1. *p.* 80.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 33.
Rosmarinblättrige Ciſtenroſe. W.
Habitat in Hiſpania. *Loefling.* ♄ (v. ſ.)

Frutex *habitu Roſmarini.* Caulis *nudus, purpuraſcens.* Folia *oppoſita, exſtipulata, ſeſſilia, linearia, ſed utrinque revoluta, laevia, marginibus baſeos hir-*

ſuta, internodiis longiora, ad apices ramorum approximata. Folia floralia ſ. ſuprema breviora, ovata, plana. Pedunculi *terminales, foliis longiores*: Floribus *aliquot luteis.* Calyces *ovati, viſcidi.*

* * *Exſtipulati ſuffruticoſi.*

23. CISTUS *umbellatus.*
C. ſuffruticoſus procumbens exſtipulatus, foliis oppoſitis linearibus, floribus umbellatis. *Sp. pl.* 739.
C. foliis linearibus non ſtipulatis, caule florali umbellato lignoſo. *Guett. ſtamp.* 2. *p.* 160. *Dalib. pariſ.* 158.
Ciſtus Ledon foliis thymi. *Bauh. pin.* 467.
Ledon X. *Cluſ. hiſt.* 1. *p.* 81.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 219.
Doldenblütige Ciſtenroſe. W.
Habitat in Gallia, Hiſpania. ♄
Umbellae pedunculatae ſunt.

24. CISTUS *Laevipes.*
C. ſuffruticoſus adſcendens exſtipulatus, foliis alternis faſciculatis filiformibus glabris, pedunculis racemoſis. *Amoen. acad.* 4. *p.* 475. *Iacq. hort. t.* 158. *Cavan. ic.* 2. *p.* 56. *t.* 173. *
C. ſuffruticoſus procumbens, foliis alternis confertis inaequalibus ſetaceis. *Ger. prov.* 394. *t.* 14.
C. humilis maſſiliotica, camphoratae tenuiſſimis foliis glabris. *Pluk. alm.* 107. *t.* 84. *f.* 6.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 220.
Glattſtrunkige Ciſtenroſe. W.
Habitat Monſpelii. ♄ (v. v.)
Ex ſingulis foliorum alis faſciculus foliorum minor. Flores *lutei.*

25. CISTUS *calycinus.*
C. ſuffruticoſus erectus exſtipulatus, foliis linearibus, pedunculis unifloris, calycibus triphyllis. *Mant.* 565.
Ciſtus Fumana variet. A. *Desfont. fl. atl.* 1. *p.* 414. *t.* 105.
C. (*ericoides*) exſtipulatus, caule fruticoſo erecto, foliis alternis imbricatis ſemicylindricis brevibus. *Cavan. ic.* 2. *p* 56. *t.* 172. *
C. humilis [Chamae-Ciſtus ericae folio, luteus et elatior. *Bauh. pin.* 465. *Pluk. alm.* 107. *t.* 83. *f.* 6.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 221.

Heidekrautartige Ciſtenroſe. W.
Habitat in Europa *auſtrali.* ♄

Fruticulus *pedalis, erectus; ramis oppoſitis, rubeſcentibus.* Folia *oppoſita, linearia, laevia, ſubtus obtuſe carinata: ex axillis foliola conferta.* Pedunculi *terminales, ſolitarii, uniflori, vix foliis longiores.* Calyx *triphyllus, laevis, aequalis.* Corolla *pentapetala, patens, flava, vix margine rubro.* Stamina 16, *breviſſima, lutea, omnia fertilia.* Piſtillum *album.* Stylus *breviſſimus.* Stigma *verrucoſum.*

Obſ. *Similis Fumanae.*

Clariſſ. Desfontaines hunc cum ſequente conjunxit. W.

26. CISTUS *Fumana.*

C. ſuffruticoſus procumbens exſtipulatus, foliis alternis, linearibus margine ſcabris, pedunculis unifloris. *Sp. pl.* 740. *Iacq. auſtr. t.* 252. *Fabric. helmſt. p.* 196. *Scop. carn. n.* 644. *Crantz. auſtr. p.* 102. *Kniph. cent.* 9. *n.* 22. *Hoffm. germ.* 185. *Roth. germ. I.* 228. *II.* 584.

C. foliis duris, linearibus, confertis, petiolis unifloris, calycibus glabris. *Hall. helv. n.* 1032.

C. caule procumbente, foliis alternis. *Fl. ſuec.* 435, 474.

C. minor, brevi vermiculatoque folio Hiſpanicus. *Barr. ic.* 285. *Hall.*

Helianthemum foliis linearibus ſaepius alternis non ſtipulatis, pedunculis unifloris, caule lignoſo. *Guett. ſtamp.* 2. *p.* 18.

C. fruticoſus procumbens, foliis aciformibus nudis, floribus auriculatis. *Sauv. monſp.* 46.

Chamae-Ciſtus 6. *Cluſ. hiſt.* 1. *p.* 75.

β. Chamae-Ciſtus, ericae folio, luteus humilior. *Bauh. pin.* 666.

Helianthemum tenuifolium glabrum, per humum ſparſum, luteo flore. *Bauh. hiſt.* 2. *p.* 18.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 221.

Dünne Ciſtenroſe. W.

Habitat in Gallia, Gotlandia, Helvetia. ♄ (v. ſ.)

Stamina *circiter* 16. *antherifera et totidem exteriora caſtrata.* Folia *laevia, margine ſpinulis ſcabris aliquot minutiſſimis.*

27. CISTUS *canus.*

C. ſuffruticoſus procumbens exſtipulatus, foliis oppoſi-

tis obovatis villoſis ſubtus tomentoſis, floribus ſubumbellatis. *Sp. pl.* 740. *Iacq. auſt. t.* 277. *Scop. carn. n.* 647. *Allion. pedem. n.* 1664. *t.* 45.

C. exſtipulatus, foliis oppoſitis, incanis nudis. *Crantz auſtr. p.* 100.

C. foliis ovatis ſubtus incanis, alis nudis; caule procumbente. *Sauv. monſp.* 148.

Chamae Ciſtus foliis myrti minoris incanis. *Bauh. pin.* 466.

Chamae Ciſtus foliis myrti tarentinae canis et cinereis. *Bauh. hiſt.* 2. *p.* 18.

Chamae Ciſtus 3. *Cluſ. hiſt.* 1. *p.* 74.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 222.

Graue Ciſtenroſe. W.

Habitat in G. Narbonenſi, Hiſpania. ♄ (v. ſ)

Folia *ovalia ſ. obovata, obtuſa, ſupra pallide cineraſcentia, ſubtus tomentoſo-cana.* Umbella *terminalis,* 4-*flora.*

*28. CISTUS *ſcabroſus.*

C. ſuffruticoſus exſtipulatus, foliis oppoſitis ovatis piloſo-ſcabris trinerviis, calycibus triphyllis. *Ait. Kew.* 2. *p.* 236.

Rauhe Ciſtenroſe. W.

Habitat in Italia, Luſitania. ♄

Caules *decumbentes teretes, pilis brevibus ſtellatis denſe adſperſi ſcabri.* Rami *breves.* Folia *ſubpetiolata trinervia uncialia.* Flores *terminales ſubpaniculati.* Calyx *triphyllus: foliola ovato-lanceolata acuminata, extus pubeſcentia pilis ſtellatis et villo longo aequalia quadrilinearia.* Petala *obovata ſubretuſa, calyce fere duplo longiora, intenſe flava, baſi lutea.* Ait. l. c.

*29. CISTUS *cinereus.*

C. ſuffruticoſus exſtipulatus, foliis oppoſitis ovatis acutis canis, calycibus obtuſis, racemis paniculatis. W.

C. exſtipulatus ſuffruticoſus, foliis oppoſitis ſubtomentoſis ovato-acutis, floribus racemoſis, racemis ſubternis terminalibus. *Cavan. ic.* 2. *p.* 33. *t.* 141.*

Aſchfarbene Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Affinis videtur praecedenti, diverſus tamen: foliis *evidenter petiolatis venoſis,* calycinis *foliolis tribus ovatis obtuſis, binis linearibus,* petalis *integerrimis. C. polyanthos huic habitu ſimillimus, ſed eſt ſtipulatus.* W.

*30. CISTUS *ocymoides.*

C. foliis obovatis trinerviis, ramulorum utrinque incanis apice reflexis, calycibus racemoſis pedunculisque glaberrimis. *Vahl. ſymb.* 3. *p.* 68. *

C. (*ocymoides*) ſubfruticoſus exſtipulatus, foliis petiolatis dorſo carinatis incanis minimis, pedunculis ramoſis umbellato-paniculatis. *Lamarck encycl.* 2. *p.* 18. *

C. folio ſampſuci Cluſii cui macula in flore. *Bauh. hiſt.* 2. *p.* 6.

C. folio ſampſuci. *Cluſ. hiſt.* 1. *p.* 72.

β. ramis laxe villoſis. *Lamarck l. c.*

C. (*ſampſucifolius*) fruticoſus exſtipulatus, foliis ovato-oblongis carinatis ſeſſilibus, pedunculis elongatis ramoſo-paniculatis. *Cavan. ic.* 1. *p.* 65. *t.* 96. *

Baſilienkrautartige Ciſtenroſe. W.

Habitat in Murcia, Baetica. ♄

31. CISTUS *italicus.*

C. ſuffruticoſus exſtipulatus, foliis oppoſitis hiſpidis: inferioribus ovatis; ſuperioribus lanceolatis, ramis patentibus. *Sp. pl.* 740.

Helianthemum, ſerpilli folio villoſo, flore pallido, italicum. *Barr. rar.* 510. *t.* 366.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 223.

Italiäniſche Ciſtenroſe. W.

Habitat in Italia. ♄

Frutex *Caule erecto, ſpithamaeo: Ramis oppoſitis, longioribus, patentibus, ſubdeflexis, rufis.* Folia *oppoſita, exſtipulata, utrinque adſperſa pilis ſtrigoſis, raris, ſubgeminis et ſubciliata: inferiora petiolata, ovata; ſuperiora ſubſeſſilia.* Racemus *terminalis.* Calyces *hiſpidi.* Corolla *pallida petalis vix emarginatis,* An *varietas C. cani?*

32. CISTUS *marifolius.*

C. ſuffruticoſus exſtipulatus divaricatus, foliis ovatis acutis oppoſitis petiolatis planis ſubtus tomentoſis, floribus racemoſis. W.

C. ſuffruticoſus exſtipulatus, foliis oppoſitis oblongis petiolatis planis ſubtus incanis. *Sp. pl.* 741. *Cavan. ic.* 2. *p.* 34. *t.* 143. *

Helianthemum luteum, thymi durioris folio. *Barr. rar.* 521. *t.* 441.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 223.

Marumblättrige Ciſtenroſe. W.

Habitat Maſſiliae, Veronae *inque* Hiſpania. ♄

Variat Foliis lanceolatis ſupra glabris, et ovalibus ſupra ſubpiloſis, ſemper Folia ſubtus denſe tomentoſa et incano-alba.

*33. CISTUS *origanifolius.*

C. ſuffruticoſus exſtipulatus adſcendens, foliis ovatis obtuſis oppoſitis petiolatis planis utrinque piloſis, racemis ſecundis. W.

C. (*origanifolius*) ſuffruticoſus exſtipulatus, foliis oppoſitis petiolatis ovatis utrinque piloſis. *Lamarck encycl. 2. p. 21.* * *Cavan. ic. 3. p. 31. t. 262. f. 1.* *

Doſtenblättrige Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Corollae *flavae vix calyce majores duplo minores, quam in praecedente.* Racemi *ſecundi geminati pedunculati piloſi ut caulis.* Folia *ſeniora utrinque piloſa viridia, juniora albida.* W.

*34. CISTUS *mollis.*

C. ſuffruticoſus exſtipulatus, foliis ſubrotundo-ovatis obtuſis petiolatis planis utrinque tomentoſis mollibus. W.

C. (*mollis*) ſuffruticoſus exſtipulatus, foliis oppoſitis ſuborbiculatis valde tomentoſis. *Cavan. ic. 3. p. 32.* *

Weiche Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Caules *ſuffruticoſi rubentes diffuſi ſemipedales ramoſi tomentoſi.* Folia *ſuborbiculata oppoſita petiolata mollia tomentoſa, nervis obliteratis.* Stipulae *nullae.* Flores *racemoſi, racemis terminalibus ſolitariis.* Corolla *triplo major quam in praecedenti.* Cavan. l. c.

*35. CISTUS *dichotomus.*

C. ſuffruticoſus exſtipulatus dichotomus, foliis ovatis acutis glabris margine revolutis oppoſitis petiolatis, floribus racemoſis. W.

C. (*dichotomus*) ſuffruticoſus exſtipulatus dichotomus, foliis ovatis oppoſitis glabris, floribus racemoſis. *Cavan. ic. 3. p. 32. t. 263. f. 1.* *

Gabeläſtige Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Folia *parva facie Thymi Piperellae, breve petiolata glabra margine revoluta.* Racemi *tenues pauciflori.* Flores *ſaturate flavi vix magnitudine Spergulae nodoſae.* W.

36. CISTUS *anglicus.*

C. ſuffruticoſus exſtipulatus procumbens, foliis oppoſitis oblongis revolutis piloſis, floribus racemoſis. *Mant.* 245.

Ciſtus hirſutus. *Hudſ. angl.* 232.

Helianthemum alpinum, folio piloſellae minoris Fuchſii. *Bauh. hiſt.* 2. *p.* 13. *Seguier. ver. p.* 196.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 224.

Engliſche Ciſtenroſe. W.

Habitat in quibusdam Angliae *locis.* ♄

Caulis *vix ſemipedalis, obliquus, totus ſcaber.* Folia *oppoſita, lanceolata* (*Hyſſopi*), *ſcabriuſcula, nec laevia, utrinque viridia.* Flores *racemoſi, albi, nutantes, ſub antheſi erecti.*

*37. CISTUS *vinealis.* W.

C. ſuffruticoſus procumbens exſtipulatus, foliis petiolatis oblongis obtuſis ſubtus incano-tomentoſis, floribus racemoſis. W.

C. caulibus duris procumbentibus, foliis ellipticis hirſutis, inferne cum calycibus tomentoſis. *Hall. helv. n.* 1035.

Weinbergs Ciſtenroſe. W.

Habitat in Germaniae *vineis, inque* Helvetia. ♄ (v. v.)

Hanc ſpeciem prope Kelme, Rulsdorf et Schmon haud procul Halae Saxonum legi, et pro C. marifolio antea habui, ſed a C. marifolio diverſiſſima foliorum forma, ramis procumbentibus totoque habitu. Linnaeus vero in deſcriptione C. marifolii hunc cum C. marifolio confundere videtur. A C. oelandico abunde diſtinctus foliis ſubtus tomentoſis albis. W.

38. CISTUS *oelandicus.*

C. ſuffruticoſus procumbens exſtipulatus, foliis oppoſitis oblongis utrinque glabris: petiolis ciliatis, petalis emarginatis. *Sp. pl.* 741. *Iacq. auſtr. t.* 399.

C. caulibus duris procumbentibus, foliis conjugatis, ellipticis, ſubhirſutis, petiolis confertis unifloris, calycibus tomentoſis. *Hall. helv. n.* 1034. *Iacq.*

C. ſuffruticoſus procumbens exſtipulatus, foliis oppoſitis oblongis utrinque glabris, petiolis ciliatis, petalis emarginatis. *Gouan. monſp.* 265. *illuſtr. p.* 32.

C. alpeſtris. *Iacq. vind.* 248. *Scop. carn. n.* 645. *t.* 23.

C. (*alpeſtris*) ſuffruticoſus, procumbens, exſtipulatus,

foliis oppoſitis ciliatis, calycibus laevibus. *Crantz auſtr. p.* 103. *t.* 6. *f.* 1.

C. (*Seguieri*) exſtipulatus, foliis caulinis inferioribus ovatis oppoſitis revolutis, ſuperioribus expanſis. *Crantz auſtr. p.* 104.

C. caule procumbente, foliis oblongis utrinque glabris, ſtipulis nullis. *Fl. ſuec.* 434. * 473. *

C. ſuffruticoſus procumbens, foliis nudis oppoſitis viridibus, floribus ſubumbellatis, calycibus hirſutis. *Ger. prov.* 396.

Helianthemum alpinum ſerpillifolio nigricante et hirſuto. *Segu. veron.* 3. *p.* 195. *t.* 6. *f.* 2.

Chamae-Ciſtus II. *Cluſ. hiſt.* 1. *p.* 73.

C. helianthemos, flore parvo luteo. *Bauh. hiſt.* 2. *p.* 17.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 224.

Oelandiſche Ciſtenroſe. W.

Habitat in rupibus Galliae, Oelandiae, Helvetiae, Auſtriae. ♄ (v. ſ.)

Calyx *lineatus pubeſcens.*

*39. CISTUS *alternifolius.*

C. ſuffruticoſus exſtipulatus, foliis alternis, pedunculis lateralibus terminalibusque ſolitariis unifloris. *Vahl ſymb.* 1. *p.* 38.

C. (*braſilienſis*) ſubfruticoſus exſtipulatus, foliis alternis ovato-oblongis villoſis ſeſſilibus, pedunculis unifloris. *Lamarck encycl.* 2. *p.* 22. *

Wechſelblättrige Ciſtenroſe. W.

Habitat in Braſilia. ♄

Fruticulus *erectus*; *ramis teretibus villoſis, cortice fuſco.* Folia *alterna exſtipulata ſeſſilia oblonga erecta plana integerrima utrinque villoſa, ſiccitate nigricantia unguicularia.* Pedunculi *verſus apicem ſolitarii, interdum bini vel terni uniflori in axillis foliorum inferiorum ramorum patentes, folio duplo longiores villoſi, apice parum craſſiores, folia duo caduca ad apicem linearia.* Calyces *ovati acuti hirſuti.* Calyx *et* Inflorescentia *C. ſalicifolii at fruticoſus.* Vahl l. c.

*** *Exſtipulati herbacei.*

*40. CISTUS *globularifolius.*

C. exſtipulatus perennis, caule ſimplici ſubnudo, foliis radicalibus petiolatis ſpathulatis obtuſis. *Lamarck encycl.* 3. *p.* 22.

Helianthemum luſitanicum globulariae folio. *Tournef. inſt.* 250.
Kugelblumenartige Ciſtenroſe. W.
Habitat in Luſitania. ♃.

Radix *perennis magna lignoſa multiceps.* Folia *petiolata ſpathulata apice rotundata et obtuſa trinervia pubeſcentia interdum fere glabra.* Caulis 4-ſ. 6-*uncialis ſimplex glabriuſculus* 2 ſ. 3 *paribus foliorum diſtantium parvorum et acutorum inſtructus.* Flores *ut in ſequenti.* Lamarck l. c.

41. CISTUS *Tuberaria.*

C. exſtipulatus perennis, foliis radicalibus ovatis trinerviis tomentoſis; caulinis glabris lanceolatis; ſummis alternis. *Sp. pl.* 741. *Cavan. ic.* 1. *p.* 65. *t.* 67. *
C. perennis, foliis humi adpreſſis villoſis; ſummis glabris. *Sauv. monſp.* 149.
C. folio plantaginis. *Bauh. pin.* 465.
Helianthemum Tuberaria. *Mill. dict. n.* 10.
Helianthemum, plantaginis folio, perenne. *Buxb. cent.* 3. *p.* 33. *t.* 63.
Tuberaria noſtras. *Bauh. hiſt.* 2. *p.* 12.
Tuberaria major. *Bauh. hiſt.* 2. *p.* 12.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 225.
Stumpfblättrige Ciſtenroſe. W.
Habitat in Galloprovincia, Hiſpania, Piſis. ♃.

*42. CISTUS *plantagineus.* W.

C. herbaceus exſtipulatus, foliis lanceolatis utrinque attenuatis trinerviis hirtis, racemis ebracteatis, petalis denticulatis. W.
Ciſtus ſerratus. *Desfont. fl. atl.* 1. *p.* 416. (*excluſo ſynonymo Cavanilliano.*)
C. (*lanceolatus*) ſuffruticoſus, foliis lanceolatis trinerviis piloſis. *Vahl. ſymb.* 2. *p.* 62?
Helianthemum creticum annuum lato Plantaginis folio, flore aureo. *Tournef. cor.* 18.
Wegetrittblättrige Ciſtenroſe. W.
Habitat in Creta *et* Africa *boreali.* ⊙ (v. ſ.)

Caulis *pedalis vel ſesquipedalis pilis longis rectis adſperſus.* Folia *radicalia oblonga acuminata in petiolum attenuata tri-vel quinquenervia utrinque hirta, ſuperne pilis ſimplicibus ſparſis adpreſſis, inferne pilis copioſioribus ſtellatis tecta.* Folia *caulina oppoſita lanceolata utrinque attenuata ſeſſilia trinervia ſeſ-*

qui-vel biuncialia hirta, piliſque ut in foliis radicalibus ſed copioſioribus obſita. Stipulae *nullae niſi in ſummo pare foliorum, quae ſemiunciales lineares hirtae.* Racemi *bini in apice caulis ut videtur ſecundi.* Pedunculi *longiſſimi filiformes.* Bracteae *nullae.* Calyx *pentaphyllus, piloſus, foliolis lanceolatis, binis exterioribus parum brevioribus et obtuſioribus.* Corolla *flava immaculata Helianthemi, petalis denticulatis.* W.

*43. CISTUS *ſerratus.*

C. herbaceus exſtipulatus, foliis oppoſitis lanceolatis trinerviis hirtis viſcoſis, radicalibus obovatis, racemis ebracteatis, petalis ſerratis. W.

C. (*ſerratus*) herbaceus exſtipulatus, foliis oppoſitis lanceolatis ovatis trinerviis, petalis ſerratis. *Cavan. ic.* 2. *p.* 57. *t.* 175. *f.* 1. *

Geſägte Ciſtenroſe. W.

Habitat in Hiſpania. ☉

Petala *flava, baſi macula magna nigra notata. A praecedente diſtinctus:* ſtatura *triplo minori*, foliis *obtuſis viſcoſis*, petalis *baſi maculatis. A C. guttato diverſus:* foliis *viſcoſis*, petalis *ſerratis baſi maculatis nec tantum punctatis.* W.

44. CISTUS *guttatus.*

C. herbaceus exſtipulatus, foliis oppoſitis oblongis trinerviis hirtis, radicalibus obovatis, racemis ebracteatis, petalis integris. W.

C. herbaceus exſtipulatus, foliis oppoſitis lanceolatis trinerviis, racemis ebracteatis. *Hudſ. angl.* 204.

C. annuus, foliis radicalibus oppoſitis lanceolatis trinerviis, ramiferis ternatis linearibus. *Sauv. monſp.* 148.

C. foliis oblongo-acuminatis non ſtipulatis, caule florali racemoſo. *Guett. ſtamp.* 2. *p.* 181. *Dalib. pariſ.* 158.

C. flore pallido, punicante macula inſignito. *Bauh. pin.* 465.

Helianthemum guttatum. *Mill. dict. n.* 18.

Helianthemum flore maculoſo. *Col. ecphr.* 2. *p.* 78. *t.* 77.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 226.

Getüpfelte Ciſtenroſe. W.

Habitat in G. Narbonenſi, Italia, Anglia. ☉ (v. v.)

Caulis *erectus, ut tota planta pilis patentibus.* Folia *ſeſſilia, oppoſitionibus* 3 *ſ.* 4. Racemus *filiformis*

ſaepe ebracteatur; deflorati flores penduli; Flos *et* Fructus *erecti.* Petalorum *baſis macula rubra.* Calyx *foliolis* 2 *patulis.*

45. CISTUS *canadenſis.*
C. herbaceus exſtipulatus, foliis omnibus alternis lanceolatis, caule adſcendente.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 227.
Canadiſche Ciſtenroſe. W.
Habitat in Canada. *Kalm.* ♃
Facies *C. Helianthemi, ſed Folia alterna.*

* * * * *Stipulati herbacei.*

*46. CISTUS *punctatus.* W.
C. herbaceus erectus pubeſcens ſtipulatus, racemo terminali, foliis oppoſitis oblongis, inferioribus obovatis. W.
Punctirte Ciſtenroſe. W.
Habitat - - - - ⊙ (v. ſ.)
Planta *annua digitalis facie C. guttati.* Caulis *erectus ſtrictus ramis erectis oppoſitis ſimplicibus caule brevioribus.* Folia *oppoſita ut omnes partes tenui pube induta, petiolata, inferiora obovata obtuſa, ſuperiora oblonga acutiuſcula.* Stipulae *lineari-lanceolatae.* Racemus *elongatus terminalis.* Pedunculi *uniflori erecti.* Bractea *ovato-lanceolata parva non ad baſin pedunculorum ſed intermedia.* Calycis *foliola bina linearia reliquis minora patula.* W.

47. CISTUS *ledifolius.*
C. herbaceus erectus glaber ſtipulatus, floribus axillaribus, pedunculis erectis calyce brevioribus. W.
C. herbaceus erectus glaber ſtipulatus, floribus ſolitariis ſubſeſſilibus, folio ternato oppoſitis. *Hort. upſ.* 144. *Sauv. monſp.* 178. *Mill. dict. n.* 20. *ſub Helianthemo.*
C. ſtipulis quaternis, foliis oblongis, caule erecto. *Hort. cliff.* 475.
C. ledi folio. *Bauh. pin.* 465.
C. annuus, foliis ledi. *Lob. ic.* 2. *p.* 118.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 227.
Porſtblättrige Ciſtenroſe. W.
Habitat in Monſpelii. ⊙ (v. v.)
Calyx corolla longior: foliolis 3 *erectis,* 2 *patulis.* Capſulae *longitudine calycis.*

48. CISTUS *ſalicifolius.*
C. herbaceus patulus villoſus ſtipulatus, floribus axillaribus, pedunculis horizontalibus calyce longioribus. W.
C. herbaceus patulus villoſus ſtipulatus, floribus racemoſis erectis: pedicellis horizontalibus. *Hort. upſ.* 144. *Cavan. ic.* 2. *p.* 35. *t.* 144. *
C. ſtipulis quaternis, foliis ſubovatis ſubtus tomentoſis, caule procumbente. *Hort. cliff.* 206.
C. folio ſalicis. *Bauh. pin.* 465. *Lob. ic.* 2. *p.* 118.
C. annuus 1. *Cluſ. hiſt.* 1. *p.* 76.
Helianthemum ſalicifolium et fugacium. *Mill. dict. n.* 21. *et* 19.
Helianthemum annuum humile, foliis ovatis, flore fugaci. *Segu. ver.* 3. *p.* 297. *t.* 6. *f.* 3. *opt.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 228.
Weidenblättrige Ciſtenroſe. W.
Habitat in Luſitania, Hiſpania. ⊙ (v. ſ.)
Capſulae *calyce longiores.*

49. CISTUS *niloticus.*
C. herbaceus ſtipulatus erectus, ſubtomentoſus, floribus racemoſis ſolitariis ſeſſilibus oppoſitifoliis. *Mant.* 246.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 229.
Nil Ciſtenroſe. W.
Habitat in Aegypto. ⊙ (v. ſ.)
Caulis *ſublignoſus, pedalis, erectus, teres, ſubtomentoſus.* Rami *radicales adſcendentes, breviores; Caulini verſus ſummitatem, alterni, erecti, pauci.* Folia *oppoſita, petiolata, elliptica, patula, ſubtomentoſa, venoſa, articulis longiora.* Stipulae *quaternae, enſiformes, folio dimidio breviores, perſiſtentes.* Racemus *terminalis, erectus, ſtrictus.* Flores *alterni, ſubſeſſiles, ſolitarii, oppoſitifolii folio cum ſtipulis* 2, *caulinis ſimili.* Flos *petalis luteis.* Calyx *pentaphyllus, erectus: foliolis interioribus* 3 *trinerviis, acuminatis; exterioribus* 2 *linearibus, brevioribus, patulis.*
Simillimus Ciſto ledifolio, *ſed multoties major, nec laevis.*

50. CISTUS *aegyptiacus.*
C. herbaceus erectus ſtipulatus, foliis lineari-lanceolatis petiolatis, calycibus inflatis corollis majoribus. *Sp. pl.* 742. *Iacq. obſ.* 3. *p.* 17. *t.* 68. *
Houttuyn Lin. Pfl. Syſt. 7. *p.* 229.

Aegyptiſche Ciſtenroſe. W.
Habitat in Aegypto. *B. Juſſieus.* ☉ (v. v.)

Calycis *folia exteriora* 2 *parva; interiora* 3 *conniventia in veſicam ovatam inflatam, nec dehiſcentia ante maturitatem fructus.* Petala 5, *lanceolata lutea, breviſſima, incluſa.* Stamina 10, *parva.* Calycis *valvulae majores* 4-*nerviae: nervis elevatis ciliatis.*

* * * * * *Stipulati ſuffruticoſi.*

51. CISTUS *ſquamatus.*

C. ſuffruticoſus ſtipulatus, foliis obtectis ſquamis orbiculatis. *Sp. pl.* 743. *Loefl. it.* 67. *Cavan ic.* 2. *p.* 32. *t.* 139. *

C. humilis, floribus compactis in verticillos, minoris halimi foliis. *Barr. ic.* 328. *Bocc. muſ.* 2. *p.* 76. *t.* 64.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 229.

Schuppige Ciſtenroſe. W.

Habitat in Hiſpania *collibus aridis.* ♄ (v. ſ.)

Suffrutex. Ramis erectiuſculis, inferne tetragonis. Folia *ovali-lanceolata, craſſiuſcula, petiolata, oppoſita aut tria utrinque.* Stipulae *minutiſſimae, racemoſae, marceſcentes, ſeſſiles. Pedunculi teretes, conferti.* Planta *tota obtecta ſquamis orbiculatis, centro depreſſis.*

52. CISTUS *Lippii.*

C. ſuffruticoſus ſtipulatus erectus, foliis alternis oppoſitisque lanceolatis ſcabris, ſpicis ſecundis. *Mant.* 245. *Vahl. ſymb.* 1. *p.* 39.

C. (*ſtipulatus*) fruticoſus, foliis confertis tomentoſis ovalibus breviter petiolatis, margine revolutis. *Forſk. deſcr.* 100.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 230.

Lippiſche Ciſtenroſe. W.

Habitat in Aegypto. ♄ (v. ſ.)

Caulis *erectus, teres, pubeſcens, albicans, bifidus ſ. parce ramoſus.* Rami *infra bifurcaturam caulis, laterales, ſimplices, albi.* Folia *paſſim alterna paſſimque oppoſita, petiolata, elliptico-lanceolata, obtuſiuſcula, ſcabra, patentia, ſubtus caneſcentia.* Stipulae *anguſto-lanceolatae, erectae, longitudine petiolorum.* Racemi *oppoſitifolii, ſecundi, erecti ſ. inclinati.* Flores *ſeſſiles, conferti.* Calyx *ovatus: foliolis alternis, linearibus.* Petala *ovata, flava, vix calyce ma-*

jora. Stamina *ſaepius* 10, *petalis breviora.* Germen *pubeſcens.* Fructus *obtuſi, calyce tecti, in ſpica ſeſſiles, ſecundi, ſubrotundi.*

Planta *ſpontanea ramis foliisque utrinque caneſcentibus.* Folia *juniora axillarum revoluta.* Stipulae *ſetaceae.* Calyces *piloſi ſericei.* Vahl l. c.

*53. CISTUS *ſeſſiliflorus.*

C. ſuffruticoſus ſtipulatus erectus, foliis alternis oppoſitisque linearibus canis margine revolutis, ſpicis ſecundis. W.

C. (*ſeſſiliflorus*) fruticoſus ſtipulatus, foliis oppoſitis alternisque linearibus cinereis margine revolutis, racemis ſecundis, capſulis exſertis pubeſcentibus. *Desfont. fl. atlant.* 1. *p.* 427. *t.* 106. *

Aehrenblütige Ciſtenroſe. W.

Habitat in Africae *borealis collibus aridis prope* Maſcar. ♄

Frutex *pedalis vel bipedalis erectus ramoſiſſimus.* Folia *ſubpetiolata linearia obtuſiuſcula caneſcentia breviſſime tomentoſa, margine ſubtus revoluta, inferiora oppoſita, media et ſuperiora alterna.* Sipulae *parvae lineares.* Flores *ſeſſiles parvi racemoſi ſecundi.* Bracteae *minutae lineari-lanceolatae.* Calyx *pubeſcens.* Corolla *flava calyce paulo longior.* Capſula *ſubrotunda pubeſcens calyce longior.* Desf. l. c.

*54. CISTUS *ellipticus.*

C. ſuffruticoſus ſtipulatus erectus, foliis oppoſitis ellipticis canis margine revolutis, ſpicis ſecundis. W.

C. (*ellipticus*) ſtipulatus cinereus fruticoſus, foliis oppoſitis ellipticis, racemis ſecundis, floribus ſeſſilibus, capſulis exſertis. *Desfont. fl. atl.* 1. *p.* 418. *t.* 107. *

Elliptiſche Ciſtenroſe. W.

Habitat in Atlante *prope* Mayane Algeriae. ♄

Caulis *fruticoſus ramoſus erectus pedalis vel bipedalis.* Rami *pubeſcentes lanugine breviſſima.* Folia *oppoſita breviter petiolata elliptica margine ſubtus reflexa cinereo-candida, villis breviſſimis denſiſſimis ſtellatis utrinque obtecta.* Stipulae *quaternae parvae lineares.* Flores *racemoſi ſecundi parvi ſeſſiles.* Petala *flava calyce paullo longiora.* Capſula *ſubrotunda pubeſcens exſerta.* Desfont. l. c.

55. CISTUS *ſurrejanus.*

C. ſuffruticoſus procumbens ſtipulatus, foliis ovato-ob-

longis ſubpiloſis, petalis lanceolatis. *Sp. pl.* 743. *Hudſ. angl.* 233.
Helianthemum ſurrejanum. *Mill. dict. n.* 15.
Helianthemum vulgare, petalis florum peranguſtis. *Dill. elth.* 177. *t.* 145. *f.* 174.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 230.
Schmalblumige Ciſtenroſe. W.
Habitat in Angliae *comitatu* Surrejano *prope* Croydon. ♄

Petalis *anguſtis lanceolatis ab omnibus diſtinctus.* W.

*56. CISTUS *polyanthos.*

C. ſuffruticoſus ſtipulatus, foliis inferioribus ſubtus incanis, caulinis utrinque viridibus ciliatis calycibus hiſpidis, racemis paniculatis. *Desfont. fl. atl.* 1. *p.* 420. *t.* 108. *
Vielblumige Ciſtenroſe. W.
Habitat in Africae *borealis arenoſis prope* Mascar. ♄

Caules *ſuffruticoſi.* Rami *plures ex eodem caeſpite pedales teretes villoſi tuberculoſo aſperi.* Folia *oppoſita petiolata, inferiora ovata obtuſa minora ſubtus candida tomentoſa, media et ſuperiora ovato-oblonga ſeu lanceolata utrinque viridantia margine ciliata obtuſa, ſubtus venoſa.* Stipulae 4 *petiolatae lineari-lanceolatae obtuſiuſculae petiolo longiores.* Pedunculi *filiformes.* Flores *racemoſi, racemis paniculatis erectis ante floreſcentiam revolutis.* Flores *parvi pedicellati, pedicellis capillaribus.* Bracteae *lineares pedicello breviores.* Calyx *villis albidis mollibus numeroſis patentibus hirſutus.* Corolla *flava calyce longior.* Capſula *parva apice villoſa.* Desf. l. c.

*57. CISTUS *glaucus.*

C. ſuffruticoſus ſtipulatus adſcendens, foliis tomentoſis glaucis margine revolutis, inferioribus ovatis, ſummis lanceolatis, racemis ſecundis. W.
C. (*glaucus*) caule ſuffruticoſo ſtipulato, foliis ovatis glaucis margine revolutis, floribus racemoſis. *Cavan. ic.* 3. *p.* 31. *t.* 261. *
Meergrüne Ciſtenroſe. W.
Habitat in Hiſpania. ♄

Petala *flava calyce majora ſubrotunda, margine crenulata.* W.

58. CISTUS *nummularius.*

C. ſuffruticoſus ſtipulatus, foliis inferioribus orbiculatis,

ſuperioribus ovatis. *Sp. pl.* 743. *Cavan. ic.* 2. *p.* 34. *t.* 142. * *Desfont. fl. atl.* 1. *p.* 423. *

C. humilis ſ. Chamae-Ciſtus nummulariae folio. *Magn. monſ.* 293.

Helianthemum nummularium. *Mill. dict. n.* 12.

Helianthemum ad nummulariam accedens. *Bauh. hiſt.* 2. *p.* 20.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 231.

Rundblättrige Ciſtenroſe. W.

Habitat Monſpelii, *inque* Hiſpania *et* Africa *boreali.* ♄

*59. CISTUS *canarienſis.*

C. ſuffruticoſus ſtipulatus procumbens, foliis oppoſitis alternisque pubeſcentibus glaucis, oblongo-obovatis acutis, racemis ſecundis. W.

C. (*canarienſis*) ſuffruticoſus ſtipulatus procumbens, foliis ſubovatis alternis et oppoſitis, racemis erectis. *Iacq. ic.* 1. *t.* 97. (*ſed foliorum color male expreſſus.*) *Iacq. miſc.* 2. *p.* 339. *

Canariſche Ciſtenroſe. W.

Habitat in Canariis *in* Fuerta ventura. ♄ (v. ſ.)

Pili *parvi ſtellati denſe tegunt utramque paginam foliorum, inde toto planta glauca et folia juniora mollia.* W.

60. CISTUS *ſerpillifolius.*

C. ſuffruticoſus ſtipulatus, foliis oblongis, calycibus laevibus. *Sp. pl.* 743. *Sauv. monſp.* 148.

Chamae-Ciſtus repens ſerpillifolia lutea. *Bauh. pin.* 466.

Chamae-Ciſtus ſerpilli folio nigricante et hirſuto, flore aureo odorato. *Bauh. hiſt.* 2. *p.* 17.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 231.

Quendelblättrige Ciſtenroſe. W.

Habitat in alpibus Europae *auſtralioris.* ♄

*61. CISTUS *violaceus.*

C. ſuffruticoſus ſtipulatus adſcendens, foliis oppoſitis ſubtomentoſis linearibus obtuſis baſi attenuatis margine revolutis, calycibus laevibus. W.

C. (*violaceus*) ſuffruticoſus ſtipulatus foliis oppoſitis linearibus ſubtomentoſis, calycibus laevibus violaceis. *Cavan. ic.* 2. *p.* 38. *t.* 147. *

Violette Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Racemi *longi erecti.* Calyx *laevis violaceo-ruber.* Petala *alba.* W.

*62. CISTUS *linearis.*

C. ſuffruticoſus ſtipulatus adſcendens ſubtomentoſus, foliis linearibus obtuſis petiolatis margine revolutis, calycibus laevibus. W.

C. (*linearis*) ſuffruticoſus ſtipulatus ſubtomentoſus, foliis linearibus, floribus racemoſis. *Cavan. ic.* 3. *p.* 8. *t.* 216. *

Linienblättrige Ciſtenroſe. W.

Habitat in Hiſpania. ♄

Calyx *laevis ſtriatus violaceus.* Petala *alba. Valde affinis praecedenti, differt vero:* Foliis *evidenter petiolatis baſi non attenuatis triplo longioribus uncialibus,* Calycinis *foliolis binis minoribus acutioribus nec obtuſis,* Petalis *obovatis et toto habitu.* W.

*63. CISTUS *laevis.*

C. ſuffruticoſus ſtipulatus erectus, foliis linearibus ſeſſilibus glabris margine revolutis carinatis, calycibus laevibus. W.

C. (*laevis*) ſuffruticoſus ſtipulatus laevis, foliis linearibus brevibus oppoſitis, margine revoluto. *Cavan. ic.* 2. *p.* 35. *t.* 145. *f.* 1. *

Glatte Ciſtenroſe. W.

Habitat in collibus Hiſpaniae. ♄

Caulis *erectus.* Folia *linearia ſeſſilia glabra, margine revoluta, coſta media inferne protuberante, apice acuta, inferiora conferta breviora, ſuperiora patentiuſcula.* Stipulae *lineares.* Calycis *foliola bina breviora ſubulata.* Corolla *calyce major ſaturate flava.* W.

*64. CISTUS *ſtrictus.*

C. ſuffruticoſus ſtipulatus erectus, foliis canis lineari-ſubulatis margine revolutis, racemis ſecundis, calycibus glabris. W.

C. (*ſtrictus*) ſtipulatus, caule ſuffruticoſo erecto, ramis oppoſitis ſtrictis, foliis ſubulatis canis. *Cavan. ic.* 3. *p.* 32. *t.* 263. *f.* 2. *

Steife Ciſtenroſe. W.

Habitat in Hiſpania *in monte* Palomera. ♄

Caulis *ſuffruticoſus erectus ſemipedalis, ramis pluribus oppoſitis ſtrictis, tomento brevi cano obductis.* Folia *oppoſita ſubſeſſilia anguſtiſſima, limboque ita re-*

voluto ut ſubulata appareant, caneſcentia, apice pilis paucis terminata. Stipulae *ſubulatae ſubrubentes.* Calyx *glaber ſtriatus.* Corolla *alba calyce major.* Cavan. l. c.

65. CISTUS *glutinoſus.*

C. ſuffruticoſus ſtipulatus, foliis linearibus oppoſitis alterniſque, pedunculis villoſis glutinoſis. *Mant.* 246. *Ger. prov.* 394.

C. ſuffruticoſus ſtipulatus, foliis minimis obovatis, limbo revoluto, pedunculis villoſis glutinoſis, capſulis dehiſcendo patulis. *Cavan. ic.* 2. *p.* 36. *t.* 145. *f.* 2. *

Chamae-Ciſtus incanus, tragorigani folio hiſpanicus. *Barr. rar.* 513. *t.* 415.

Helianthemum thymi folio incano. *Bauh. hiſt.* 2. *p.* 19.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 232.

Klebrige Ciſtenroſe. W.

Habitat in Europa *auſtrali.* ♄

Caulis *ſuperne pubeſcenti-viſcidus.* Folia *ſuperiora ſemper alterna magisque villoſa.* Racemus *pauciflorus, villoſo-glutinoſus.* Petala *parva, lutea.*

Petala *calyce minora.* W.

66. CISTUS *thymifolius.*

C. ſuffruticoſus ſtipulatus procumbens, foliis linearibus oppoſitis breviſſimis congeſtis. *Sp. pl.* 743. *Sauv. monſp.* 148.

Chamae-Ciſtus luteus, thymi folio, oliganthes. *Barr. ic.* 444.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 232.

Thymianblättrige Ciſtenroſe. W.

Habitat in G. Narbonenſi, Hiſpania. ♄ (v. ſ.)

67. CISTUS *piloſus.*

C. ſuffruticoſus ſtipulatus erectiuſculus, foliis linearibus ſubtus biſulcatis incanis, calycibus laevibus. *Sp. pl.* 744. *Sauv. monſp.* 147. *Mill. dict. n.* 1. *Allion. ped n.* 1672. *t.* 45. *f.* 1. 2.

C. ſuffruticoſus erectus, foliis majoribus petiolatis revolutis ſubtus incanis, caeteris linearibus. *Ger. prov.* 395.

Chamae Ciſtus foliis thymi incanis. *Bauh. pin.* 466.

Chamae-Ciſtus 4. *Cluſ. hiſt.* 1. *p* 74.

β. Ciſtus ſtipulis quaternis, foliis lineari-ovalibus incanis, calycibus tomentoſis. *Sauv. monſp.* 148.

Helianthemum flore albo, folio anguſto hirſuto. *Bauh. hiſt.* 2. *p.* 17.

γ. Ciſtus foliis villoſis lanceolatis, axillis folioſis, ſtipulis ſubulatis. *Sauv. monſp.* 148.

Helianthemum ſampſucifolium. *Mill. dict. n.* 7.

Helianthemum ſ. Ciſtus humilis, folio ſampſuci, capitulis valde hirſutis. *Bauh. hiſt.* 2. *p.* 20.

δ. Ciſtus foliis minoribus et incanis. *Bauh. pin.* 466. *Ger.*

C. omnino candicans, flore nunc albo nunc pallido. *Bauh. hiſt.* 2. *p.* 16.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 232.

Behaarte Ciſtenroſe. W.

Habitat in Gallia. ♄ (v. ſ.)

Caules *erectiuſculi.* Bracteae *ſolitariae, ad latera pedicellorum. Gouan.* Flores *albi.*

Ad hanc ſpeciem facile omnes ſpecies parvas albo flore, etiam C. appenninum refert Gerard.

*68. CISTUS *lavandulifolius.*

C. ſuffruticoſus ſtipulatus, foliis lanceolato-linearibus tomentoſis, calycibus racemoſis tomentoſis ſecundis pendulis. *Vahl. ſymb.* 1. *p.* 39. *

C. (*lavandulaefolius*) ſuffruticoſus erectus ſtipulatus, foliis lanceolatis margine revolutis ſubincanis, racemis incurvis terminalibus, floribus confertis. *Lamarck. encycl.* 2. *p.* 25. * *Desfont. fl. atl.* 1. *p.* 417. *

C. (*ſyriacus*) ſuffruticoſus exſtipulatus erectus, foliis lanceolatis ad oras revolutis, floribus racemoſis. *Iacq. ic. rar.* 1. *t.* 96. *Collect.* 1. *p.* 98. *

C. folio lavandulae Cluſii forte idem cum Helianthemo recto. *Bauh. hiſt.* 2. *p.* 5.

C. folio ſpicae. *Bauh pin.* 465.

C. lavendulae latifoliae facie. *Barrel. ic.* 288.

C. folio lavandulae. *Cluſ. hiſt.* 1. *p.* 72.

β. Ciſtus lavandulaefolius. *Iacq. Collect. ſuppl. p.* 115. *t.* 10. *f.* 2 *

Lavendelblättrige Ciſtenroſe. W.

Habitat in Hiſpaniae, Maſſiliae, Tuneti *et* Syriae *collibus aridis.* ♄ (v. ſ. ſ. fl.)

69. CISTUS *racemoſus.*

C. ſuffruticoſus ſtipulatus, foliis linearibus, calycibus racemoſis ſecundis nervoſo-angulatis glabris. *Vahl ſymb.* 1. *p.* 39. *

C. ſuffruticoſus ſtipulatus, foliis lanceolato-linearibus ſubtus tomentoſis. *Mant.* 76. *Cavan. ic.* 2. *p.* 33. *t.* 140. *

C. lavandulae folio, thyrſoides. *Barr. rar. ic.* 293. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 233.

Traubige Ciſtenroſe. W.

Habitat in Hiſpania. ♄ (v. ſ.)

Caulis *ſtatura Rosmarini, vix tomentoſus.* Folia *linearia ſ. anguſto-lanceolata, ſubtus leviſſime tomentoſa, longiuſcula.* Stipulae *lineares.* Racemi *longiſſimi, terminales, ſecundi.* Calyces *angulati, laeves.* *Differt a C. lavandulifolio:* foliis *anguſtioribus brevioribus minus tomentoſis absque mucrone in apice, nec faſciculo foliorum in axillis,* calycibus *minoribus glabris ex nervis elevatis angulatis.* Vahl l. c.

*70. CISTUS *ciliatus.*

C. ſuffruticoſus procumbens ſtipulatus, ramis tomentoſis, foliis anguſto-lanceolatis villoſis, calycibus membranaceis, angulis ciliatis. *Desfont. fl. atl.* 1. *p.* 421. *t.* 109. *

Gefranzte Ciſtenroſe. W.

Habitat in collibus aridis arenoſis Africae *borealis prope* Caſlam. ♄

Caules *ſuffruticoſi decumbentes baſi ramoſi pedales.* Rami *tomentoſi incani teretes ſimplices.* Folia *oppoſita breviſſime petiolata anguſto-lanceolata, ſuperne hirſuta, ſubtus caneſcentia tomentoſa margine revoluta.* Stipulae 4 *lineares petiolo longiores.* Flores *racemoſi terminales ante floreſcentiam revoluti.* Bracteae *lineari-lanceolatae.* Calyx *anguloſus, angulis elevatis obliquis fuſcis glanduloſis piloſis, foliola* 2 *exteriora minuta linearia.* Corolla *roſea paullo major quam in C. Helianthemo.* Capſula *ſubrotunda calyce tecta.* Desfont. l. c.

*71. CISTUS *anguſtifolius.*

C. ſuffruticoſus ſtipulatus diffuſus, foliis lanceolatis, calycibus hirſutis. *Murr. ſyſt. veg. ed.* 14. *p.* 500. *Jacq. hort.* 3. *t.* 53.

Schmalblättrige Ciſtenroſe. W.

Habitat - - - - ♄ (v. ſ.)

Folia *oppoſita breviter petiolata lanceolata acutiuſcula aſpera.* Petala *flava ad baſin fulva.* Capſula *ovata hirſuta.* W.

72. CISTUS *Helianthemum.*

C. ſuffruticoſus procumbens, ſtipulis lanceolatis, foliis oblongis revolutis ſubpiloſis. *Sp. pl.* 744. *Scop. carn. n.* 649. *Pollich. pal. n.* 511. *Hoffm. germ.* 185. *Fl. dan. t.* 101. *Kniph. cent.* 12. *n.* 27. *Roth. germ. I.* 229. *II.* 586.

C. ſtipulis quaternis, foliis oblongis utrinque nudis, caule procumbente. *Hort. cliff.* 206. *Fl. ſuec.* 433, 472.

C. foliis conjugatis, ellipticis hirſutis, integerrimis, petiolis unifloris ſubhirſutis. *Hall. helv. n* 1033.

C. ſtipulatus, foliis ſubhirſutis, ellipticis, obtuſe acuminatis. *Crantz. auſtr. p.* 100.

Helianthemum Chamae-Ciſtus. *Mill. dict. n.* 1.

Helianthemum germanicum. *Tabern.* 1062.

Chamae-Ciſtus vulgaris, flore luteo. *Bauh. pin.* 465. *Loeſ. pruſſ* 43. *t.* 8.

Flos ſolis, Panax chironium. *Cam. epit.* 501.

β. Ciſtus grandiflorus. *Scop. carn. n.* 648. *t.* 25. *

γ. flore albo. *Ait. Kew.* 2. *p.* 239.

δ. flore rubro.

C. (*roſeus*) ſuffruticoſus exſtipulatus procumbens, foliis oppoſitis oblongis ad oras revolutis, utrinque ſubincanis. *Murr. ſyſt. ed.* 14. *p.* 498. *Iacq. hort.* 3. *t.* 65. *Allion. ped. n.* 1675. *t.* 45. *f.* 4.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 234.

Gemeine Ciſtenroſe. W.

Habitat in Europae *paſcuis ſiccis.* ♄ (v. v.)

Petala ſuborbiculata, integerrima, flava, annulo receptaculum ambiente ſaepius fulvo. Racemus *ante floreſcentiam nutans.* Folia *pilis raris adſperſa.*

Variat valde, altitudine caulis, foliis glabris et pubeſcentibus, atque floris colore. W.

*73. CISTUS *mutabilis.*

C. ſuffruticoſus ſtipulatus procumbens, ſtipulis lanceolatis, foliis oblongis glabris planis. *Iacq. ic.* 1. *t.* 99. *Miſc.* 2. *p.* 340. *

Veränderliche Ciſtenroſe. W.

Habitat - - - - ♄ (v. v.)

Differt a praecedente: foliis *planis,* caule *altiore et quod frigore deſtruatur. Variat corollis pallide flavis et roſeis.* W.

*74. CISTUS *foetidus.*

C. ſuffruticoſus ſtipulatus procumbens, ſtipulis lanceolatis, foliis oblongis hirſutis ſcabris. *Iacq. ic.* 1. *t.* 98. *Miſc.* 2. *p.* 341. *

Stinkende Ciſtenroſe. W.

Habitat - - - - ♄ (v. v.)

Odor totius plantae foetidus Bryoniae. Corolla *C. Helianthemi alba.* W.

*75. CISTUS *croceus.*

C. ſuffruticoſus ſtipulatus pubeſcens, villis breviſſimis ſtellatis, foliis ellipticis obtuſis. *Desfont. fl. atl.* 1. *p.* 422. *t.* 110. *

Ciſtus humilis. *Cluſ. hiſp.* 151.

Helianthemum fruteſcens folio Majoranae incano. *Tournef. inſt.* 249.

Majoranblättrige Ciſtenroſe. W.

Habitat in Hiſpania *et* Atlante *prope* Tlemſen. ♄ (v. ſ.)

Frutex *pedalis.* Rami *plures ex eodem ceſpite ſimplices erecti teretes tomentoſi caneſcentes.* Folia *oppoſita petiolata, inferne caneſcentia, margine revoluta, utrinque pubeſcentia, villis confertiſſimis breviſſimis ſtellatis, inferiora minora ſubrotunda, media elliptica obtuſa, ſuperiora lanceolata acutiuſcula.* Stipulae 4 *ſubulatae petiolo paullo longiores.* Racemi *ante floreſcentiam convoluti.* Bracteae *lanceolatae pubeſcentes longitudine pedicellorum.* Calyx *pubeſcens anguloſus flaveſcens.* Corolla *magnitudine C. Helianthemi crocea.* Desfont. l. c.

Affinis C. glauco, ſed diſtinctus, foliorum *forma,* caule *duro fruteſcente,* petalis *croceis integerrimis.* W.

76. CISTUS *hirtus.*

C. ſuffruticoſus ſtipulatus, foliis ovatis, calycibus hiſpidis. *Sp. pl.* 744. *Kniph. cent.* 12. *n.* 28. *Cavan. ic.* 2. *p.* 37. *t.* 146. *

C. foliis lanceolato-linearibus integerrimis ſubtus incanis. *Gron. orient.* 196.

C. foliis oblongo-linearibus margine revolutis. *Le Monier. obſ.* 229. *Sauv. monſp.* 148.

Ciſtus Ledon roriſmarini foliis ſubtus incanis. *Bauh. pin.* 467.

Helianthemum, anguſto ſerpilli folio, villoſum, flore aureo, italicum. *Barrel. rar.* 411. *t.* 488.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 235.
Rauhkelchige Ciſtenroſe. W.
Habitat in Hiſpania, Narbona. ♄ (v. ſ.)

77. CISTUS *apenninus.*
C. ſuffruticoſus ſtipulatus patulus, foliis lanceolatis hirtis. *Sp. pl.* 744.
Helianthemum germanicum et apenninum. *Mill. dict. n.* 2. *et* 4.
Helianthemum album germanicum. *Tabern. icon.* 1062.
Helianthemum ſaxatile, foliis et caulibus incanis oblongis, floribus albis. *Mentz. pug.* 8. *f.* 3. *Dill. elth.* 170.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 236.
Apenniniſche Ciſtenroſe. W.
Habitat in Apenninis, Italiae *montibus.* ♄
Folia *ſupra viridia hirta, ſubtus incana.* Flores *albi.* Frutex *pedalis ramoſo-patulus.*

78. CISTUS *polifolius.*
C. ſuffruticoſus ſtipulatus procumbens, foliis oblongo-ovatis incanis, calycibus laevibus, petalis ſerratis. *Sp. pl.* 745. *Hudſ. angl.* 234.
C. humilis alpinus durior, polii noſtratis folio candicante. *Pluk. alm.* 107. *t.* 23. *f.* 6.
Chamae-Ciſtus montanus, polii folio. *Raj. angl.* 4. *p.* 274. *t.* 274. *f.* 2.
Helianthemum montanum, polii folio incano, flore candido. *Dill. elth.* 175. *t.* 145. *f.* 172.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 236.
Poleyblättrige Ciſtenroſe. W.
Habitat in aeſtuario Sabrinae Anglorum *ad* Brentdown. ♄ (v. ſ.)
Flores *albi.*

79. CISTUS *arabicus.*
C. ſuffruticoſus ſtipulatus procumbens, foliis linearibus pedunculorum alternis, ramulorum confertis. *Vahl. ſymb.* 2. *p.* 62. *t.* 55.
C. ſuffruticoſus ſtipulatus, foliis alternis lanceolatis planis laevibus. *Sp. pl.* 745. *Amoen. acad.* 4. *p.* 275. *Cent.* 39. *Desf. fl. atl.* 1. *p.* 419.*
Helianthemum creticum linariae folio, flore croceo. *Tournef. cor.* 18.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 237.

Arabiſche Ciſtenroſe. W.
Habitat in Arabia, Creta *et* Africa *boreali.* ♄ (v. ſ.)

Rami *aphylli procumbentes demum adſcendentes, ſaepe pedales teretes glabri tenues.* Ramuli *a baſi usque copioſi pollicares alterni diſtantes patentiſſimi, inferiores ſteriles, ſupremi floriferi ſimpliciſſimi elongati cinerei pubeſcentes.* Folia *in ramulis ſterilibus conferta bilinearia declinata linearia rigidiuſcula avenia obtuſiuſcula, ultima tomento ſubcinerea.* Stipulae *minutae ovatae.* Folia *ramorum floriferorum latiora et longiora pubeſcentia.* Stipulae *lanceolatae.* Racemus *terminalis 3-ſ. 4-florus ebracteatus.* Pedicelli *diſtantes piloſo-ſubviſcoſi verſus apicem, ſub flore parum craſſiores.* Calyx *piloſus ſubviſcoſus, foliola interiora trinervia, interſtitiis membranaceis, exteriora lanceolata.* Vahl l. c.

Ciſtorum hiſtoria maxime omnium obſcura ob varietatum copiam, illuſtrabitur ſi determinent autoptae in ſingulis:

1. *Caudicem arboreſcentem, fruticuloſum, annuum ſ. perennem.*
2. *Caulem erectum ſ. decumbentem.*
3. *Folia oppoſita ſ. alterna: cum figura eorundem.*
4. *Stipulas binas ſ. nullas.*
5. *Pedunculum uniflorum ſ. multiflorum: nudum ſ. bracteatum.*
6. *Petalorum figuram.*
7. *Capſulas 5-loculares ſ. 3-valves.*
8. *Calycem aequalem ſ. inaequalem.*

1049. VALLEA.

Gen. plant. edit. Schreb. n. 914.

Cal. 3-ſ. 4-phyllus. *Petala* 4 ſ. 5 trifida. *Stigma* 4-ſ. 5-fidum. *Capſ.* 2-locularis polyſperma.

1. VALLEA *ſtipularis.*

V. ſtipularis. *Lin. ſuppl.* 266. *Mutis amer. v. 7. t.* 10.
Herzblättrige Vallea. W.
Habitat in Nova Granada *ſub temperie frigida.* ♄

Arbor *biorgyalis.* Folia *alterna petiolata cordata integerrima ſubtus ſubtomentoſa.* Petioli *foliis brevio-*

res. Stipulae *ſeſſiles reniformes amplexicaules integerrimae majuſculae.* Panicula *trifaria ramorum terminalis.* Flores *ſanguinei.*

Numerus partium fructificationis communiter quaternarius, variat quinario.

*1050. BONNETIA.

Gen. pl. ed. Schreb. n. 915.

Cal. 5-phyllus. *Cor.* 5-petala. *Capſ.* 3-locularis 3-valvis polyſperma.

*1. BONNETIA *paluſtris.*

B. paluſtris. *Vahl eclog.* 2. *p.* 42. *

Mahurea paluſtris. *Aubl. guj.* 1. *p.* 558. *t.* 222. *

Stumpfe Bonnetie. W.

Habitat in locis paludoſis Cajennae, Gujanae. ♄

Arbor *quindecimpedalis.* Folia *alterna petiolata elliptica obtuſa integerrima reticulato-venoſa glabra.* Racemus *terminalis.* Pedunculi *ſubgeminati.* Flores *purpurei.* W.

1051. PROCKIA.

Gen. plant. edit. Schreb. n. 916. *et* 1609.

Cal. 3-phyllus praeter foliola 2 baſeos. *Cor.* 0. *Bacca* 5-angularis, polyſperma.

1. PROCKIA *Crucis.*

P. foliis cordato-ovatis dentatis, pedunculis terminalibus ſubracemoſis. W.

P. crucis. *Sp. pl.* 745. * *Vahl. ſymb.* 3. *p.* 69. *t.* 64. *.

Prockia. *Brown*

Houttuyn Lin. Pfl. Syſt. 4. *p.* 34.

Weſtindiſche Prockie. W.

Habitat in inſula St. Crucis. ♄

Frutex *Ramis teretibus.* Folia *alterna, petiolata, ovata, acuminata, ſerrata, glabra.* Stipulae *lineares.* Flores *nonnulli, terminales, alterni, pedicellati: Bracteis linearibus.*

*2. PROCKIA *ſerrata.*

P. foliis oblongo-ovatis ſerratis acuminatis, pedunculis lateralibus aggregatis unifloris.

Lightfootia ſerrata. *Swartz prod.* 83. *Vahl ſymb.* 3 *p.* 69.
Sägeblättrige Prockie. W.
Habitat in inſula Montſerrat. ♄ (v. ſ.)

*3. PROCKIA *theaeformis.*

P. foliis lanceolato - ellipticis ſerratis obtuſiuſculis, pedunculis axillaribus ſubſolitariis unifloris.
Lightfootia theaeformis. *Vahl ſymb.* 3. *p.* 69. *
Theeförmige Prockie. W.
Habitat in inſula Borboniae. ♄

*4. PROCKIA *integrifolia.*

P. foliis ſubcoriaceis oblongis obovatisque emarginatis ſubintegerrimis, pedunculis ſubaggregatis unifloris.
Lightfootia integrifolia. *Vahl ſymb.* 3. *p.* 70.*
Ganzblättrige Prockie. W.
Habitat in inſula Mauritii. ♄

Lightfootia Swartzii a Prockia non differt, niſi ſtigmate ſeſſili; nota haud ſufficiens pro charactere generico. W.

1052. CORCHORUS.

Gen. pl. ed. Schreb. n. 917.

Cor. 5-petala. *Cal.* 5-phyllus, deciduus. *Capſ.* plurivalvis, loculamentoſa.

1. CORCHORUS *olitorius.*

C. capſulis oblongis ventricoſis, foliorum ſerraturis infimis ſetaceis. *Fl. zeyl.* 213. *Hort. upſ.* 147. *Gron. orient.* 170. *Mill. dict. n.* 1.
C. foliorum infimis ſerraturis maximis reflexis *Hort. cliff.* 209. *Roy. lugdb.* 478.
C plinii. *Bauh. hiſt.* 317.
Corchorus. *Comm. hort.* 47. *t.* 12.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 238. *t.* 54. *f.* 1.
Gemüſeartiger Corchorus. W.
Habitat in Aſia, Africa, America. ⊙ (v. v.)

Caulis *laevis, viridis.* Stipulae *ſimplices baſi rubrae.* Folia *trinervia, ovata dentibus ſetaceis.* Capſulae *fuſiformes, 5-loculares, ſulcatae, ſubquinque dentatae ſ. obtuſae.*

2. CORCHORUS *trilocularis.*

C. capſulis trilocularibus trivalvibus triquetris: angulis bifidis ſcabris, foliis oblongis: ſerraturis infimis ſetaceis. *Mant.* 77. *Iacq. hort.* 173. *Vahl ſymb.* 2. *p.* 63.
C. aeſtuans. *Forsk. deſcr.* 101.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 240.
Dreyfächriger Corchorus. W.
Habitat in Arabia. *Forskaehl.* ⊙ (v. v.)

Caules *laeves, erecti, pedales, teretes, virides.* Stipulae *ſetaceae, parvae.* Folia *alterna, petiolata, lineata, nuda, ſubtus ſcabra, lanceolata, undato-ſerrata dentibus ſetaceis.* Pedunculi *ſuboppoſitifolii, breves, bifidi ſ. biflori.* Calyces *angulati.* Petala *anguſta, lutea.* Capſulae *lineares, triquetro-priſmaticae, ſcabrae, angulis linea ſtriatis, apice obtuſae.*

3. CORCHORUS *tridens.*

C. capſulis linearibus teretiuſculis ſcabris, foliorum ſerraturis infimis ſetaceis. *Mant.* 566.
C. americanus, anguſto barbato folio. *Pluk. phyt. t.* 127. *f.* 4. *Burm. ind.* 123. *t.* 37. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 240.
Dreyzähnige Corchorus. W.
Habitat in India.

Caulis *laevis, viridis.* Stipulae *tripartitae, ſetaceae.* Folia *lineata, lanceolata, undato-ſerrata dentibus ſetaceis.* Capſulae *lineares, ſcabrae.* Styli 3, *divaricatiſſimi, bifidi.*

4. CORCHORUS *aeſtuans.*

C. capſulis oblongis trilocularibus trivalvis ſexſulcatis ſexcuſpidatis, foliis cordatis: ſerraturis infimis ſetaceis. *Mant.* 565. *Iacq. hort. t.* 85. *Mill. dict. n.* 2.
C. capſulis oblongis ſexſulcatis ſexcuſpidatis foliis, cordatis infimis ſerraturis ſetaceis. *Loefl. it.* 224. *n.* 112. *β. Sp. pl.* 2. *p.* 746.
Alcea cibaria ſ. Corchorus americana, carpini foliis, ſextuplici capſula longiore. *Pluk. phyt.* 127. *f.* 3.
Triumfetta ſubvilloſa, foliis rotundioribus undulatis atque dentatis, proſtremis in ſetas inermes. *Brown. jam.* 232. *t.* 25. *f.* 1.
Houttuyn Lin. Pfl. Syſt. 6. *p.* 241.
Heiſser Corchorus. W.
Habitat in America *calidiore.* ♄

Caulis *ſcabriuſculus, purpuraſcens, ramis divaricatis.* Stipulae *ſetaceae.* Folia *trinervia, cordato-ovata: dentibus ſetaceis.* Capſulae *lineares, 6-angulares, angulis compreſſis: paribus approximatis.* Styli 3, *bifidi.* Capſulae *dentes terminales divaricatiſſimi.*

*5. CORCHORUS *acutangulus.*

C. capſulis priſmatico-cuneatis acutangulis tridentatis, foliis ovatis baſi ſubuniſetoſis, petiolis hiſpidis. *Lamarck. encycl.* 2. *p.* 104. *

Lyſimachia non pappoſa meliſſophylla maderaſpatenſis. *Pluk. phytog. t.* 44. *f.* 1.

Scharfeckiger Corchorus. W.

Habitat in India *orientali.* ♄ (v. ſ.)

Caulis *ramoſus ut petioli hiſpidus.* Folia *alterna longe petiolata ovata obtuſa dentata, dente infimo ad unum latus tantum quandoque acuminato ſetaceo, trinervia venoſa ſcabriuſcula.* Stipulae 2 *ſetaceae.* Flores *flavi parvi plerumque geminati axillares.* Capſulae *baſi attenuatae 5-angulares angulis binis acutioribus et magis prominentibus.* W.

*6. CORCHORUS *faſcicularis.*

C. capſulis oblongis ſubſeſſilibus piloſis mucronatis ſubgeminis, foliis lanceolato-oblongis glabris, ſerraturis aequalibus. *Vahl. ſymb.* 3. *p.* 69.

C. (*faſcicularis*) capſulis conoideis lanatis faſciculatis ſubſeſſilibus, foliis oblongo-ellipticis petiolatis baſi muticis. *Lamarck. encycl.* 2. *p.* 105. *

Euphraſiae affinis indiana meliſſae turcicae foliis alternis, capſulis pubeſcentibus trifariam junctis. *Pluk. amalth.* 85. *t.* 439. *f.* 6.

Büſchliger Corchorus. W.

Habitat in India *orientali.* ♄

Rami *teretes glabri.* Folia *petiolata oblonga anguſta, utrinque glabra pollicaria obtuſa, ſerraturis infimis non ſetaceis.* Stipulae *ut in congeneribus.* Fructus *gemini vel terni ſubſeſſiles ad latus folii, vix unguiculares, piloſo-cinerei trivalves.* Vahl l. c.

7. CORCHORUS *capſularis.*

C. capſulis ſobrotundis depreſſis rugoſis, foliorum ſerraturis infimis ſetaceis. *Fl. zeyl.* 214. * *Mill. dict. n.* 3. *Kniph. cent.* 4. *n.* 21.

C. foliorum infimis ſerraturis minoribus. *Hort. cliff.* 210. *Roy. lugdb.* 478.

Alcea olitoria ſ. Corchorus americanus, praelongis foliis, capſula ſtriata ſubrotunda brevi. *Pluk. alm.* 18. *t.* 255. *f.* 4.
Gania ſativa. *Rumph. amb.* 5. *p.* 212. *t.* 78. *f.* 1.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 242.
Rundfrüchtiger Corchorus. W.
Habitat in India. ⊙

Calyx *foliolis concavis, incurvatis, extus punctatis, corolla brevioribus.* Petala *emarginata.*

*8. CORCHORUS *ſcandens.*

C. foliis ovatis ſetaceo-ſerratis oppoſitis, caule ramisque flexuoſo-ſcandentibus. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 335.
Kletternder Corchorus. W.
Habitat in Japonia.

Caulis *teres ſcandens ramoſus.* Rami *oppoſiti ſimiles divaricati.* Folia *oppoſita breviſſime petiolata, baſi rotundata ovata acuminata ſerrata, ſerraturis ſetigeris, pollicaria.* Flos *in ramulis terminalis ſolitarius flavus.* Thunb. l. c.

*9. CORCHORUS *ſerratus.*

C. foliis oblongis ſerratis cuſpidatis, ramis glabris. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 335.
Geſägter Corchorus. W.
Habitat in Japonia.

Caulis *erectus glaber ramoſus.* Rami *teretes purpurei glabri erecti. Folia alterna petiolata oblonga ſerrata, ſerraturis magnis apice ſetaceis, cuſpidata, ſupra tenuiſſime villoſo-ſcabra, ſubtus glabra bipollicaria patula.* Thunb. l. c.

10. CORCHORUS *hirſutus.*

C. capſulis ſubrotundis lanatis, foliis ovatis obtuſis tomentoſis aequaliter ſerratis. *Sp. pl.* 747. *Mill. dict. n.* 8. *Jacq. amer.* 165.
C. fruteſcens, foliis ellipticis tomentoſis crenatis, capſulis ovato-oblongis lanuginoſis umbellatis. *Lamarck. encycl.* 2. *p.* 105. *
C. americanus lanuginoſus, folio chamaedryos, ſiliqua tricapſulari lanuginoſa et ſubaſpera. *Breyn. prod.* 3. *p.* 56.
Guazuma frutex chamaedryfolia, fructu lanuginoſo, major et minor. *Plum. gen.* 36. *ic.* 104.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 242.
Zottiger Corchorus. W.
Habitat in America *meridionali.* ♄ (v. ſ.)

*11. CORCHORUS *japonicus.*

C. capſulis rotundis glabris, foliis duplicato-ſerratis. *Thunb. jap.* 227. *
α. flore ſimplici.
Teito vulgo Jamma Buki. *Kaempf. amoen.* 844.
β. flore pleno.
Jamma Buki altera. *Kaempf. amoen.* 845.
Corchorus flore pleno e Japonia. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 245. *t.* 45. *f.* 2.
Japaniſcher Corchorus. W.
Habitat in Japonia *juxta* Nagaſaki. ♄

*12. CORCHORUS *flexuoſus.*

C. foliis duplicato-ſerratis cuſpidatis, caule flexuoſo. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 335.
Geknieter Corchorus. W.
Habitat in Japonia.

Caulis *teres a foliis decurrentibus ſubangulatus flexuoſus glaber erectus bipedalis.* Folia *alterna petiolata oblique cordata ovata cuſpidata duplicato-ſerrata villoſa patula bipollicaria.* Flores *in ramis terminales lutei.* Thunb. l. c.

13. CORCHORUS *hirtus.*

C. capſulis oblongis cauleque piloſis, foliis oblongis aequaliter ſerratis. *Sp. pl.* 747.
C. folio ulmi, major. *Plum. ſpec.* 7. *ic.* 303. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 243.
Rauher Corchorus. W.
Habitat in America *meridionali.*

14. CORCHORUS *ſiliquoſus.*

C. capſulis linearibus compreſſis bivalvibus, foliis lanceolatis aequaliter ſerratis. *Mill. dict. n.* 7. *Kniph. cent.* 4. *n.* 22. *Swartz. obſ.* 219. * *Jacq. hort.* 3. *p.* 34. *t.* 59.
C. folio ulmi minor. *Plum. ſpec.* 7. *ic.* 103. *f.* 1.
C. Coreta. *Amoen. acad.* 5. *p.* 398. *
Corchoro affinis chamaedryos folio, flore ſtamineo, ſeminibus atris quadrangulis duplici ſerie diſpoſitis. *Sloan. jam.* 50. *hiſt.* 1. *p.* 145. *t.* 94. *f.* 1. *Raj. ſuppl.* 516.

Coreta foliis minoribus ovatis crenatis, floribus ſingularibus. *Brown. jam.* 147.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 244.
Schotenbringender Corchorus W.
Habitat in America *meridionali.* ♃ (v. ſ.)

Caulis *fere fruticoſus, erectus, paniculatus, ſubpubeſcens.* Folia *lanceolata ſ ovato-oblonga, ſerraturis aequalibus absque dente elongato ad baſin* Pedunculi *uniflori, lateriflori, absque bracteis* Germina *hirta.* Siliquae *bivalves, compreſſae.* Flores Vernales *apetali, tetrandri, calyce 4-phyllo.* Autumnales 5-*petali, polyandri, calyce* 5-*phyllo.*

1053. SEGUIERIA.

Gen. pl. ed Schreb. n. 918.

Cal. 5-phyllus. *Cor.* 0. *Capſ.* 1-ſperma: ala magna terminata, alulisque lateralibus.

1. SEGUIERIA *americana.*
S. americana. *Sp. pl.* 747.
Seguieria aculeata. *Iacq. amer.* 170. *Loefl. it.* 191.
Houttuyn Lin. Pfl. Syſt. 4. *p.* 35.
Americaniſche Seguierie. W.
Habitat in America *meridionali, e. g.* Carthagenae *ad vias.* ♄

Folia *alterna, petiolata, elliptica, emarginata.* Aculei *juxta folium, recurvi.* Flores *racemoſi, albi, foetidi.*

*1054. SWARTZIA.

Rittera. Gen. pl. ed. Schreb. n. 919.

Cal. 4-phyllus. *Petalum* 1 laterale planum. *Legumen* uniloculare bivalve. *Semina* arillata.

*1. SWARTZIA *ſimplicifolia.*
S. foliis ſimplicibus, petalo ſubrotundo ovato calyce majore, polyandra.
Rittera ſimplex *Vahl. ſymb.* 2. *p* 60. *
Poſſira (*ſimplex*) foliis ſimplicibus oblongis. *Swartz. prod.* 82.
Einfachblättrige Swartzie. W.
Habitat in inſulis Caribaeis. ♄

*2. SWARTZIA *grandiflora.* †

S. foliis ſimplicibus oblongo-ovatis, pedunculis ſubtrifloris, petalo ſubrotundo reniformi maximo, leguminibus oblongis.

Rittera grandiflora. *Vahl eclog.* 2. *p.* 37. * *Vahl plant. amer. dec.* 1. *t.* 9.

Groſsblütige Swartzie. W.

Habitat in inſula Trinitatis. ♄

Tam multa ſimilia habet haec ſpecies cum S. ſimplicifolia, ut limites infallibiles tradere fere nequeam, ambae autem inter ſe collatae differre videntur. Folia quidem in S. grandiflora anguſtiora magis acuminata et conſiſtentia tenuiora, petioli apex dente obſcuro, nullo modo autem per petiolum decurrente, pedunculi plerumque triflori tantum, petala majora. Sed notae hae omnes nimis levidenſes ſunt pro ſpecie indubitabili ſtabilienda. Characteres certiores forte petendi a petalo et legumine, cum vero in ſpecimine meo S. ſimplicifoliae parum bono, prior et poſterior plane deſideratur, neceſſitas me cogit aliis relinquere ulterius inquirere, num ſpecies diverſae revera ſint, nec ne. Vahl l. c.

*3. SWARTZIA *dodecandra.*

S. foliis ſimplicibus, floribus dodecandris, petalo oblongo longitudine calycis.

Rittera dodecandra. *Vahl ſymb.* 2. *p.* 60. *t.* 34. *

Zwölffädige Swartzie. W.

Habitat in America *meridionali.* ♄

*4. SWARTZIA *triphylla.*

S. foliis ternatis, petiolis marginatis.

Poſſira triphylla. *Swartz prod.* 82.

Poſſira arboreſcens. *Aubl. guj.* 2. *p.* 934. *t.* 355. *

Rittera. *Schreb. gen. pl. n.* 919.

Dreyblättrige Swartzie. W.

Habitat in Gujana, Cajenna. ♄

*5. SWARTZIA *pinnata.*

S. foliis pinnatis, petiolo communi tereti. W.

Rittera (*pinnata*) foliis pinnatis. *Vahl eclog.* 2. *p.* 38. *

Gefiederte Swartzie. W.

Habitat in inſula Trinitatis. ♄

*6. SWARTZIA *alata.* W.

S. foliis pinnatis, petiolo communi alato. W.

Swartzia. *Schreb. gen. pl. n.* 1227.

Tounatea gujanenſis. *Aubl. guj.* 1. *p.* 550. *t.* 218. *
Geflügelte Swartzie. W.
Habitat in Gujanae *ſylvis.* ♄

Rittera et Swartzia Schreberi eſt unum genus, hinc ſuaſu Clariſſ. Vahl nomen Swartziae ſervavi, in memoriam amici aeſtumatiſſimi et Botanici holmienſis celeberrimi. W.

DIGYNIA.

1055. PAEONIA.

Gen. plant. ed. Schreb. n. 920.

Cal. 5-phyllus. *Petala* 5. *Styli* 0. *Capſ.* polyſpermae.

1. PAEONIA *officinalis.*

P. foliis decompoſitis nudis, foliolis lobatis, lobis lato-lanceolatis, capſulis rectiuſculis tomentoſis. *Retz. obſ.* 3. *p.* 35. *

P. foliolis oblongis. *Hort. cliff.* 212. *Hort. upſ.* 149. *Mat. med.* 137. *Sauv. monſp.* 307. *Scop. carn. n.* 650. *Regn. bot.*

P. radice glandulosa, foliis duplicato-pinnatis, pinnis ellipticis et trilobis. *Hall. helv. n.* 1187.

P. femina. *Mill. dict. n.* 2. *Kniph. cent.* 6. *n.* 69. *Knorr. del.* 1. *t. R.* 3. *et. t. R.* 2. *flore pleno.*

P. communis ſ. femina. *Bauh. pin.* 323.

P. femina. *Fuchſ. hiſt.* 202. *Blackw. t.* 65. *Matth.* 915. *Lob. ic.* 682. *Moriſ. hiſt.* 3. *ſ.* 12. *p.* 455. *t.* 1. *f.* 7. 13. 15.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 245.

Gemeine Paeonie. W.

Habitat in nemoribus montium Idae, Helvetiae. ♄ (v. v.)

Flores explicati regulares, ſed aeſtivatione omnino irregulares ſunt.

*2. PAEONIA *corallina.*

P. foliis biternatis, foliolis ovatis integris nudis, capſulis recurvatis tomentoſis. *Retz. obſ.* 3. *p.* 34. *

P. (*integra*) foliis biternatis, foliolis integris ovatis, corolla regulari integerrima. *Murr. comment. goett.* 1784. *et* 1785. *p.* 92.

Paeonia officinalis β. *Sp. pl.* 747.

P. folio nigricante ſplendido, quae mas. *Bauh. pin.* 323. *Moriſ. hiſt.* 3. *ſ.* 12. *p.* 454. *t.* 1. *f.* 1.
P. mas. *Lob. ic.* 684. *Blackw. t.* 245.
P. mas flore purpureo. *Besl. eyſt. vern. ord. VI. f.* 10.
Ganzblättrige Paeonie. W.
Habitat in Europa *auſtrali*, Sibiria. ♃ (v. v.)

*3. PAEONIA *albiflora*.
P. foliis biternatis, foliolis ovato-lanceolatis integris nudis, capſulis recurvatis glabris. W.
P. (*albiflora*) foliis ternatis, foliolis lobatis nitentibus, germinibus ternis glabris. *Pall. roſſ.* 2. *p.* 90. *t.* 84. *
P. lacteo flore, foliis utrinque viridantibus et ſplendentibus. *Amman. ruth.* 77. *n.* 103. *Gmel. ſib.* 4. *p.* 184.
Weiſse Paeonie. W.
Habitat in Sibiria *transbaicalenſi*. ♃ (v. ſ.)
Folium *biternatum, foliolis tribus utrinque lateralibus baſi ſaepius cohaerentibus.* W.

*4. PAEONIA *humilis*.
P. foliis biternatis, foliolis tripartito-laciniatis ſubtus ſubvilloſis, capſulis erectiuſculis piloſis. *Retz. obſ.* 2. *p.* 35. *
P. tenuis laciniata ſubtus pubeſcens flore purpureo. *Bauh. pin.* 323. *Moriſ. hiſt.* 3. *ſ.* 12. *p.* 455. *t.* 1. *f.* 8.
P. hiſpanica ſemine nata ſive II. *Cluſ. hiſt.* 1. *p.* 279.
P. pumila foemina. *Lob. ic.* 683.
Niedrige Paeonie. W.
Habitat in Hiſpania. ♃ (v. v.)
Dignoſcitur, licet P. officinali ſimilis ſit, caule *humiliore,* foliis *ſubtus pubeſcentibus,* capſulis *pubeſcentibus nec villoſis.* W.

5. PAEONIA *anomala*.
P. foliis biternatis, foliolis multipartitis nudis, laciniis lanceolatis, capſulis depreſſis glabris. W.
P. calyce folioſo, capſulis glabris depreſſis. *Mant.* 247.
P. fructibus quinque glabris patentibus. *Gmel. ſib.* 4. *p.* 184. *t.* 72.
P. (*laciniata*) foliis biternatis, foliolis acute laciniatis, germinibus glabris. *Pall. roſſ.* 2. *p.* 93. *t.* 85. *
Houttuyn Lin. Pfl. Syſt. 7. *p.* 247.
Unregelmäſsige Paeonie. W.
Habitat in omni Sibiria. ♃ (v. v.)
Differt a praecedenti Foliis *anomalis ſ. inaequaliter diviſis;* Calycis *foliolis apice lanceolatis longitudine*

corollae vel ultra; Germinibus *glabris, plerumque quinis.*

*6. PAEONIA *hybrida.*

P. foliis biternatis, foliolis multipartitis nudis, laciniis linearibus, germinibus pubeſcentibus. W.

P. (*hybrida*) foliis ternato-multifidis, laciniis linearibus, germinibus pubeſcentibus. *Pall. roſſ.* 2. *p.* 94. *t.* 86. *

Baſtard Paeonie. W.

Habitat - - - - ♃

Planta *teſte Clariſſ. Pallas in horto Petropolitano e ſeminibus P. tenuifoliae eadem areola cum P. anomala creſcentis orta, et quae nunquam ſemina perfecta dedit.* W.

7. PAEONIA *tenuifolia.*

P. foliis biternatis, foliolis multipartitis nudis, laciniis lineari-ſubulatis, capſulis tomentoſis. W.

P. foliolis linearibus multipartitis. *Lin. dec.* 9. *t.* 5. *Gmel. ſib.* 4. *p.* 185. *t.* 73. *Pall. roſſ.* 2. *p.* 95 *t.* 87. *

P. laciniis foliorum linearibus. *Zinn. goett.* 127.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 248.

Feinblättrige Paeonie. W.

Habitat in Ucrania, Sibiria. ♃ (v. v.)

Folia *in caule plurima, nec tantum* 3 *ſ.* 4. *ut in P. officinali.* Germina 3, *purpuraſcentia, veſtita pilis purpureis, nec albis P. officinalis.*

Fructus 2 *ſ.* 3. *Gmel.*

Piſtilla 3 *in omnibus Paeoniis plerumque vidi.* W.

1056. CURATELLA.

Gen. pl. ed. Schreb. n. 921.

Cal. 5-phyllus. *Petala* 4. *Styli* 2. *Capſ.* 2-partita: loculis diſpermis.

1. CURATELLA *americana.*

C. americana. *Sp. pl.* 748. *Aubl. guj.* 1. *p.* 579. *t.* 232. *

Curatella. *Loefl. it.* 260. *

Houttuyn Lin. Pfl. Syſt. 4. *p.* 36.

Americaniſche Curatelle. W.

Habitat in America *meridionali.* ♄

Statura Coccolobae. Folia *alterna, ſubſeſſilia, oblonga.*

1057. FOTHERGILLA.

Gen. plant. ed. Schreb. n. 922.

Cal. truncatus, integerrimus. *Cor.* 0. *Germen* bifidum. *Capſ.* 2 - locularis. *Sem.* ſolitaria, oſſea.

1. FOTHERGILLA *alnifolia.*

F. alnifolia. *Lin ſuppl.* 267. *Willd. arb.* 113. *
F. Gardeni. *Syſt. veg.* 13. *p.* 418.
α. foliis obtuſis. *Ait. Kew.* 2. *p.* 241.
β. foliis acutis. *Ait. l. c.*
F. Gardeni. *Iacq. ic. rar.* 1. *t.* 100. *Collect.* 1. *p.* 79. *
Houttuyn Lin. Pfl. Syſt. 4. *p.* 37.
Elſenblättrige Fothergille. W.
Habitat in Carolina. ♄ (v. v.)

Arbuſcula *Foliis fructuque Hamameli ſimilis*, Flore *diverſiſſima. D. Garden.*

Arbor *facie et foliis Betulae Alni.* Folia *alterna petiolata cuneiformia integra apice ſerrata, ſerraturis majoribus paucioribus, ſupra viridia, ſubtus cana, juniora ſubtus albido-tomentoſa.* Flores *ſpicati terminales congeſti, quaſi in capitulum oblongum, albi.* Capſulae *magnae ovatae hirſutiſſimae acutae biloculares.* Valvulae *aperiuntur in cuſpides quatuor dum ſemina matura ſunt.* Flores *primo vere e gemmis terminalibus ante folia proveniunt.* Suppl.

*1058. TRICHOCARPUS.

Gen. plant. edit. Schreb. n. 923.

Cal. 4 - f. 5-partitus. *Cor.* 0. *Styl.* 2 bifidi. *Capſ.* ſetoſa 1 - locularis, 4 - valvis polyſperma.

*1. TRICHOCARPUS *laurifolia.* W.

Ablania gujanenſis. *Aubl. guj.* 1. *p.* 585. *t.* 234. *
Lorbeerblättrige Haarfrucht. W.
Habitat in ſylvis Gujanae. ♄

Arbor 50-*pedalis.* Folia *ſparſa coriacea oblonga acuta integerrima venoſa glabra petiolata.* Corymbi *pauciflori ſubdichotomi laterales.* W.

*1059. LACIS.

Gen. pl. ed. Schreb. n. 924.

Cal. 0. *Cor.* 0. *Styli* 2. *Capſ.* 1-locularis bivalvis polyſperma.

*1. LACIS *fluviatilis.*

Mourera fluviatilis. *Aubl. guj.* 1. *p.* 562. *t.* 233. *
Waſſer-Schlitzblatt. W.
Habitat in fluviis Gujanae. ♃

Radix *repens.* Caulis *herbaceus ſimplex aſper.* Folia *alterna ſeſſilia utrinque glabra atro-viridia ampla profunde ſinuata, laciniis pinnatifido-ſinuatis, margine criſpa, venoſa.* Aculeus *ſolitarius carnoſus rigidus in venarum folii axillis.* Racemus *terminalis ſimplex.* Pedunculus *communis ſemiteres.* W.

TRIGYNIA.

*1060. HOMALIUM.

Gen. pl. ed. Schreb. n. 925.

Cal. 6-7-partitus. *Cor.* 6-7-petala. *Stamina* 21 per tria aggregata. *Capſ.* unilocularis polyſperma.

*1. HOMALIUM *racemoſum.*

H. foliis ſerratis, racemis axillaribus terminalibusque, floribus pedunculatis. *Swartz. prod.* 86.
Homalium racemoſum. *Iacq. amer.* 170. *t.* 183. *f.* 72.
Traubiges Homalium. W.
Habitat in Martinica, Jamaica. ♄

*2. HOMALIUM *Racoubea.*

H. foliis dentatis coriaceis, racemis terminalibus, floribus ſubſeſſilibus. *Swartz. prod.* 86.
Racoubea gujanenſis. *Aubl. guj.* 1. *p.* 590. *t.* 236. *
Aehrenblütiges Homalium. W.
Habitat in ſylvis Gujanae. ♄

Genus *Blackwelliae affine, tamen* petalorum *numero cum laciniis calycinis aequali et* ſtaminibus *per tria aggregatis diſtinctum.* W.

1061. DELPHINIUM.

Gen. plant. ed. Schreb. n. 924.

Cal. 0. *Petala* 5. *Nectarium* bifidum, poſtice cornutum. *Siliquae* 3 ſ. 1.

* *Unicapſulares.*

1. DELPHINIUM *Conſolida.*

D. nectariis monophyllis, caule ſubdiviſo. *Hort. cliff.* 212. *Fl. ſuec.* 440; 476. *Mat. med.* 138. *Roy. lugdb.* 482. *Dalib. pariſ.* 158. *Crantz. auſtr. p.* 131. *Scop. carn. n.* 651. *Pollich. pal. n.* 512. *Fl. dan.* 683. *Ludw. ect. t.* 54. *Kniph. cent.* 2. *n.* 20. *Blackw. t.* 26. *Knorr. del.* 1. *t. R.* 17. *Hoffm. germ.* 186. *Roth. germ. I.* 230. *II.* 589.

D. uniſiliquum, caule brachiato diffuſo, floribus ſparſis. *Hall. helv. n.* 1203.

D. nectariis acuminatis ſubcurvatis pedunculo longioribus, foliis lineari-pinnatifidis. *Neck. gallob. p.* 233.

Conſolida regalis arvenſis. *Bauh. pin.* 142. *Tabernaem.* 62.

Conſolida regalis. *Cam. epit.* 521.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 249.

Feld-Ritterſporn. W.

Habitat in agris Europae *reſtibilibus.* ⊙ (v. v.)

Variat *flore pleno.* R.

2. DELPHINIUM *Ajacis.*

E. nectariis monophyllis, caule ſimplici. *Hort. cliff.* 213. *Hort. upſ.* 150. *Roy. lugdb.* 482. *Mill. dict. n.* 2. *Hoffm. germ.* 186.

D. uniſiliquum, caule recto ſpicato. *Hall. helv. n.* 120.

Conſolida regalis hortenſis, flore majore et ſimplici et minore. *Bauh. pin.* 142.

Conſolida regalis flore majore et multiplici. *Bauh. pin.* 142.

Flos regius. *Dod. pempt.* 252.

Delphinium ſativum. *Riv. t.* 123.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 250.

Garten-Ritterſporn. W.

Habitat in Helvetia *ad maceries.* ⊙ (v. v.)

Dic quibus in terris inſcripti nomine Regum
Naſcantur flores, et phyllida ſolus habeto.

Ecce ſuos gemitus foliis inſcripſit et Aia. Virg. eclog. Variat *flore pleno et ſemipleno. Riv.*

3. DELPHINIUM *aconiti.*

D. nectariis monophyllis antice quadridentatis, capſulis ſolitariis, ramulis unifloris. *Mant.* 77. *Vahl ſymb.* 1. *p.* 40. *t.* 13.

D. orientale annuum, flore ſingulari. *Tournef. cor.* 30.

Aconitum (*monogynum*) unicapſulare nectario unico, caule foliisque multifidis villoſis, flore pallide violaceo. *Forsk. cat. pl. conſt. p.* 27. *n.* 248.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 251.

Eiſenhutartiger Ritterſporn. W.

Habitat in Dardanellis. *Forskaehl.* ☉

Statura *D. ſegetum ſ. peregrini, corollis Aconiti.* Caulis *pedalis, paniculatus, ramoſus, caneſcens, pubeſcens.* Folia *pedato-multifida, linearia, caneſcentia: ſumma tantum tripartita.* Flores *terminales, ſolitarii, pedunculati, parvi, lividi, intus pupuraſcente virideque varii.* Nectarium exterius, *tubuloſum, reliquis longius, apice obtuſiſſimum, baſi elongatum, ut labium anterius horizontale in medio collocetur.* Nect. interius *baſi utrinque lobo obtuſo amplectitur ſtamina, ſed antice et ſuperius ſpatio ab exteriore prominet dentibus quatuor ſubaequalibus, margine linea purpuraſente inſcriptis: apex hujus intra exteriorem, obtuſiſſimus, craſſiuſculus, recurvatus collo coarctato.* Petala *inferiora* 4, *ovato-lanceolata, aequalia, vireſcentia lateribus purpuraſcentibus.* Stamina *longitudine petalorum purpuraſcentia.* Germen *unicum.* H. U.

** *Tricapſulares.*

4. DELPHINIUM *ambiguum.*

D. nectariis monophyllis, corollis hexapetalis, capſulis ternis, foliis multipartitis *Sp. pl.* 749. *Mill. dict. n.* 3.

D. elatius, ſimplici flore. *Cluſ. hiſt.* 2. *p.* 206.

Conſolida regalis, flore minore. *Bauh. pin.* 142.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 252.

Zweifelhafter Ritterſporn. W.

Habitat in Mauritania. ☉

Statura *D. Ajacis, ſed magis caneſcens.* Rami *divaricati in caule ſimplici.* Corollae *caeruleae, extus virides.* Nectarium *monophyllum, bifidum lobis acutiuſculis. Calcar ſaepe apice bidentatum.* Differt a *D.*

monogynis numero capſularum; a D. trigynis Nectario interiore monophyllo ſ. calcare interiore non bipartito; adeoque Staphiſagriae genere ſeparari nequeunt. Piſtilla *variant in eadem planta* 1 *et* 3, *ut vix conſtans numerus.*

5. DELPHINIUM *peregrinum.*

D. nectariis diphyllis, corollis enneapetalis, foliis multipartitis, obtuſis. *Sp. pl.* 749. *Hort. cliff.* 213. *Mill. dict. n.* 4. *Allion. pedem. n.* 1508. *t.* 25. *f.* 3. *Desfont. fl. atl.* 1. *p.* 426. *

D. nectariis diphyllis, floribus ſolitariis, foliis multipartitis: foliolis linearibus acuminatis. *Allion. nic.* 200. *taur.* 71. *

Conſolida regalis latifolia, parvo flore. *Bauh. pin.* 142. *prodr.* 74. *t.* 74. *Moriſ. hiſt.* 3. *p.* 466. *ſ.* 12. *t.* 4. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 252.

Fremder Ritterſporn. W.

Habitat in Italia, Sicilia, Melita, Palaeſtina. ☉ (v. ſ.)

Caulis *ſimplex: Ramis virgatis, nudis, ſtrictis.* Folia *D. Ajacis facie.* Bracteae *ſubulatae.* Corolla *caerulea, praeter Nectaria et quatuor petala inferiora, habet praeterea Petala* 2 *lateralia ſubrotunda, unguibus longis affixa, quae deflectuntur verſus petala exteriora ſ. infima, et iis incumbunt.* Nectaria *nulla habent labella deflexa.* Capſulae *ternae.*

6. DELPHINIUM *grandiflorum.*

D. nectariis diphyllis: labellis integris, floribus ſubſolitariis, foliis compoſitis lineari-multipartitis. *Sp. pl.* 749. *Mill. dict. n.* 6. *ic. t.* 250. *f.* 1. *Gmel. ſib.* 4. *t.* 78.

D. nectariis diphyllis labellis integris, floribus ſubſolitariis, foliis compoſitis lineari-multipartitis. *Hort. upſ.* 150. * *Sp. pl. ed.* 2. *p.* 749.

D. luſitanicum glabrum, aconiti folio. *Rolof. hort.* 61. *t.* 3.

D. elatius ſubincanum perenne, floribus amplis azureis. *Amm. ruth.* 175. *Mill. dict. t.* 119.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 253.

Groſsblütiger Ritterſporn. W.

Habitat in Sibiria. ♃ (v. v.)

Capſulae incanae ſunt.

*7. DELPHINIUM *intermedium.*

D. nectariis diphyllis, labellis ovatis bifidis, laciniis ova-

tis, foliis tripartitis, laciniis trifidis inciſis. *Ait. Kew.* 2. *p.* 243.

Delphinium elatum. *Mattuſch. enum. ſtirp. ſileſ. p.* 132.

D. nectariis diphyllis, labellis integris, floribus ſpicatis, foliis palmatis multifidis glabris. *Mill. ic. p.* 79. *t.* 119.

β. Aconitum lycoctonum flore Delphinii II. ſileſiacum. *Cluſ. hiſt.* 2. *p.* 94.

Aconitum coeruleum hirſutum, flore coſolidae regalis. *Bauh. pin.* 183.

Mittlerer Ritterſporn. W.

Habitat in Sileſiae *montibus.* ♃ (v. v.)

Folia *radicalia quinqueloba, lobis inciſo-dentatis, laciniis oblongis. Varietas* α. *eſt glabra,* β. *vero foliis ſubtus pubeſcentibus.* W.

8. DELPHINIUM *elatum.*

D. nectariis diphyllis, labellis ovatis emarginatis, laciniis breviſſimis inaequalibus, foliis ſubpeltatis tripartitis, laciniis multifidis. *Ait. Kew.* 2. *p.* 244.

D. nectariis diphyllis, labellis bifidis apice barbatis, foliis inciſis, caule recto. *Hort. upſ.* 151. * *Kniph. cent.* 6. *n.* 35.

D. nectariis diphyllis, foliis peltatis multipartitis acutis. *Hort. cliff.* 213.

D. triſiliquum, foliis ſemiquinquelobis, acute ſerratis, petiolis interioribus quaternis, barbatis. *Hall. helv. n.* 1201.

D. perenne, aconiti folio ampliori, floribus caeruleis. *Amm. ruth.* 174. *Hall. goett.* 93. *Gmel. ſib.* 4. *t.* 77.

Aconitum caeruleum glabrum flore conſolidae regalis. *Bauh. pin.* 183.

Aconitum lycoctonum flore Delphinii II. *Cluſ. hiſt.* 2. *p.* 95.

Lycoctonum flore Delphinii. *Dod. pempt.* 441.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 254.

Hoher Ritterſporn. W.

Habitat in Sibiria, Helvetia. ♃ (v. v.)

Folia *radicalia profunde quinquepartita, laciniis trifidis inciſis linearibus.* W.

*9. DELPHINIUM *hybridum.*

D. nectariis diphyllis, labellis ovatis bifidis, laciniis breviſſimis inaequalibus, foliis tripartitis, laciniis trifidis linearibus utrinque pubeſcentibus. W.

Delphinium hybridum. *Stephan in litteris.*

Baſtard Ritterſporn. W.

Habitat in Sibiria. (v. ſ.)

Folia *utrinque, ut tota planta, pubeſcentia quinque-partita, laciniis trifidis linearibus.* Folia *caulina tri-partita, laciniis anguſtisſimis, intermedia trifida, la-teralibus bifidis. Simillimum praecedenti, ſed totum pubeſcens, et flores duplo minores. An varietas?* W.

*10. DELPHINIUM *exáltatum.*

D. nectariis diphyllis, labellis oblongis bifidis, laciniis lanceolatis aequalibus, foliis tripartitis, laciniis trifidis. *Ait. Kew.* 2. *p.* 244.

D. nectariis diphyllis, labellis bifidis apice barbatis, foliis trilobis inciſis, caule erecto. *Mill. ic. p.* 167. *t.* 250. *f.* 2.

Americaniſcher Ritterſporn. W.

Habitat in America *boreali.* ♃ (v. ſ.)

Maxime affine D. urceolato, differt vero: foliis *gla-bris planis,* caule *glabro purpureo.* Ait. l. c.

Folia *tripartita, laciniis trifidis lanceolatis.* W.

*11. DELPHINIUM *urceolatum.*

D. nectariis diphyllis, labellis oblongis bifidis, laciniis lanceolatis aequalibus, foliis peltato-concavis triparti-tis pubeſcentibus, laciniis trifidis. W.

D. (*urceolatum*) nectariis diphyllis, labellis bifidis apice barbatis, foliis palmatis urceolatis. *Iacq. ic. rar.* 1. *t.* 101. *collect.* 1. *p.* 153.

Hohlblättriger Ritterſporn. W.

Habitat - - - - ♃ (v. ſ.)

Folia *baſi ſuperne concava et ſubtus pubeſcentia, cete-rum forma praecedentis.* W.

12. DELPHINIUM *puniceum.*

D. nectariis diphyllis, labellis oblongis bifidis, laciniis lanceolatis aequalibus, germinibus pedunculisque vil-loſis, foliis tripartitis, laciniis multifidis. W.

D. labellis bipartitis piloſis, nectarii cornu recto, foliis multipartitis, bracteis calycinis nullis. *Lin. ſuppl.* 267.

Delphinium puniceum. *Pall. itin.* 3. *p.* 96.

Rothbrauner Ritterſporn. W.

Habitat in Sibiria. ♃ (v. v.)

Differt ab omnibus reliquis ſpeciebus fuſco-rubente colore florum ut in Veratro nigro.

In omnibus D. eláto ſimillima, ſed differt: Caule, Petiolis *et* Pedunculis *pubeſcentibus;* Petalis *corollae aequalibus,* Corolla *magis regulari quam in D. elato,*

labellis profunde partitis rectis, Cornu *recto laevi longitudine pedicellorum*, Bractea *ſubulata utrinque ad baſin corollae nulla. Primum credidi varietates eſſe, poſtea vero comparatis partibus vidi differentias. Bracteas in omnibus floribus hujus ſpeciei a me examinatis vidi.* W.

13. DELPHINIUM *Staphiſagria.*

D. nectariis tetraphyllis petalo brevioribus, foliis palmatis: lobis obtuſis. *Mat. med.* 138. *Regn. bot.*

D. nectariis diphyllis petalo brevioribus, foliis palmatis: lobis obtuſis. *Hort. cliff.* 213. *Hort. upſ.* 150. *Roy. lugdb.* 482. *Sauv. monſp.* 214. *Sp. pl.* 2. *p.* 750. *Mill. dict. n.* 8. *Scopoli. carn. n.* 652.

Staphiſagria. *Bauh. pin.* 324. *Dod. pempt* 366. *Blackw. t.* 265.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 255.

Scharfer Ritterſporn. W.

Habitat in Iſtria, Dalmatia, Calabria, Apulia, Creta, Galloprovincio. ♂ (v. f.)

Foliorum *lobi trifidi.* Nectarii *cornu breviſſimum, obtuſum.*

Varietas *foliis albo-venoſis differt* Foliis *inciſis*, 7-*partitis*, *laciniis acuminatis*; Corollae *calcari non obſoleto*, *ſed longitudine pedicelli.*

*14. DELPHINIUM *pentagynum.*

D. nectariis diphyllis, labellis cuneiformibus inaequalibus bifidis, laciniis cuneatis, foliis multipartitis linearibus, piſtillis quinis. W.

D. piſtillis quinis, foliis multifariam decompoſitis, caule ſuperne pubeſcente, floribus heptapetalis, nectario gemello. *Desfont. fl. atl.* 1. *p.* 427. *t.* 111. *

D. calcaribus internis bifidis, capſulis quinis, foliis palmato-multifidis glabris. *Lamarck. encycl.* 2. *p.* 262. *

D. luſitanicum glabrum Aconiti folio. *Tournef. inſt.* 426.

Fünfgriffliger Ritterſporn. W.

Habitat inter ſegetes Algeriae *inque* Luſitania. ⊙

Caulis *pedalis vel tripedalis ramoſus erectus laeviter flexuoſus, ſuperne pubeſcens lanugine breviſſima denſa.* Folia *D. Ajacis leviſſime pubeſcentia aut glabra, radicalia circumſcriptione rotunda, longe petiolata, caulina flabelliformia multifariam decompoſita, laciniis linearibus inaequalibus acutis divergentibus.*

Petioli *baſi vaginantes.* Flores *racemoſi ſinguli pedicellati, pedicellis pubeſcentibus erectis, bracteola ſubulata duplo triplove longioribus.* Foliola *bina alterna conformia, infra ſinguli pedicelli apicem.* Calyx *nullus.* Corolla *magnitudine D. Ajacis caerulea.* Germina 5 *rarius* 3 *villoſa.* Styli *et* Capſulae *totidem polyſpermae.* Desfont. l. c.

1062. ACONITUM.

Gen. pl. ed. Schreb. n. 928.

Cal. o. *Petala* 5, ſupremo fornicato. *Nectaria* 2, pedunculata, recurva. *Siliquae* 3 ſ. 5. (†)

* *Corollis flavis vel albis.*

1. ACONITUM *Lycoctonum.*

A. cucnlli calcare ſpirali, labio ovato emarginato, corollae galea conica elongata, foliis palmatis pubeſcentibus, laciniis trifidis dentatis. W.

A. foliis palmatis multifidis villoſis. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 188. *t.* 81. *Scop. carn. n.* 653. *Pall. it.* 1. *p.* 72. *Iacq. auſtr. t.* 380. *Blackw. t.* 563. *Kniph. cent.* 5. *n.* 4. *Crantz. auſtr. p.* 131. *n.* 3. *Matruſch. ſil. n.* 384. *Gunn. norv. n.* 14. *Roth. germ. I.* 231. *II.* 590. *Koelle Aconit.* 23. * *Hoffm. germ.* 188.

A. foliis hirſutis, ſemitrilobis, lobis duplicato-trifidis. *Hall. helv. n.* 1200.

A. foliis peltatis multifidis, petalo ſupremo cylindraceo. *Fl. lapp.* 211. *Hort. cliff.* 213. *Fl. ſuec.* 441, 476. *Roy. lugdb.* 482.

A. lycoctonum luteum. *Bauh. pin.* 183.

A. lycoctonum, flore luteo. *Besl. eyſt. aeſt.* 25. *f.* 2.

Napellus flore luteo. *Riv. t.* 487.

Aconitum II. *Cam. epit.* 827.

A. lycoctonum vulgare flore luteo. *Cluſ. hiſt.* 2. *p* 94.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 257.

(†) *Nectarium Aconiti eſt cucullus pedunculatus, cujus baſis ſurſum ſpectans eſt calcar, et cujus apex eſt labium. Forma vero cuculli in diverſis ſpeciebus diverſe obſervatur et characteres non ſpernendos ſuppeditat.* W.

Wolfs-Eiſenhut. W.
Habitat in alpibus Galliae, Helvetiae, Sileſiae, Auſtriae, Italiae. ♃ (v. v.)

*2. ACONITUM *japonicum.* †
A. foliis trifido-palmatis, laciniis inciſis obtuſis. *Thunb. jap.* 231.
Iapaniſcher Eiſenhut. W.
Habitat in Japonia. ♃
Caulis *teres glaber herbaceus.* Folia *petiolata palmato-trifida, lobi laterales bifidi, medius trifidus, omnes obtuſi inciſo-dentati, dentibus rotundatis cum acumine.* Florum *racemus brevis. Differt ab A. Lycoctono:* foliis *tantum profunde trifidis,* laciniis *inciſis obtuſis rotundato-dentatis.* Thunb. l. c.

3. ACONITUM *pyrenaicum.*
A. cuculli calcare recto obtuſo, labio ſpathulato emarginato, galea conica, foliis digitato-quinquepartitis, laciniis tripartito-dentatis cuneiformibus. W.
A. foliis multipartitis: laciniis linearibus incumbentibus ſquarroſis. *Hort. upſ.* 152. * *Mill. dict. n.* 8. *Pall. it.* 2. *p.* 316. *Koelle Aconit.* 24. *
A. pyrenaicum luteum, foliorum ſegmentis ſibi invicem incumbentibus. *Raj. europ.* 307.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 260.
Pyrenäiſcher Eiſenhut. W.
Habitat in Sibiria, Tataria, Pyrenaeis. ♃ (v. ſ.)
Racemus ante antheſin nutat.

*4. ACONITUM *ochroleucum.* W.
A. cuculli calcare arcuato obtuſo, labio lanceolato emarginato, galea conica elongata, foliis ſubtus pubeſcentibus, palmato-quinquelobis, laciniis tripartitis inciſo-dentatis. W.
Aconitum flore albo. *Stephan. litt.*
Gelbweiſser Eiſenhut. W.
Habitat in Sibiria *ad* Caucaſum. ♃ (v. ſ.)
Caulis *teres pubeſcens.* Folia *radicalia petiolata ampla ſuperne glabra, ſubtus tenuiſſime pubeſcentia, palmato-quinqueloba, laciniis trifidis inciſo-dentatis, dentibus lanceolatis.* Folia *caulina ſubpeltata parva breve petiolata.* Racemus *terminalis ſimplex, alii racemi pedunculati ex ſupremis foliorum axillis ſolitarii.* Corollae *galea ſeſſilis conica elongata medio an-*

guſtata apice obtuſa, inferne dilatata acuminata. Petala *bina inferiora oblonga obtuſa ciliata.* Cuculli *calcar anguſtum arcuato - uncinatum apice obtuſum, labium lanceolatum rectum obtuſum emarginatum.* Piſtilla *tria.* Flores *ex toto pubeſcentes ochroleuci.* Habitu *A. ſeptentrionali ſimile.* W.

5. ACONITUM *Anthora.*

A. cuculli calcare refracto uncinato, labio adſcendente obcordato, galea conica rotundata, foliis multifidis, laciniis linearibus. W.

A. floribus pentagynis, foliorum laciniis linearibus. *Mat. med. p.* 139. *Mill. dict. n.* 4. *Gmel. ſib.* 4. *p.* 190. *Jacq. auſtr. t.* 382. *Scop. carn. n.* 655. *Ludw. ect. t.* 104. *Blackw. t.* 562. *Kniph. cent.* 4. *n.* 1. *bot. Koelle Aconit.* 24. *

A. foliorum laciniis linearibus diſtinctis ubique ejusdem latitudinis. *Hort. cliff.* 214. *Hort. upſ.* 152. *Roy. lugdb.* 483.

A. foliis hirſutis, ſemitrilobis, lobis duplicato - trifidis. *Hall. helv. n.* 1199.

A. ſalutiferum ſ. Anthora. *Bauh. pin.* 184.

Anthora ſ. Antithora. *Cam. epit.* 837. *Riv. t.* 486. *Mill. ic. t.* 12.

Anthora vulgaris. *Cluſ. hiſt.* 2. *p.* 98.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 266.

Feinblättriger Eiſenhut. W.

Habitat in alpibus Pyrenaeis, Helveticis, Taurinis, Allobrogicis. ♃ (v. v.)

*6. ACONITUM *album.*

A. cuculli calcare uncinato obtuſo, labio breviſſimo reflexo bifido, galea conica late unguiculata, foliis tripartitis, laciniis ovato - cuneiformibus trifidis dentatis. W.

A. (*album*) floribus pentagynis, foliis glabris tripartitis, laciniis acute inciſis, petali ſupremi ungue lateralibus longiore. *Ait. Kew.* 2. *p.* 246.

Aconitum orientale. *Mill. dict.*

A. lycoctonum orientale flore magno albo. *Tournef. cor.* 30.

Weiſser Eiſenhut. W.

Habitat in Oriente ♃ (v. v.)

Habitu *A. variegato haud diſſimile.* W.

* * *Corollis coeruleis.*

*7. ACONITUM *ſeptentrionale.*

A. cuculli calcare revoluto, labio ſpathulato ſubemarginato, galea conica elongata, foliis palmatis pubeſcentibus, laciniis trifidis divaricatis dentatis. W.

Aconitum ſeptentrionale. *Koelle Aconit.* 22.

Aconitum Lycoctonum β. flore coeruleo. *Lin. ſyſt. veg. ed. R.* 2. *p.* 615. *Stroem. ſöndm.* 1. *p.* 67. *Gunn. norv. n.* 14. β. *Gort. ingr.* 84. *Fl. dan. t.* 123. *Fabr. it. norv.* 119.

A. Lycoctonum floribus pallide purpureis. *Gmel. ſib.* 4. *p.* 189.

Calceolus lapponum ſ. Braſſica rangiferorum. *Scheff. lapp.* 360.

Nördlicher Eiſenhut. W.

Habitat in Norvegia, Suecia, Ruſſia, Sibiria. ♃ (v. ſ.)

Simillimum A. Lycoctono, ſed altius, foliorum *dentes longiores et acutiores*, corollarum *galeae longiores*, nectaria *magis obliqua.* Koelle. l. c.

8. ACONITUM *Napellus.*

A. cuculli calcare recto obtuſo, labio lanceolato adſcendente bifido, galea convexa, foliis nitidis quinquepartitis, laciniis tripartitis inciſis linearibus. W.

A. foliorum laciniis linearibus ſuperne latioribus linea exaratis. *Hort. cliff.* 214. *Fl. ſuec.* 442, 477. *Gron. virg.* 165. *Mat. med.* 138. *Gmel. ſib.* 4. *p.* 189. *itin.* 1. *p.* 8. *Hoffm. germ.* 187. *Koelle Aconit.* 14. *c. icone.*

A. caule ſimplici, ſpica denſa, petiolis unifloris, caſſidis mucrone brevi. *Hall. helv. n.* 1197.

A. coeruleum ſ. Napellus 1. *Bauh. pin.* 183.

A. lycoctonum VI. Napellus vulgaris. *Cluſ. hiſt.* 2. *p.* 96.

Napellus verus. *Lob. hiſt.* 387.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 258.

Wahrer Eiſenhut. W.

Habitat in alpibus Praegratenſibus, Helvetiae, *inque* Suecia, Sibiria. ♃ (v. ſ.)

Caulis *ſimpliciſſimus firmus, inferne denſe folioſus.* Spica *florum terminalis brevis denſe compacta, ſerius ad baſin ſubracemoſa, ramulis e ſummorum foliorum axillis brevibus paucifloris.* Pedunculi *uniflori.* Galea *humilius fornicata, antrorſum breviter roſtrata, tandem glabra, cum reliquis petalis ſaturatiſſime atroviolacea.* Folia *petiolata digitato - quinquepartita,*

ſegmentis anguſtis lineari-lanceolatis laciniatis, laciniis remote dentatis, dentibus linearibus, ſupra ex nigro ſaturate viridia, ſubtus pallide aquoſe virentia, utrinque nuda laevia nitida. Koelle l. c.

*9. ACONITUM *neomontanum.*

A. cuculli calcare adunco obtuſo, labio lanceolato revoluto bifido, galea convexa, foliis nitidis digitato-quinquepartitis, laciniis lato-lanceolatis breviſſime dentatis, pedunculis glabris. W.

Aconitum neomontanum. *Koelle Aconit.* 16.

A. (*neomontanum*) caule elongato, ſpica racemoſo-ramoſa, galea humili breviter roſtrata, foliis palmato-quinquepartitis, ſegmentis profunde inciſis latioribus. *Hoffm. germ.* 187.

Aconitum Napellus. *Roth. germ. I.* 231. *II.* 591. *Iacq. auſt.* 4. *p.* 42. *t.* 381.

Aconitum. *Act. holm.* 1739. *p.* 41. *t.* 2.

A. purpureum ſeu Napellus 3. *Bauh. pin.* 183. *Rivin. t.* 488.

A. lycoctonum V. neubergenſe. *Cluſ. hiſt.* 2. *p.* 96.

Neubergiſcher Eiſenhut. W.

Habitat alpeſtribus ſylvaticis, Carinthiae, Carnioliae, *inque montoſis* Germaniae. ♃ (v. v.)

Caulis *elongatus firmus, ſpica florum longa laxa inferne racemoſo-ramoſa, pedunculis unifloris.* Flores *ſaturate violacei, galea humili fornicata antrorſum breviter roſtrata et ipſa tandem glabra.* Folia *in inferiore caule laxius ſparſa petiolata ampla ſubpalmato-quinquepartita, ſegmentis lateſcentibus, profunde inciſo-laciniatis, laciniis lato-lanceolatis dentatis, dentibus brevibus latiuſculis acutis, ſupra ſaturate viridia, ſubtus pallida aquoſe virentia, utrinque nuda laevia nitentia.* Koelle l. c.

*10. ACONITUM *tauricum.*

A. cuculli calcare adunco obtuſo, labio lanceolato revoluto bifido, galea convexa, foliis nitidis digitato-quinquepartitis, laciniis lanceolatis trifido-inciſis, pedunculis pubeſcentibus. W.

A. (*tauricum*) caule ſubſimplici, racemo florum brevi compacto, galea humili breviter roſtrata, antrorſum repando-ſubfornicata, foliis nitentibus quinquepartitis, ſegmentis oblongis latiuſculis laciniatis, laciniis remote inciſo-dentatis. *Wulffen in Iacq. Collect.* 2

p. 112. * *Iacq. ic. rar.* 3. *t.* 492. *Koelle Aconit.* 15. * *Hoffm. germ.* 187.

A. violaceum ſ. Napellus 2. *Bauh. pin.* 183.

A. Napello ſimile, ſed minus, coeruleum, praecocius. *Bauh. hiſt.* 3. *p.* 656.

A. lycoctonum IV. tauricum. *Cluſ. hiſt.* 2. *p.* 95.

Behaarter Eiſenhut. W.

Habitat in alpeſtribus apricis Taureri Malnizenſis *et* Radſtadienſis, *inque* Hungaria. ♃ (v. v.)

Simillimum praecedenti, differt vero: galea *obtuſiore,* laciniis *foliorum latioribus profundioribus,* pedunculisque *pubeſcentibus. In hortis ramoſius.* W.

*11. ACONITUM *volubile.*

A. cuculli calcare adunco obtuſo, labio obcordato adſcendente, foliis opacis quinquepartitis, laciniis cuneiformibus trifidis dentatis, caule volubili. W.

Aconitum volubile. *Koelle Aconit.* 21. * *Pall. hort. demidof.*

Windender Eiſenhut. W.

Habitat in Sibiria. ♃ (v. v.)

Ab omnibus cognitis et a binis praecedentibus affinibus ſatis diſtinctum caule volubili. W.

*12. ACONITUM *cernuum.*

A. cuculli calcare ſubadunco obtuſo, labio lanceolato emarginato, galea convexa, racemo nutante, foliis opacis digitato-quinquepartitis, laciniis trifidis inciſodentatis. W.

Aconitum cernuum. *Koelle Aconit.* 17.

A. lycoctonum coma nutante. *Cluſ. hiſt.* 2. *p.* 97.

Hängender Eiſenhut. W.

Habitat in alpinis rupeſtribus abruptis Taureri Naſſfeldenſis *et* Radſtadienſis. ♃ (v. ſ.)

Folia *opaca nunquam nitida.* Racemus *non ſemper compoſitus, ante perfectam vero floreſcentiam nutans, deinde vero cernuus.* W.

13. ACONITUM *variegatum.*

A. cuculli calcare adunco obtuſo, labio oblongo emarginato adſcendente, galea conica, foliis nitidis tripartitis, laciniis trifidis cuneiformibus dentatis. W.

A. floribus pentagynis, foliorum laciniis ſemipartitis ſuperne latioribus. *Syſt. veg.* 419. *Koelle Aconit.* 18. * *Hoffm. germ.* 188.

A. foliis multifidis: laciniis ſemipartitis ſuperne latis. *Hort. cliff.* 214. *Hort. upſ.* 151. *Roy. lugdb.* 482. *Sp. pl.* 2. *p.* 750.

A. caeruleum minus ſ. Napellus minor. *Bauh. pin.* 183.

A. caeruleum parvum. *Dod. pempt.* 441. *Dalech. hiſt.* 1743.

A. lycoctonum X. *Cluſ. hiſt.* 2. *p.* 97.

Lycoctonum coeruleum parvum facie Napelli. *Lob. hiſt.* 386.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 261.

Bunter Eiſenhut. W.

Habitat in Italiae, Bohemiae, Carnioliae *montibus.* ♃ (v. ſ.)

Racemus *ſubſimplex.* Pedunculi *inferiores longiſſimi.* W.

14. ACONITUM *Cammarum.*

A. cucullí calcare adunco obtuſo, labio lanceolato recto acute emarginato, galea conica elongata, foliis opacis quinquepartitis, laciniis oblongo-cuneiformibus trifidis dentatis. W.

A. floribus ſubpentagynis, foliorum laciniis cuneiformibus inciſis acutis. *Sp. pl.* 751. *Mill. dict. n.* 9. *Koelle Aconit.* 17. * *Iacq. auſt.* 5. *p.* 11. *t.* 424. *Hoffm. germ.* 188. *Roth. germ. II.* 592. *Blackw. t.* 561.

A. caule ramoſo, paniculato, petiolis multifloris, caſſidis mucrone longo. *Hall. helv. n.* 1198.

A. caeruleo-purpureum, flore maximo ſ. Napellus 4. *Bauh. pin.* 183.

A. lycoctonum IX. judenbergenſe. *Cluſ. hiſt.* 2. *p.* 97.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 262.

Langhelmiger Eiſenhut. W.

Habitat in Stiria, Taurero. ♃ (v. ſ.)

Corollae *galea mucrone elongato.*

Folia *ſuperiora ſubſeſſilia.* W.

15. ACONITUM *uncinatum.*

A. floribus ſubpentagynis, foliis multilobis, corollarum galea longius extenſa. *Sp. pl.* 750. *Koelle Aconit.* 22.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 264.

Hakenförmiger Eiſenhut. W.

Habitat in Philadelphia. ♃

Folia *triloba ſ. quinqueloba, angulato-dentata, nuda.* Flores *caerulei, ſingulares, apice galeae uncinato, recto et longius, quam ipſe unguis galeae, extenſo.*

Haec americana flore fere A. Cammari, ſed foliis propior A. Napello.

TETRAGYNIA.

1063. WINTERA.

Gen. plant. edit. Schreb. n. 929.

Cal. 3-lobus. *Petala* 6. ſ. 12. *Germina* clavata. *Styli* 0. *Baccae* 4 ſ. 8 obovatae.

1. WINTERA *aromatica.*

W. pedunculis axillaribus aggregatis ſubtrifloris, floribus tetragynis. W.

W. (*aromatica*) pedunculis aggregatis terminalibus. *Forſt. Commentat. goett.* 9. *p.* 34. *t.* 7. *

Drymis (*Winteri*) pedunculis aggregatis terminalibus, piſtillis quatuor. *Lin. ſuppl.* 269. *Forſt. gen. n.* 42. *Forſt. Act. upſ.* 3. *p.* 181. * *Medicinal. obſerv.* 5. *p.* 41. *t.* 1. *Miller faſc.* (*figura pulcherrima.*)

Periclymenum rectum foliis laurinis, cortice aromatico acri. *Act. angl.* 1693. *n.* 204. *p.* 922. *t.* 1.

Gewürzhafte Wintere. W.

Habitat in Freti Magellanici *vallibus ſoli expoſitis.* ♄

Arbor *ſempervirens. Ex hac cortex Winteranus Officinarum verus, diverſiſſimus a Canella alba officinali figura, qualitatibus et vi.*

2. WINTERA *granadenſis.*

W. pedunculis axillaribus trifloris elongatis, floribus octogynis. W.

Drymis (*granadenſis*) pedunculis axillaribus elongatis trifloris, piſtillis octo. *Lin. ſuppl.* 269.

Dreyblumige Wintere. W.

Habitat in Nova Granada. *Mutis.* ♄

Hanc ſolam varietatem praecedentis a climate calidiori cauſſatam eſſe puto. Rami *longiores.* Folia *oblonga ſubtus magis glauca et fere albida longiora.* Pedunculi *axillares ſolitarii bini ſ. terni filiformes elongati, foliis tamen breviores, ſupra medium trifidi.* Pedicelli *uniflori.* Calyx *praecedentis.* Petala *ut in praecedente ſed* 12 *magis oblonga, unde flores majores, interiora exterioribus minora.* Stamina *praecedentis.*

Germina 8, *a quo numero flores rarius variant.* Baccae *totidem ac piſtilla.* Semina *ſubbina nitida. Corticis ſapor idem cum praecedente.*

3. WINTERA *axillaris.*

W. pedunculis axillaribus congeſtis unifloris, floribus tetragynis. *Forſt. prod. n.* 229.

Drymis (*axillaris*) pedunculis ternis axillaribus, piſtillo unico. *Lin. ſuppl.* 270. *Forſt. gen. n.* 42. *Forſt. Act. upſ.* 3. *p.* 182. *

Südſee-Wintere. W.

Habitat in Novae Zeelandiae *nemoribus.* ♄

Sapor totius plantae et inprimis corticis acerrimus eſt et pungens. Floret menſe Novembri. Proxima affinitas Micheliis.

Cur Linnaeus junior hanc monogynam fecit, non video. Forſt. l. c.

1064. TETRACERA.

Gen. plant. ed. Schreb. n. 936.

Cal. 5-ſ. 6-phyllus. *Cor.* 4-ſ. 5-petala. *Filamenta* ſuperne dilatata utrinque antherifera. *Capſulae* 4 latere dehiſcentes. *Semen* baſi arillatum.

* *Floribus monogynis.*

1. TETRACERA *ſarmentoſa.*

T. foliis oblongis ſerratis ſcabris, floribus monogynis. *Vahl ſymb.* 3 *p.* 70.

Delima ſarmentoſa. *Sp. pl.* 736. *Fl. zeyl.* 205. * *Amoen. acad.* 1. *p.* 403. *Burm. ind. p.* 122. *t.* 37. *f.* 1.

Frutex indicus ſarmentoſus, foliis hiſpidis rigidis. *Burm. zeyl.* 101.

Piripu. *Rheed. mal.* 7. *p.* 101. *t.* 54.

Houttuyn Lin. Pfl. Syſt. 4. *p.* 17.

Kletternder Raſpelſtrauch. W.

Habitat in Zeylona. ♄

Calycis *foliola ſaepe* 6. Capſula *ovata attenuata glabra, longitudinaliter altero latere dehiſcens.* Semen 1 *parvum nigrum, arillo denticulato inferne cinctum.* Vahl l. c.

*2. TETRACERA *tomentoſa.* W.
T. foliis ovatis acuminatis dentatis ſuperne glabris ſubtus tomentoſis, floribus monogynis. W.
Tigarea (*dentata*) foliis ovato-oblongis acuminatis, ſubtus tomentoſis. *Aubl. guj.* 2. *p.* 920. *t.* 351. *
Filziger Raſpelſtrauch. W.
Habitat in ſylvis Gujanae, Cajennae. ♄

*3. TETRACERA *aſpera.* W.
T. foliis ſubrotundis ſubrepandis ſcabris, floribus monogynis. W.
Tigarea (*aſpera*) foliis ſubrotundis undulatis, floribus racemoſis. *Aubl. guj.* 2. *p.* 918. *t.* 350. *
Rhininm. *Schreb gen. n.* 1545.
Scharfer Raſpelſtrauch. W.
Habitat in ſylvis Gujanae *et* Cajennae. ♄

*4. TETRACERA *Doliocarpus.* W. †
T. foliis oblongis acuminatis apice dentatis, pedunculis axillaribus unifloris, floribus monogynis. W.
Doliocarpus caule ſcandente, foliis ovatis dentatis, pedunculis lateralibus unifloris. *Roland. Act. holm.* 1756. *p.* 249. *t.* 9. *
Giftiger Raſpelſtrauch. W.
Habitat in Surinamo. ♄

*5. TETRACERA *ſtricta.* W. †
T. foliis ovato-lanceolatis dentatis, floribus terminalibus monogynis, caule ſtricto. W.
Doliocarpus caule ſtricto, foliis deflexis ovato-lanceolatis dentatis, floribus terminalibus. *Roland. Act. holm.* 1756. *p.* 249. *
Aufrechter Raſpelſtrauch. W.
Habitat in Surinamo. ♄

*6. TETRACERA *Calinea.* W.
T. foliis oblongis acuminatis integerrimis glabris, pedunculis multifloris lateralibus, floribus monogynis. W.
Calinea ſcandens. *Aubl. guj.* 1. *p.* 556. *t.* 221. *
Seitenblütiger Raſpelſtrauch. W.
Habitat in ſylvis Gujanae. ♄

*7. TETRACERA *obovata.* W.
T. foliis obovatis integerrimis glabris, pedunculis ſubcorymboſis lateralibus, floribus monogynis. W.
Soramia gujanenſis. *Aubl. guj.* 1. *p.* 552. *t.* 219. *
Mappia. *Schreb. gen. n.* 1755.

Doldentraubiger Raſpelſtrauch. W.
Habitat in Gujana *ad ripas fluvii* Sinemari. ♄

*8. TETRACERA *nitida.*
T. foliis lanceolato-oblongis ſcabris integerrimis, floribus monogynis. *Vahl ſymb.* 3. *p.* 70. *
Glänzender Raſpelſtrauch. W.
Habitat in inſula Trinitatis? ♄

* * *Floribus ſubtetragynis.*

*9. TETRACERA *Euryandra.*
T. foliis oblongis obtuſis laevibus integerrimis, floribus trigynis. *Vahl ſymb.* 3. *p.* 71.
Euryandra ſcandens. *Forſt. prod. n.* 228. *Forſt. gen. n.* 41.
Dreygriffliger Raſpelſtrauch. W.
Habitat in Nova Caledonia. ♄
Folia *petiolata ſubtus pallidiora bipollicaria et ultra.* Vahl l. c.

10. TETRACERA *volubilis.*
T. foliis ſcaberrimis ſerratis, floribus tetragynis. *Vahl ſymb.* 3. *p.* 71. *
Tetracera. *Hort. cliff.* 214. * *Mill. dict.*
Petraea floribus ſpicatis, ſcabris lauri foliis. *Amm. hort.* 581.
Fagus americana, ulmi ampliſſimis foliis, capſulis bigemellis. *Pluk. amalt.* 87.
Arbor americana convolvulacea platyphyllos barbadenſibus dicta, foliis ſerratis. *Pluk. alm.* 48. *t.* 146. *f.* 1.?
Houttuyn Lin. Pfl. Syſt. 2. *p.* 58.
Windender Raſpelſtrauch. W.
Habitat in America *meridionali.* ♄.

*11. TETRACERA *laevis.*
T. foliis oblongis laevibus ſubintegerrimis acuminatis, floribus terminalibus. *Vahl ſymb.* 3. *p.* 71. *
Glatter Raſpelſtrauch. W.
Habitat in India *orientali.* ♄
Folia *petiolata alterna baſi attenuata apice acuminata, ſuperne dentibus nonnullis obſolete ſerrata, bi-ſ. tripollicaria venoſa utrinque glabra.* Petiolus *breviſſimus.* Pedunculi *terminales ſolitarii vel gemini pollicares.* Flores *ſubracemoſi, pedicellis uni-ſ. bifloris.* Calyx *foliolis* 6 *ſubrotundis.* Stylos *tantum* 2 *obſervavi*

ſubulatos. Capſulae *quatuor unguiculares ſubrotundae ventricoſae mucronatae glaberrimae nitidae.* Semen *unicum parvum nigrum nitidum, arillo albido margine denticulato ad medium fere tectum.* Vahl l. c.

*12. TETRACERA *alnifolia.* W.
T. foliis oblongis acutis ſubintegerrimis ſubtus ſcabriuſculis: panicula terminali. W.
Elſenblättriger Raſpelſtrauch. W.
Habitat in Guinea. ♄ (v. ſ.)

Rami *teretes glabri.* Folia *alterna petiolata coriacea oblonga baſi attenuata apice rotundata acuta, margine plerumque integerrima interdum ad apicem dente uno alterove obſoleto inſtructa, venoſa, venulis parallelo-reticulatis, ſuperne nitida glabra inferne ſcabriuſcula.* Calyx *tetraphyllus.* Corolla *videtur pentapetala.* Filamenta *apice parum dilatata.* Anthera *in utroque latere.* Capſulae 4 *monoſpermae.* Semen 1 *nigrum arillo albido ex toto tectum.* Panicula *ſimplex terminalis.* Pedunculi *triflori.* W.

Aſſa indica Houttuyn Lin. Pfl. Syſt. 4. *p.* 40. *t.* 26. *f.* 1. *videtur hujus generis ſpecies.* W.

1065. CARYOCAR.

Gen. pl. ed. Schreb. n. 931.

Cal. 5-partitus. *Pet.* 5. *Styli* ſaepius 4. *Drupa* nucleis 4 ſulcis reticulatis.

1. CARYOCAR *nuciferum.*
C. foliis ternatis. W.
C. nuciferum. *Mant.* 247.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 60.
Nuſstragender Caryocar. W.
Habitat in Berbices *et* Eſſequebo. ♄

Arbor *procera.* Folia *ternata.*

Calyx Corolla*que purpurea.* Drupa *magnitudine capitis.* Nuclei *eſculenti amygdali ſaporis. Allamand.*

*2. CARYOCAR *butyroſum.* W. †
C. foliis quinatis utrinque glabris. W.
Pekea (*butyroſa*) fructu laevi, foliis digitatis utrinque glabris. *Aubl. guj.* 1. *p.* 594. *t.* 238. *
Rhizobolus. *Schreb. gen. n.* 932.
Caſtanea peruviana. *Cluſ. exot.* 129.

Glatter Caryocar. W.
Habitat in ſylvis Gujanae. ♄

*3. CARYOCAR *tomentoſum.* W. †
C. foliis quinatis ſubtus tomentoſis. W.
Pekea (*tuberculoſa*) foliis digitatis ſubtus tomentoſis. *Aubl. guj.* 1. 597. *t.* 239.
Filziger Caryocar. W.
Habitat in Gujanae *ſylvis.* ♄

Totum Pekeae genus huc pertinere videtur, forte etiam Saouari Aubletii. W.

*1066. WAHLBOMIA.

Thunb. Act. holm. 1790.

Cal. 4-phyllus. *Corolla* 4-petala. *Fructus* oblongus. *Styli* perſiſtentes.

*1. WAHLBOMIA *indica.*
W. indica. *Thunb. Act. holm.* 1790. *p.* 203. *t.* 9. *
Indiſche Wahlbomie. W.
Habitat in Java. ♄

Rami *alterni teretiuſculi cinereo-villoſi erecto patuli.* Folia *alterna petiolata elliptico-ovata acuminata, inferne integra, ſuperne ſerrata, ſupra viridia, ſubtus pallidiora, parum villoſa patentia digitalia.* Petioli *breves vix unguiculares.* Flores *in ramulis terminales ſubumbellati.* Umbella *ſimplex et compoſita ſubquadriflora.* Petioli *et* Pedicelli *filiformes villoſi patuli.* Thunb. l. c.

Tetraceris affinis et vix ſeparanda, auctoritate tantum Illuſt. Thunbergii diſtinxi, qui fructum mihi ignotum forte examinavit. W.

1067. CIMICIFUGA.

Gen. pl. ed. Schreb. n. 933.

Cal. 4-phyllus. *Cor.* nectariis 4, urceolatis. *Capſ.* 4. *Sem.* ſquamoſa.

1. CIMICIFUGA *foetida.*
C. foetida. *Syſt. nat. ed.* 12. *p.* 659.

Actaea (*Cimicifuga*) racemis paniculatis, fructibus quadricapſularibus. *Sp. pl.* 2. *p.* 722. *Amoen. acad.* 2. *p.* 54. *Amoen. acad.* 7. *t.* 6. *f.* 2. *Gmel. ſib.* 4. *p.* 181. *t.* 70.

Thalictroides foetidiſſimum, criſtophorianae facie. *Amm. ruth.* 102.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 265.

Stinkendes Wanzenkraut. W.

Habitat in Sibiria. ♃

Simillima *Actaeae racemoſae.* Variat *omnium maxime numero* Calycis, Corollae, Piſtillorum, *ut neque ſexus conſtans ſit, neque proportio.*

P E N T A G Y N I A.

1068. A Q U I L E G I A.

Gen. pl. ed. Schreb. n. 934.

Cal. o. *Petala* 5. *Nectaria* 5, corniculata, inter petala. *Capſ.* 5, diſtinctae.

1. AQUILEGIA *viſcoſa.*

A. nectariis incurvis: caule ſubnudo ſubunifloro viſcoſo-piloſo, foliis ſubtrilobis. *Mant.* 77. *Gouan. monſp.* 267. *

A. caule ſubnudo ſubunifloro, foliis ternatis, foliolis petiolatis ſubtrilobis. *Gouan. illuſtr.* 33. *t.* 19. R.

A. montana flore parvo Thalictri folio. *Bauh. prodr.* 75. *pin.* 144. *huc pertinet monente De la Cheval in Act. Helv. VIII. p.* 146. *t.* 5. R.

A. viſcoſa. *Magn. monſp.* 24. *hort.* 21.

A. hirſuta, flore viſcoſo. *Tournef. inſt.* 428.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 267.

Klebrige Akeley. W.

Habitat Monſpelii, *nec alibi etiamnum obſervata.* ♃ (v. ſ.)

Planta *tota pilis glandulofis, ſeſquipalmaris.* Folia *radicalia tripartita, laciniis trilobis;* caulina 1 ſ 2 *ternata: foliolis lanceolatis.* (Nectaria *incurva. Gouan.* R.)

2. AQUILEGIA *vulgaris.*

A. nectariis incurvis. *Hort. upſ.* 152. *Mat. med.* 159.

Fl. ſuec. 2. *n.* 478. *Crantz. auſtr. p.* 130. *Scop. carn. n.* 656. *Pollich. pal. n.* 513. *Fl. dan. t.* 695. *Ludw. ect. t.* 181. *Blackw. t.* 409. *Kniph. cent.* 5. *n.* 15. *Knorr. del.* 1. *tab. A. Hoffm. germ.* 188. *Roth. germ. I.* 231. *II.* 592.

Aquilegia. *Hort. cliff.* 215. *Roy. lugdb.* 482. *Dalib. pariſ.* 160.

A. foliis trilobatis, lobis petiolatis tripartitis, ſubrotundis, obtuſe dentatis. *Hall. helv. n.* 1195.

A. perigoniis et piſtillis pubeſcentibus, nectariis incurvis. *Ehrh. Beitr.* 7. *p.* 145.

A. (*ſylveſtris*) foliis compoſitis alternis ternatis: nectario incurvo. *Neck. gallob. p.* 234.

A. ſylveſtris. *Bauh. pin.* 144.

β. hortenſis ſimplex. *Bauh pin* 144.

γ. A. hortenſis multiplex, flore magno. *C. B.* 144.

δ. A. hortenſis multiplex, flore inverſo. *C. B.* 144.

ε. A. flore roſeo multiplici. *C. B.* 144.

ζ A. degener vireſcens. *Bauh. pin.* 144.

η. A. *vulgaris ſpecioſa*, nectariis coeruleis apice luteis. *Ait. Kew.* 2. *p.* 247.

A. (*bicolor*) perigoniis et piſtillis glabris, nectariis incurvis. *Ehrh. Beitr.* 7. *p.* 146.

ϑ. A. *vulgaris daurica*, foliis ſubtus pubeſcentibus, flore minori. W.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 268.

Gemeine Akeley. W.

Habitat in Europae *nemoribus ſaxoſis*, η. *et* ϑ. *in* Sibiria. ♃ (v. v.)

Sibirica *varietas* η. *conſtans*, *humilior*, *coerulea nectariorum oris albis*. Viſcoſa *monſpelienſis e ſeminibus Gouani enatis nil ab A. vulgari differebat. Syſt. Veg. p.* 420.

3. AQUILEGIA *alpina*.

A. nectariis rectis petalo lanceolato brevioribus. *Mill. dict. n.* 2. *Gmel. ſib.* 4. *p.* 185.

A. foliis trilobatis, lobis tripartitis linearibus, caule paucifloro. *Hall. helv. n.* 1196.

A. (*alpina*) perigoniis et piſtillis pubeſcentibus, nectariis fere rectis. *Ehrh. Beitr.* 7. *p.* 146.

A. montana, magno flore. *Bauh. pin.* 144. *prodr.* 75. *Burſ. VII.* 108. *Bauh. hiſt.* 3. *p.* 484. *Raj. hiſt.* 707.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 269.

Alpen-Akeley. W.
Habitat in Helvetia. ♂ (v. ſ.)

Folia *duplicato-ternata, parva: foliolis multifidis: lobis ſublinearibus, obtuſis, facie et teneritudine ſequentis. Nectarii cornicula erectiuſcula, brevia, petalis dimidio breviora. Petala lanceolata, longiſſima.*

Nectaria *curva eſſe, ut in A. vulgari, et* Foliorum *lacinias lineares, nec rotundas; habitare in* Helvetiae *dictione Aquilegienſi. Haller.*

An de A. viſcoſa *loquitur Hallerus?* R.

4. AQUILEGIA *canadenſis.*

A. nectariis rectis, ſtaminibus corolla longioribus. *Hort. upſ.* 153. *Mill. dict. n.* 3. *et ic. t.* 47. *Kniph. cent.* 11. *n.* 8.

A. corolla ſimplici, nectariis ferè rectis. *Gron. virg.* 59.

A. pumila praecox canadenſis. *Corn. canad.* 60. *t.* 60. *Lob. ic.* 26.

A. praecox canadenſis, flore externe rubicundo: medio luteo. *Moriſ. hiſt.* 3. *p.* 457. *ſ.* 12. *t.* 2. *f.* 4.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 270.
Canadiſche Akeley. W.
Habitat in Virginia, Canada, Dauria. *Pall. itin.* 3. *p.* 330. ♃ (v. v.)

*5. AQUILEGIA *viridiflora.*

A. nectariis rectis, apice incraſſato ſubinflexo, ſtaminibus corollam ſubaequantibus. *Ait. Kew.* 2. *p.* 248. *Pall. Act. petrop.* 1779. *p.* 260. *t.* 11. *

A. nectariis rectis conniventibus, petalo lanceolato aequalibus. *Murr. comment. goett.* 1780. *p.* 8. *t.* 2. * *Jacq. ic. rar.* 1. *t.* 102. *Collect.* 1. *p.* 35. *

A. nectariis rectis petalo rugoſo longioribus. *Retz. obſ.* 3. *p.* 34. *

β. nectariis coeruleſcentibus. W.
Grünblütige Akeley. W.
Habitat in Sibiria. ♃ (v. v.)

1069. NIGELLA.

Gen. plant. edit. Schreb. n. 935.

Cal. 0. *Petala* 5. *Nectar.* 5, trifida, intra corollam. *Capſ.* 5, convexae.

* *Pentagynae.*

1. NIGELLA *damaſcena.*
N. floribus involucro folioſo cinctis. *Hort. cliff.* 215. *Hort. upſ.* 153. *Roy. lugdb.* 481. *Mill. dict. n.* 2. *et ic. t.* 187. *f.* 2. *Blackw. t.* 558. *Kniph. cent* 10. *n.* 66. *Curt. mag.* 22. *Hoffm. germ.* 188. *Roth. germ. I.* 231. *II.* 593.
N. flore foliis nidulato. *Crantz. auſtr. p.* 133.
N. anguſtifolia, flore majore ſimplici caeruleo. *Bauh. pin.* 145.
Melanthium ſylveſtre. *Matth. dioſc.* 529.
β. Nigella flore majore pleno caeruleo. *Bauh. pin.* 145. *Knorr. del.* 1. *t. S.* 21.
γ. Nigella caule humiliore, pedunculo brevi, petalis albis minoribus. W.
Nigella coarctata. *Hortulan.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 271.
Garten-Nigelle. W.
Habitat inter ſegetes Europae *auſtralis.* ⊙ (v. v.)
Involucrum *approximatum flori*, 5-*phyllum*, *flore longius.* Corolla *ſaepius multiplicata.*

2. NIGELLA *ſativa.*
N. piſtillis quinis, capſulis muricatis ſubrotundis, foliis ſubpiloſis. *Hort. upſ.* 154. *Mat. med.* 139. *Boehm. lipſ.* 173. *Mill. dict. n.* 3. *et. ic. t.* 187. *f.* 1. *Ludw. ect. t.* 80. *Kniph. cent.* 7. *n.* 63. *Hoffm. germ.* 189. *Roth. germ. I.* 232. *II.* 594.
N. flore minore ſimplici candido. *Bauh. pin.* 145.
Melanthium ſativum. *Cam. epit.* 551.
Nigella flore minore pleno et albo. *Bauh. pin.* 146.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 272.
Gemeine Nigelle. W.
Habitat in Germania, Aegypto, Creta. ⊙ (v. v.)

3. NIGELLA *arvenſis.*
E. piſtillis quinis, petalis integris, capſulis turbinatis. *Sp. pl.* 753. *Scop. carn. n.* 657. *Pollich. pal. n.* 514. *Hoffm. germ.* 189. *Roth. germ. I.* 232. *II.* 574.

N. flore foliis nudo, piſtillis corollam aequantibus. *Hort. cliff.* 215. *Roy. lugdb.* 481. *Dalib. pariſ.* 160. *Boehm. lipſ.* 174.
N. flore nudo, ſiliquarum cornubus longiſſimis. *Hall. helv. n.* 1194.
N. flore foliis nudo. *Crantz. auſtr. p.* 132. *n.* 1.
N. arvenſis cornuta. *Bauh. pin.* 145.
Melanthium ſylveſtre alterum. *Cam. epit.* 553.
Houttuyn Lin. Pfl. Syſt. 6. *p.* 274.
Acker-Nigelle. W.
Habitat in Germaniae, Galliae, Italiae *agris.* ☉ (v. v.)
Hujus Capſula inferne anguſtior, praecedentis inferne ventricoſior, ſuperne utraque contra.
Styli *et* Germina *quinque, ſex, ſeptem. Pollich. l. c.* R.

*** Decagynae.*

4. NIGELLA *hiſpanica.*
N. piſtillis denis corollam aequantibus. *Hort. upſ.* 154. *Sauv. monſp.* 253. *Mill. dict. n.* 6. *Desfont. fl. atl.* 1. *p.* 430. *t.* 112. *
N. latifolia, flore majore ſimplici coeruleo. *Bauh. pin.* 145. *prodr.* 75. *Moriſ. hiſt.* 3. *p.* 516. *ſ.* 12. *t.* 18. *f.* 8.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 275.
Spaniſche Nigelle. W.
Habitat in Hiſpania, Monſpelii. ☉ (v. v.)

5. NIGELLA *orientalis.*
N. piſtillis denis corolla longioribus. *Hort. cliff.* 215.* *Hort. upſ.* 153. *Roy. lugdb.* 481. *Mill. dict. n.* 7.
N. chalepenſis lutea, corniculis longioribus. *Moriſ. hiſt.* 3. *p.* 516. *ſ.* 12. *t.* 18. *f.* 10. *Raj. app.* 525.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 275.
Orientaliſche Nigelle. W.
Habitat circum Alepum. ☉ (v. v.)

1070. REAUMURIA.

Gen. plant. edit. Schreb. n. 936.

Cal. 5-phyllus. *Pet.* 5. *Capſ.* 5-locularis, 5-valvis, polyſperma. *Sem.* lanata.

1. REAUMURIA *vermiculata.*
R. foliis ſubulatis ſemiteretibus. W.
R. vermiculata. *Sp. pl.* 754. *Forsk. deſcr.* 101. * *La-*

marck. illuſt. gen. t. 489. *f.* 1. *Desfont. fl. atlant.* 1. *p.* 431. *

Sedum ſiculum vermiculatum, flore ſaxifragae albae, ſemine villoſo. *Bocc. ſicc. t.* 6. *f.* 7. *Moriſ. hiſt.* 3. *p.* 481. *ſ.* 12. *t.* 9. *f.* 6.

Sedum minus arboreſcens vermiculatum. *Lob. ic.* 380.

Sedum minus fruticoſum. *Bauh. pin.* 284.

Kali vermiculatum, albo et amplo ſedi roſei flore. *Barr. ic.* 888.

Kali arabicum primum genus. *Rauw. it.* 37. *t.* 37. *Bauh. pin.* 289.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 276.

Pfriemförmige Reaumürie. W.

Habitat in Aegypti, Syriae, Siciliae *litoribus. De Roque.* ⊙

Statura *Salſolae ſ. Tamariſci.* Folia *ſparſa, linearia, carnoſa, ſubtus convexa, acuta, ſeſſilia, patentia, punctis roridis adſperſa.* Flores *terminales, ſeſſiles inter folia, verſus apicem majora.* *Corollae albae: Lamina ciliata, lateribus petalorum utrinque adnexa, conſtituit nectarium.* Flores *inferi ſunt.*

Capſula 1-*locularis*, 5-*valvis. Forskaehl. l. c.*

*2. REAUMURIA *hypericoides.* W. †

R. foliis ellipticis planis. W.

Reaumuria. *Lamarck. illuſt. gen. t.* 489. *f.* 2.

Hypericum (*alternifolium*) floribus pentagynis, calycibus foliaceis, caule herbaceo, foliis alternis ovatis acutis. *Billard. pl. ſyriac. dec.* 2. *p.* 17. *t.* 10. *

Johanniskrautartige Reaumürie. W.

Habitat in aridis juxta Syriae *deſertum.* ♃

Neque nectaria neque ſemina lanata in deſcriptione indicavit Clariſſ. Billardiere, tamen habitus idem cum Reaumuria vermiculata. W.

Affinitas huic generi eſt maxima cum Hypericis, et ab hoc nectarii praeſentia, ſeminibusque lanatis differre videtur. W.

POLYGYNIA.

1071. DILLENIA.

Gen. plant. edit. Schreb. n. 939.

Cal. 5-phyllus. *Petala* 5. *Capſ.* polyſpermae, connatae, pulpa repletae.

*1. DILLENIA *ſcandens.*

D. foliis lanceolatis integerrimis mucronatis caule ſcandente. W.

Dillenia humilis. *Donn cat. hort. cantab.* 64.

Rankende Dillenie. W.

Habitat in Nova Hollandia. ♄ (v. ſ.)

Frutex *quadripedalis ſcandens, ramis teretiuſculis, junioribus pubeſcentibus.* Folia *alterna petiolata biuncialia et ultra lanceolata utrinque attenuata, integerrima, apice mucronata, venoſa, ſuperne glabra, ſubtus pilis ſparſis tecta.* Flores *ſubſeſſiles ſolitarii terminales magnitudine fere Magnoliae glaucae.* Calycis *foliola ovato-lanceolata magna ſericeo-villoſa.* W.

*2. DILLENIA *integra.*

D. foliis oblongo-obovatis obtuſis ſubſerrulatis obtuſis, pedunculis unifloris. W.

D. (*integra*) foliis obovatis obtuſis ſubintegris, pedunculis unifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 199. *t.* 18.*

Ganzblättrige Dillenie. W.

Habitat in Zeylona. ♄

Arbor *ramis alternis rugoſis fuſcis glabris.* Folia *alterna petiolata obovata obtuſa, a medio ad apicem ſerrulata, ſerraturis obſoletis vixque manifeſtis, utrinque glabra coriacea, ſupra viridia, ſubtus pallidiora, reticulato-venoſa, ſubſpithamaea, palmam lata.* Petioli *ſemiteretes canaliculati villoſi pollicares.* Flores *in ultimis ramulis terminales ſubſolitarii pedunculati.* Thunb. l. c.

3. DILLENIA *ſpecioſa.*

D. foliis oblongis rotundato-acutis denticulatis, pedunculis unifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 200.

Dillenia indica. *Sp. pl.* 745. *Hort. cliff.* 221.

Syalita. *Rheed. mal.* 3. *p.* 39. *t.* 38. 39. *Raj. hiſt.* 1707.

Malva roſea malabarica, Syalita dicta. *Pluk. mant.* 124. *Burm. ind.* 124.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 61.
Prächtige Dillenie. W.
Habitat in Malabaria, Java. ♄

Arbor *excelſa, ramis craſſis rugoſis cinereis glabris.* Folia *alterna petiolata oblongo-rotundata obtuſa cum acumine, undato-denticulata, ſupra glabra, ſubtus obſcura, fuſceſcentia, ſubpedalia, palmam lata.* Petiolus *craſſus brevis vix pollicaris.* Flores *terminales in ramulis ſolitarii pedunculati.* Thunb. l. c.

*4. DILLENIA *elliptica.*

D. foliis elliptico-ovatis acutis ſerratis, pedunculis unifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 200.
Songium. *Rumph. amb.* 2. *p.* 140. *t.* 45.
Elliptiſche Dillenie. W.
Habitat in Amboina, Celebe. ♄

Folia *alterna petiolata elliptica acuminata, argute ſerrata venoſa, venis oppoſitis parallelis.* Flores *terminales ſolitarii pedunculati.* Thunb. l. c.

*5. DILLENIA *ſerrata.*

D. foliis elliptico-ovatis acutis ſerratis, pedunculis trifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 201.
Sangius. *Rumph. amb.* 2. *p.* 142. *t.* 46.
Geſägte Dillenie. W.
Habitat in Java, Celebe. ♄

Folia *alterna petiolata elliptica acuta, argute ſerrata venoſa, venis ſuboppoſitis et alternis parallelis.* Flores *in pedunculis lateralibus terni, pedicellati.* Thunb. l. c.

*6. DILLENIA *pentagyna.*

D. foliis oblongis utrinque acutis ſerratis, pedunculis unifloris aggregatis lateralibus, floribus pentagynis. W.
Dillenia pentagyna. *Roxb. corom.* 1. *p.* 21. *t.* 20. *
Fünfgriffige Dillenie. W.
Habitat in montibus Coromandeliae. ♄

Folia *alterna petiolata pedalia et ultra ſemipedem lata, oblonga utrinque acuta, argute ſerrata utrinque glabra, parallelo-venoſa, venis ſuboppoſitis.* Petioli *biunciales canaliculati baſi dilatati.* Pedunculi *uniflori aggregati* 4, 6, *vel* 9 *e gemmulis lateralibus in ramis praeteriti anni.* Calycis *foliola ſubrotunda ob-*

tuſa corolla duplo breviora. Corolla *flava hujus generis minima, magnitudine Calthae, petalis lanceolatis.* Stamina *numeroſa, quorum decem quadruplo reliquis longiora.* Styli *quinque.* W.

*7. DILLENIA *retuſa.*

D. foliis obovatis remote dentatis, apice truncatis, pedunculis unifloris. W.

D. (*retuſa* foliis obovatis truncatis ſerratis, pedunculis unifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 200. *t.* 19.

Abgeſtutzte Dillenie. W.

Habitat in Zeylonae *ſylvis.* ♄

Arbor *ramis alternis rugoſis fuſcis glabris.* Folia *alterna approximata petiolata obovata, inferne attenuata integra, ſuperne remote ſerrata, ſerraturis obſoletis, apice truncata ſubretuſa coriacea utrinque glabra, venoſa, venis parallelis alternis, ſurſum curvatis, inter venas tenuiſſime reticulata, patentia, palmam fere lata et duplo longiora.* Petioli *ſemiteretes canaliculati baſi hirſuti vix pollicares.* Flos *terminalis ſolitarius pedunculatus.* Thunb. l. c.

Folia *remote dentata baſi et apice integerrima, apice truncata.* W.

*8. DILLENIA *dentata.*

D. foliis ſubrotundis, repando - dentatis, pedunculis multifloris. W.

D. (*dentata*) foliis ovatis retuſis dentatis, pedunculis trifloris. *Thunb. Act. Soc. Lin. Lond.* 1. *p.* 201. *t.* 20. *Vahl. ſymb.* 3. *p.* 72.

Wormia (*triquetra*) petiolis triquetris alternis, latere ſuperiore transverſim ſtriatis. *Rottb. in Nov. Act. Hafn.* 2, *p.* 532. *t.* 3. *

Arbor zeylanica fructu acido. *Burm. zeyl.* 30.

Ghodhapara. *Herm. zeyl.* 49 *Lin. fl. zeyl.* 612.

Rundblättrige Dillenie. W.

Habitat in Zeylona. ♄ (v. ſ. ſ. fl.)

Arbor *ramis alternis rugoſis cinereis glabris.* Folia *alterna petiolata ovata obtuſiſſima ſeu emarginato-retuſa inferne integerrima, ſuperne ad apicem dentata coriacea venoſa, venis ſuboppoſitis parallelis, ſurſum directis, utrinque glabra, ſupra viridia ſubtus obſcuriora palmaria.* Petioli *angulati glabri folio paulo breviores.* Flores *in ramis terminales racemoſi ſubterni.* Pedicelli *alterni glabri ſemipollicares.* Thunb. l. c.

Pedunculi *ſaepe 4-8-flori, baſi ramoſi.* Vahl l. c. Folia *ſubrotundo-elliptica repando-dentata, apice obtuſiſſima et dentibus majoribus inſtructa.* W.

1072. ILLICIUM.

Gen. plant. edit. Schreb. n. 940.

Cal. 6-phyllus. *Petala* 27. *Capſ.* plures in orbem digeſtae, bivalves, 1-ſpermae.

1. ILLICIUM *aniſatum.*
I. petalis interioribus lineari-ſubulatis. W.
I. floribus flaveſcentibus. *Sp. pl.* 664. *Mat. med. p.* 140. *Thunb. jap.* 235. * *Loureiro cochin.* 432. *
Somo vulgo Skimmi. *Kaempf. amoen.* 880. *t.* 881.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 65.
Aechter Sternanis. W.
Habitat in Japonia, China. ♄

2. ILLICIUM *floridanum.* †
I. petalis interioribus lanceolatis. W.
I. floribus rubris. *Mant.* 395. *Ellis act. angl.* 1770. *vol.* 60. *p.* 524. *t.* 12.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 69.
Unächter Sternanis. W.
Habitat in Florida. ♄ (v. ſ.)
Flores *rubri.* Fructus *odoratiſſimi.*

1073. LIRIODENDRON.

Gen. plant. ed. Schreb. n. 941.

Cal. 3-phyllus. *Petala* 9. *Samarae* imbricatae in ſtrobilum.

1. LIRIODENDRON *Tulipifera.*
L. foliis trilobis truncatis, calyce triphyllo. W.
L. foliis lobatis. *Hort. cliff.* 223. *Hort. upſ.* 154. *Gron. virg.* 60. *Roy. lugdb.* 494. *Kalm. it.* 2. *p.* 322. *Trew. ehr. t.* 10. *Buttn. cun.* 229. * *Du Roi. harbk.* 1. *p.* 374. *Wangenh. amer.* 32. *t.* 13. *f.* 32. *Willd. arb.* 173.
Tulipifera Liriodendron. *Mill. dict.*
Tulipifera arbor virginiana. *Herm. lugdb.* 612. *t.* 613.
Tulipifera virginiana, tripartito aceris folio; media laci-

cinia velut abſciſſa. *Pluk. alm.* 379. *t.* 117. *f.* 15. *et t.* 248. *f.* 7. *Cateſb. car.* 1. *p.* 48. *t.* 48. *Raj. hiſt.* 1798. *Du Ham. arb. tom.* 2. *t.* 102.

β. Tulipifera caroliniana, foliis productioribus magis angulofis. *Pluk. alm.* 379. *t.* 68. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 70.

Virginiſcher Tulpenbaum. W.

Habitat in America *ſeptentrionali.* ♄ (v. v.)

*2. LIRIODENDRON *Coco.* †

L. foliis ovatis, calyce triphyllo. W.

L. (*Coco*) foliis ovatis nitidis, floribus ſolitariis, calycibus triphyllis, corollis hexapetalis. *Loureiro cochin.* 424. *

Wohlriechender Tulpenbaum. W.

Habitat in Cochinchina, China. ♄

*3. LIRIODENDRON *Figo.* †

L. foliis lanceolatis, calyce monophyllo ſpathaceo. W.

L. (*Figo*) foliis incurvis reflexis, floribus ſolitariis, calycibus ſpathaceis, corollis hexapetalis. *Loureiro cochin.* 424. *

Krummblättriger Tulpenbaum. W.

Habitat in China. ♄

4. LIRIODENDRON *Liliifera.* †

L. foliis oblongo-lanceolatis, calyce nullo. W.

L. foliis lanceolatis. *Sp. pl.* 755. *Burm. ind.* 124. *Loureiro cochin.* 424. *

Sampacca montana. *Rumph. amb.* 2. *p.* 204. *t.* 69.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 73.

Vielblütiger Tulpenbaum. W.

Habitat in Amboinae *montibus.* ♄

1074. MAGNOLIA.

Gen. pl. ed. Schreb. n. 942.

Cal. 3-phyllus. *Petala* 9. *Capſ.* 2-valves, imbricatae. *Sem.* baccata, pendula.

1. MAGNOLIA *grandiflora.*

M. foliis perennantibus oblongis, petalis obovatis. *Ait. Kew.* 2. *p.* 251. *Willd. arb.* 188.

M. foliis lanceolatis perennantibus. *Sp. pl.* 755.

α. *elliptica* foliis oblongo-ellipticis coriaceis, floribus ſubcontractis. *Ait. l. c.*
M. foliis lanceolatis perſiſtentibus, caule erecto arboreo. *Mill. dict. n. 2. et ic. t.* 172.
M. altiſſima, flore ingenti candido. *Cateſb. car. p.* 80 *t.* 80.
M. maximo flore, foliis ſubtus ſerrugineis. *Trew. ehr. t.* 33.
M. altiſſima, laurocераſi folio, flore ingenti candido. *Amoen. angl. t.* 14.
β. *obovata*, foliis obovato-oblongis, floribus expanſis. *Ait. l. c.*
γ. *lanceolata*, foliis oblongo-lanceolatis apice flexis, floribus ſubcontractis. *Ait. l. c.*
Houttuyn Lin. Pfl. Syſt. 2. *p.* 75.
Groſsblütige Magnolie. W.
Habitat in Florida, Carolina. ♄ (v. v.)
Folia *coriacea, antiquiora purpuraſcenti-pubeſcentia.* Corolla *omnium arborum facile maxima. Munchh. L. B.*

*2. MAGNOLIA *Plumieri.*
M. foliis perennantibus ovato-ſubrotundis utrinque glabris. *Swartz. prod.* 87.
M. ampliſſimo flore albo, fructu coeruleo. *Plum. gen.* 38 *ic. t.* 161.
Plumierſche Magnolie. W.
Habitat in inſulis St. Lucia, Martinica, Guadeloupe. ♄

3. MAGNOLIA *glauca.*
M. foliis ellipticis obtuſis ſubtus glaucis, petalis obovatis. W.
M. foliis ovato-oblongis ſubtus glaucis. *Sp. pl.* 755. *Mill. dict. n.* 1. *Du Roi. harbk.* 1. *p.* 399. *Wangenh. amer.* 60. *t.* 19. *f.* 46. *Willd. arb.* 189.
α. *latifolia*, foliis deciduis. *Ait. Kew.* 2. *p.* 251.
M. foliis ovato-lanceolatis. *Hort. cliff.* 222. *Gron. virg.* 61. *Kalm. it.* 2. *p.* 324.
M. laurifolio ſubtus albicante. *Dill. elth.* 207. *t.* 168. *f.* 205. *Cateſb. car.* 1. *p.* 39. *t.* 39. *Trew. ehr. t.* 9.
Tulipifera virginiana, laurinis foliis averſa parte rore caeruleo tinctis, coni-baccifera. *Pluk. alm.* 379. *t.* 68. *f.* 4.
Laurus tulipifera, baccis calyculatis. *Raj. hiſt.* 1690. *et* 1798. *n.* 4.

β. *longifolia*, foliis perennantibus. *Ait. l. c.*
Houttuyn Lin. Pfl. Syſt. 2. *p.* 77.
Graue Magnolie. W.
Habitat in Virginia, Penſylvania, Carolina. ♄ (v. v.)
Folia *ſubtus albido-coeruleſcentia.*

*4. MAGNOLIA *obovata.*
M. foliis obovatis acutis reticulato-venoſis, petalis oblongis obtuſis. W.
M. (*obovata*) foliis obovatis ſubtus parallelo-nervoſis reticulatis. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 336.
M. (*purpurea*) floribus hexapetalis, petalis extus purpureis. *Curt. mag.* 390. *
Magnolia glauca β. *Thunb. jap.* 237.
Mokkwurea. *Kaempf. amoen.* 845. *Banks icon. Kaempf. t.* 43. 44.
Rothe Magnolie. W.
Habitat in Japonia, China. ♄
Folia *obovato-oblonga integerrima glabra, ſubtus parallelo-nervoſa et tenuiſſime reticulata, nervis villoſis, palmaria usque pedalia.* Thunb. l. c.
Folia *non ſunt nervoſa, ſed reticulato-venoſa.* W.

*5. MAGNOLIA *tomentoſa.*
M. foliis ovatis acutis ſubtus tomentoſis, petalis oblongis obtuſis. W.
M. (*tomentoſa*) foliis ellipticis ſubtus tomentoſis. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 336.
Magnolia glauca α. *Thunb. jap.* 236. *
Kobus. *Bancks. ic. Kaempf. t.* 42.
Filzige Magnolie. W.
Habitat in Japonia. ♄
Folia *elliptico-lanceolata acuta petiolata integerrima, ſupra glabra, ſubtus ſericeo-tomentoſa palmaria.* Pedunculi *breves reflexi ſulcati, craſſi tomentoſi.* Thunb. l. c.

6. MAGNOLIA *acuminata.*
M. foliis ovato-oblongis acuminatis. *Sp. pl.* 756. *Willd. arb.* 189.
M. foliis ovato-lanceolatis acuminatis annuis, petalis obtuſis. *Mill. dict. n.* 4.
M. flore albo, folio majore acuminato haud albicante. *Cateſb. car.* 3. *p.* 15. *t.* 15. *Gron. virg.* 61. *Kalm. it.* 2. *p.* 324.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 79.

Zugeſpitzte Magnolie. W.
Habitat in Penſylvania. ♄ (v. ſ.)
Folia *utrinque viridia dimidio anguſtiora.* Lignum *flavum. Munchhauſ.*

7. MAGNOLIA *tripetala.*
M. foliis lanceolatis, petalis exterioribus dependentibus. *Sp. pl.* 756.
M. (*umbrella*) foliis lanceolato-ovatis umbellatim confertis patentiſſimis, petalis exterioribus dependentibus. *Lamarck. encycl* 3. *p.* 644. *
M. foliis lanceolatis ampliſſimis, petalis exterioribus dependentibus. *Mill. dict. n.* 3.
M. ampliſſimo flore albo, fructu coccineo. *Cateſb. car.* 2. *p.* 61. *t.* 61.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 80.
Dreyblättrige Magnolie. W.
Habitat in Carolina, *rarius in* Virginia. ♄
Flores *magnitudine M. grandiflorae, fragrantiſſimi.* Folia *umbellatim poſita,* 30 *pollices longa. Munchh.*

*8. MAGNOLIA *auriculata.*
M. foliis ſpathulato-ovatis acutis baſi cordatis, petalis unguiculatis. W.
M. (*auriculata*) foliis obovatis baſi attenuatis auriculatis. *Lamarck. encycl.* 3. *p.* 645. *
M. (*Fraſeri*) foliis obovato-lanceolatis baſi auriculatis. *Walt. carol.* 159. *c. icone.*
M. (*pyramidata*) foliis ovalibus oblongis acuminatis baſi auriculatis, ſtrobilo oblongo ovato. *Bartram. itin. ed. germ.* 390.
Geöhrte Magnolie. W.
Habitat in Carolina *et* Georgia, Americes. ♄
Foliis *a reliquis diſtinctiſſima, quae ſpathulato-ovata acuta, baſi cordato-ſagittata.* W.

*1075. NELUMBIUM.

Juſſ. gen. pl. ed. Uſteri p. 76.

Cal. 4-ſ. 5-phyllus. *Cor.* polypetala *Nuces* monoſpermae ſtylo perſiſtenti coronatae, receptaculo truncato immerſae.

1. NELUMBIUM *ſpecioſum.* W.
N. foliis peltatis orbiculatis integerrimis, pedunculis petiolisque muricatis, corolla polypetala. W.

Nymphaea (*Nelumbo*) foliis peltatis undique integris. *Sp. pl.* 730. *Hort. cliff.* 203. *Fl. zeyl.* 193. *Lerche in Nov. Act. A. N. C.* 5. *app. p.* 191. *Loureiro cochin.* 416. * *Thunb. jap.* 223.

Nymphaea indica Faba aegyptia dicta, flore incarnato. *Herm. par.* 205. *t.* 205.

Nymphaea fabifera, indiae paludibus gaudens, foliis umbilicatis amplis, pediculis ſpinoſis, flore roſeo purpureo. *Pluk. alm.* 267. *t.* 322. *f.* 1. *opt.*

Umbilicus aquaticus maximus chinenſis. *Moriſ. hiſt.* 3. *p.* 620. *S.* 15. *t.* 4.

Taratti. *Rumph. amb.* 6. *p.* 168. *t.* 73.

Bem-Tamara. *Rheed. mal.* 11. *p.* 61. *t.* 31.

β. Tamara. *Rheed. mal.* 11. *p.* 59. *t.* 30.

γ. Nymphaea foliis orbiculatis peltatis ſubtus radiatis. *Brown. jam.* 243.

Nymphaea (*Nelumbo*) foliis peltatis undique integris, calyce quadrifido, corolla multiplici alba, loculis monoſpermis. *Walt. carol.* 155.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 214.

Prächtige Nelumbo. W.

Habitat in Chinae, Indiae, Perſiae *aquis.* ♃

Varietas β. *habet* petiolos pedunculos*que glabros*, calyci*sque ſtructura eſt ignota, an ſpecies diſtincta? Varietas* γ. *etiam differre videtur quum* pedunculi *et* petioli *non muricati, et* calyx *quadrifidus ſit, qui tetraphyllus in planta chinenſi.* W.

*2. NELUMBIUM *luteum.* W. †

N. foliis peltatis orbiculatis integerrimis, pedunculis petiolisque glabris, corolla polypetala. W.

Nymphaea Nelumbo β. *Sp. pl.* 730. *Bartr. itin. ed. germ.* 390.

Nymphaea affinis glandifera virginiana umbilicato folio, flore luteo pleno. *Moriſ. hiſt.* 3. *p.* 514. *Raj. ſuppl.* 631.

Nymphaea glandifera virginiana lacteſcens, minori umbilicato folio, pediculo laevi, flore amplo luteo-palleſcente. *Pluk. alm.* 267.

Gelbe Nelumbo. W.

Habitat in Virginiae, Carolinae, Floridae *fluviis.* ♃

Ulterius ab autoptis ut ſequentia examinandum. W.

*3. NELUMBIUM *pentapetalum.* W. †

N. foliis peltatis orbiculatis integerrimis, calyce pentaphyllo, corolla pentapetala. W.

Nymphaea (*pentapetala*) foliis peltatis undique integris, calyce pentaphyllo, corolla magna pentapetala alba, loculis pericarpii monoſpermis. *Walt. carol.* 155.
Fünfblättrige Nelumbo. W.
Habitat in Carolina. ♃

*4. NELUMBIUM *reniforme.* W. †
N. foliis reniformibus, corolla polypetala. W.
Nymphaea (*reniformis*) foliis reniformibus, corollis polypetalis, loculis monoſpermis. *Walt. carol.* 155.
Nierenblättrige Nelumbo. W.
Habitat in Carolina. ♃

Nelumbii genus habitu Nymphaeae eſt proximum, ſed piſtilli ſtructura eſt longe alia. Plerique Botanici crediderunt partem centralem floris Nelumbii eſſe germen unicum cum ſtigmate ſeſſili; at pars illa centralis eſt receptaculum obverſe conicum apice truncatum et foveis excavatum. In qualibet fovea obſervatur germen unicum cum ſtylo brevi. Poſt floreſcentiam receptaculum, reſpectu ſpeciei, evadit ſpongioſum, ſuberoſum vel lignoſum et germina numeroſa mutantur in nuces ſtylo perſiſtenti inſtructae. Nelumbium itaque differt a Nymphaea numero piſtillorum receptaculo inſidentium atque fructu; eſt illi affinitas aliqua quoad fructus ſtructuram cum Fragaria. W.

1076. MICHELIA.

Gen. pl. ed. Schreb. n. 943.

Cal. 3-phyllus. *Petala* 15. *Baccae* multae, 4-ſpermae.

1. MICHELIA *Champaca.*
M. foliis lanceolatis. *Fl. zeyl.* 144. *Oſb. it.* 93.
Sampacca. *Rumph. amb.* 2. *p.* 199. *t.* 67. 68.
Champacam. *Rheed. mal.* 1. *p.* 31. *t.* 19. *Raj. hiſt.* 1641. *Burm. ind.* 124.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 81.
Wohlriechende Michelie. W.
Habitat in India. ♄
Flores *lutei, ſuaveolentiſſimi.*

2. MICHELIA *Tſiampaca.*
M. foliis lanceolato-ovatis. *Mant.* 78.
M. evonymoides. *Burm. ind.* 124.
Sampacca ſylveſtris. *Rumph. amb.* 2. *p.* 202. *t.* 68.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 82.
Geruchloſe Michelie. W.
Habitat in India. ♄
Flores *albicantes ſ. luteſcentes, minus olentes.* Folia *latiora ſeu lanceolato-ovata, petiolata, et etiamnum tenella, more ſalicum, pube ſericea.*

1077. UVARIA.

Gen. plant. ed. Schreb. n. 944.

Cal. 5-phyllus. *Petala* 6. *Baccae* numeroſae, pendulae, 4-ſpermae.

1. UVARIA *zeylanica.*
U. foliis lanceolatis acuminatis, pedunculo unifloro ſolitario, petalis ſubrotundis obtuſis aequalibus. W.
U. foliis integerrimis. *Sp. pl.* 756. *Mat. med. p.* 140.
Uvaria. *Fl. zeyl.* 224. *
Funis muſarius. *Rumph. amb.* 4. *p.* 78. *t.* 42.
Narum-panel. *Rheed. mal.* 2. *p.* 11. *t.* 9. *Raj. hiſt.* 1636.
Houttuyn Lin. Pfl. Syſt. 3. *p.* 84.
Zeylaniſcher Traubenbaum. W.
Habitat in India. ♄ (v. ſ. ſ. fl.)

*2. UVARIA *lanceolata.*
U. foliis lanceolatis integerrimis, floribus axillaribus ſolitariis, ramis virgatis. *Swartz. prod.* 87.
Arbor foliis ovatis glabris utrinque acuminatis, floribus umbellulatis, umbellulis ſparſis. *Brown. jam.* 370.
Lanzettenblättriger Traubenbaum. W.
Habitat in Jamaica. ♄

*3. UVARIA *ceraſoides.*
U. foliis lanceolatis acutis ſubtus pubeſcentibus, pedunculis unifloris ſolitariis, petalis ovatis acutis aequalibus. W.
Uvaria ceraſoides. *Roxb. corom.* 1. *p.* 30. *t.* 33. *
Kirſchartiger Traubenbaum. W.
Habitat in Indiae *orientalis montibus.* ♄
Baccae *monoſpermae atro-rubrae.* W.

*4. UVARIA *ſuberoſa.*
U. foliis oblongis acutis glabris, pedunculis unifloris ſolitariis, petalis tribus interioribus lanceolatis. W.

Uvaria ſuberoſa. *Roxb. corom.* 1. *p.* 31. *t.* 34. *
Korkrindiger Traubenbaum. W.
Habitat in India *orientali.* ♄.

Cortex *ſuberoſus rimoſus.* Folia *breviſſime petiolata, baſi inaequalia.* Corollae *petala tria exteriora ovata viridia duplo interioribus breviora, tria interiora lanceolata albida.* Baccae *atro-rubrae monoſpermae.* W.

*5. UVARIA *tomentoſa.*

U. foliis oblongis acutis tomentoſis, pedunculis unifloris ſolitariis, petalis tribus interioribus ovatis. W.
Uvaria tomentoſa. *Roxb. corom.* 1. *p.* 31. *t.* 35. *
Filziger Traubenbaum. W.
Habitat in Indiae *orientalis montibus.* ♄

Petala *tria exteriora lineari-ſubulata, interioribus triplo breviora vireſcentia; interiora ovata acutiuſcula rufeſcentia.* Baccae *tetraſpermae magnitudine figura et colore Pruni minoris.* W.

*6. UVARIA *odorata.*

U. foliis ovato-lanceolatis, pedunculis unifloris ſolitariis, petalis lineari-lanceolatis longiſſimis. W.
U. (*odorata*) foliis ovato-oblongis acuminatis integerrimis, petalis lanceolato-linearibus acutis planis longiſſimis. *Lamarck. encycl.* 1. *p.* 589. *
Cananga. *Rumph. amb.* 2. *p.* 195. *t.* 65.
Wohlriechender Traubenbaum. W.
Habitat in Java *et* China. ♄

*7. UVARIA *monoſperma.*

U. foliis ellipticis acuminatis ſubtus ferrugineis, pedunculis unifloris aggregatis, petalis ovatis acutis. W.
U. (*monoſperma*) foliis ovato-oblongis acuminatis integerrimis, petalis ovatis acutis, capſulis monoſpermis. *Lamarck. encycl.* 1. *p.* 590. *
Cananga Ouregou. *Aubl. guj.* 1. *p.* 608. *t.* 244. *
Einſamiger Traubenbaum. W.
Habitat in ſylvis remotis Gujanae. ♄

Capſulae *nec Baccae monoſpermae ellipticae, affinis Xylopiae, ſed capſula neque tetragona neque bivalvis.* W.

*8. UVARIA *lutea.*

U. foliis oblongis acutis nitidis, pedunculis trifloris ſolitariis, petalis ovatis obtuſis. W.

Uvaria lutea. *Roxb. corom.* 1. *p.* 32. *t.* 36. *

U. (*coriacea*) foliis lanceolato-ellipticis acuminatis planis, pedunculis oppoſitifoliis ſubbifloris, petalis oblongis. *Vahl ſymb.* 3. *p.* 72.?

Gelber Traubenbaum. W.

Habitat in Indiae *orientalis montibus.* ♄

Folia *coriacea nitida breve petiolata.* Pedunculi *uni-bi-vel triflori ſolitarii foliis oppoſiti.* Petala *rufeſcentia.* Baccae *oblongae baſi attenuatae luteae hexaſpermae magnitudine Pruni ſpinoſae. Uvaria coriacea Clariſſ. Vahlii ab hac non differre videtur.* W.

*9. UVARIA *ligularis.*

U. foliis ovatis acutis, pedunculis multifloris ſolitariis, petalis linearibus acutis longiſſimis. W.

U. (*ligularis*) foliis ovatis acutis integerrimis, petalis linearibus. *Lamarck encycl.* 1. *p.* 590.

Cananga ſylveſtris anguſtifolia. *Rumph. amb.* 2. *p.* 198. *t.* 66. *f.* 2.

Langblumiger Traubenbaum. W.

Habitat in Amboina. ♄.

Accedit quoad petalorum formam ad U. odoratam. W.

*10. UVARIA *longifolia.*

U. foliis lanceolatis margine undulatis, pedunculis umbellatis, petalis lanceolatis acutis. W.

U. (*longifolia*) foliis petalisque lanceolatis, umbellis lateralibus ſubpedunculatis. *Vahl ſymb.* 3. *p.* 72. *

U. (*longifolia*) foliis longis anguſto-lanceolatis acutiſſimis margine undulatis, floribus lateralibus umbellatis, petalis acutis. *Lamarck. encycl.* 1. *p.* 500. *

Sonnerat. itin. 2. *p.* 233. *t.* 131. *

Langblättriger Traubenbaum. W.

Habitat in India *orientali.* ♄

11. UVARIA *japonica.*

U. foliis oblongis acuminatis ſerratis, pedunculis unifloris ſolitariis, petalis ſubrotundis. W.

U. foliis ſerratis. *Sp. pl.* 756. *Thunb. jap.* 237. *

Frutex viſcoſus procumbens, folio telephii vulgaris aemulo, fructu racemoſo. *Kaempf. amoen.* 476. *t.* 477.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 87.

Japaniſcher Traubenbaum. W.

Habitat in Japonia. ♄

Ab omnibus diſtincta: caule *ſcandente,* foliis *ſerratis et* receptaculo *fructuum magno.* W.

1078. ANNONA.

Gen. plant. ed. Schreb. n. 945.

Cal. 3-phyllus. *Petala* 6. *Bacca* polyſperma, ſubrotunda, cortice ſquamato.

1. ANNONA *muricata.*

A. foliis ovali-lanceolatis glabris acutis, fructibus muricatis, petalis ovatis, interioribus obtuſis brevioribus. *Ait. Kew.* 2. *p.* 252.

A. foliis ovali-lanceolatis glabris nitidis planis, pomis muricatis. *Hort. cliff.* 222. * *Gron. virg.* 61. *Roy. lugdb.* 494. *Mill. dict. n.* 2. *Iacq. obſ.* 1. *p.* 10. *t.* 5. *Merian. ſur. t.* 14. *Swartz. obſ.* 220. *

A. foliis oblongo-ovatis nitidis, fructibus ſpinis mollibus tumentibus obſitis. *Brown. jam.* 254.

A. indica latifolia, fructu ſquamoſo aſpero. *Pluk. alm.* 31. *t.* 134. *f.* 2.

A. indica fructu conoide viridi ſquamis veluti aculeato. *Pluk. alm.* 32. *t.* 135. *f.* 2.

A. maxima, foliis latis ſplendentibus, fructu maximo viridi conoide: tuberculis ſ. ſpinulis innocentibus aſpero. *Sloan. jam.* 203. *hiſt.* 2. *p.* 166. *t.* 225.

Anona. *Comm. hort.* 1. *p.* 133. *t.* 69. *Merian. ſurin.* 14. *t.* 14.

Guanabanus fructu molliter aculeato. *Plum. gen.* 43. *t.* 143. *f.* 1.

Araticu Ponhe. *Marcgr. braſ.* 93.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 89.

Stachlicher Flaſchenbaum. W.

Habitat in America *calidiore.* ♄ (v. ſ.)

Cor. 3-*petala, extus ſcabra*: Nectarii *petala* 3, *minora, extus alba laevia, intus rubra.* Fructus *cordatus, muricatus aculeis recurvis, non eſculentus.*

Varietas hujus ſpeciei in Jamaica mihi obvia: foliis *inodoris*, floribus *majoribus fulvis*, fructibus *ſphaericis mucronatis.* Swartz. l. c.

*2. ANNONA *tripetala.*

A. foliis ovatis acutis ſubtus pubeſcentibus, floribus tripetalis, petalis lanceolatis coriaceis tomentoſis. *Ait. Kew.* 2. *p.* 252. *Wendl. obſ.* 24. *t.* 3. *f.* 24.

A. (*Cherimolia*) foliis ovalibus mollibus ſubtus pubeſcentibus, junioribus tomentoſo-ferrugineis, pedunculis tomentoſis flore ſubbrevioribus. *Lamarck. encycl.* 2. *p.* 124. *

A. foliis ovatis acutis, flore albido ungue purpureo, fructu coniformi tuberoſo nigricante. *Trew. Ehret. p.* 16. *t.* 49.
Guanabanus Perſeae folio, flore intus albo, exterius vireſcente, fructu nigricante ſquamato, vulgo Cherimolia. *Feuill. peruv.* 3. *p.* 24. *t.* 17.
Dreyblättriger Flaſchenbaum. W.
Habitat in Peru. ♄ (v. v.)

3. ANNONA *ſquamoſa.*
A. foliis oblongis acutis glabris, fructibus obtuſe ſquamoſis, petalis exterioribus lanceolatis, interioribus minutis. *Ait. Kew.* 2. *p.* 253.
A. foliis oblongis ſubundulatis, fructibus obtuſe ſubſquamatis. *Sp. pl.* 757. *Iacq. obſ.* 1. *p.* 13. *t.* 6. *f.* 1. *Mill. dict. n.* 3. *Swartz. obſ.* 221. *
A. indica, fructu ex viridi-luteo, cortice ſquamato aſpero, nucleis nigricantibus parvis. *Pluk. alm.* 31. *t.* 134. *f.* 3.
A. foliis oblongo-ovatis undulatis venoſis, floribus tripetalis fructibus mammillatis. *Brown. jam.* 256.
Annona tuberoſa. *Rumph. amb.* 1. *p.* 138. *t.* 46.
Annona foliis odoratis minoribus, fructu conoide ſquamoſo parvo dulci. *Sloan. jam.* 205. *hiſt.* 2. *p.* 168. *t.* 227. *Raj. dendr.* 77.
Atamaram. *Rheed. mal.* 3. *p.* 21. *t.* 29.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 95. *t.* 12. *f.* 2.
Schuppiger Flaſchenbaum. W.
Habitat in America *meridionali.* ♄
Fructus *ſubovatus papillis ſubimbricatis obtuſis.*

*4. ANNONA *paludoſa.*
A. foliis oblongis acuminatis tomentoſis ſubtus rufeſcentibus, fructibus muricato-ſquamoſis, petalis ſubrotundo-ovatis acutis interioribus brevioribus. W.
A. (*paludoſa*) foliis villoſis ſupra viridibus, ſubtus rufeſcentibus, fructu luteo tuberculato. *Aubl. guj.* 1. *p.* 611. *t.* 246. *
Guanabanus fructu turbinato minori luteo. *Barrer. aequin.* 53.
Moraſt-Flaſchenbaum. W.
Habitat in pratis paludoſis Gujanae. ♄

5. ANNONA *reticulata.*
A. foliis oblongo-lanceolatis acutis glabris, fructibus

ovatis reticulato-areolatis, petalis exterioribus lanceolatis, interioribus minutis. *Ait. Kew.* 2. *p.* 253.

A. foliis lanceolatis, fructibus ovatis reticulato-areolatis. *Sp. pl.* 757. *Mill. dict. n.* 1. *Iacq. obſ.* 1. *p.* 14. *t.* 6. *f.* 2. * *Swartz. obſ.* 222. *

A. foliis oblongis undulatis venoſis, fructibus areolatis. *Brown. jam.* 256.

Anona maxima foliis oblongis anguſtis, fructu maximo luteo conoide: cortice glabro in areolas angulares diſtincto. *Sloan. jam.* 264. *hiſt.* 2. *p.* 167. *t.* 226. *Cateſb. car.* 2. *p.* 86. *t.* 86. *Raj. dendr.* 77.

Anona maram. *Rheed. mal.* 3. *p.* 23. *t.* 30. 31.

Guanabanus fructu aureo et molliter aculeato. *Plum. gen.* 43.

β. A. (*mucoſa*) foliis oblongis, fructibus areolatis, corollis extimis monopetalis. *Iacq. obſ.* 1. *p.* 16. *

Anona. *Rumph. amb.* 1. *p.* 136. *t.* 45.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 98.

Netzfrüchtiger Flaſchenbaum. W.

Habitat in America *meridionali.* ♄ (v. ſ. ſ. fl.)

*6. ANNONA *longifolia.*

A. foliis lanceolatis acuminatis, fructibus ſubrotundis punctatis et reticulato-areolatis, petalis ovato-lanceolatis, interioribus minoribus. W.

A. (*longifolia*) fructu rubro punctato et reticulato. *Aubl. guj.* 1. *p.* 615. *t.* 248. *

Langblättriger Flaſchenbaum. W.

Habitat in Gujana. ♄

*7. ANNONA *punctata.*

A. foliis oblongis acuminatis, fructibus globoſis ſtellato-punctatis glabris, petalis ovatis acutis, interioribus duplo minoribus. W.

A. (*punctata*) fructu laevi punctato carne rubeſcente. *Aubl. guj.* 1. *p.* 614. *t.* 247. *

Punktirter Flaſchenbaum. W.

Habitat in ſylvis Gujanae. ♄

8. ANNONA *hexapetala.*

A. foliis elliptico-oblongis acutis glabris, petalis ſpathulatis aequalibus acutis. *Ait. Kew.* 2. *p.* 253.

A. floribus hexapetalis. *Lin. ſuppl.* 270.

Sechsblättriger Flaſchenbaum. W.

Habitat in China *et* India *orientali.* ♄

Arbor *ſimillima pluribus Annonis.* Folia *elliptica*

glabra. Calyx *triphyllus minutus.* Petala *ſex aequalia oblonga.* Fructus *mihi ignotus.*

9. ANNONA *paluſtris.*

A. foliis oblongis obtuſiuſculis glabris, fructibus areolatis, petalis exterioribus majoribus ſubrotundis. W.

A. foliis oblongis obtuſiuſculis glabris, fructibus areolatis. *Sp. pl.* 757. *Swartz. obſ.* 223. *

A. uliginoſa, foliis nitidis ovatis, fructibus areolatis odoratis. *Brown. jam.* 256.

A. foliis oblongis obtuſis glabris, fructu rotundo cortice glabro. *Mill. dict.*

Anona aquatica, foliis laurinis atro-virentibus, fructu minore conoide luteo, cortice glabro in areolas diſtincto. *Sloan. jam.* 205. *hiſt.* 2. *p.* 169. *t.* 228. *f.* 1. *Raj. dendr.* 78.

Anona americana juxta fluviorum ripas innaſcens pyriformi fructu. *Pluk. alm.* 32. *t.* 240. *f.* 6.

Guanabanus paluſtris fructu laevi viridi. *Plum. mſſ.*

Houttuyn Lin. Pfl. Syſt. 3. *p.* 102.

Sumpf-Flaſchenbaum. W.

Habitat in America *ad aquas.* ♄ (v. ſ.)

10. ANNONA *glabra.*

A. foliis lanceolato-ovatis, fructibus conoidibus glabris. *Sp. pl.* 758. *Du Roi. harbk.* 1. *p.* 62.

Anona maxima, foliis latis, fructu maximo luteo conoide: cortice glabro. *Cateſb. car.* 2. *p.* 64. *t.* 64.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 103.

Glatter Flaſchenbaum. W.

Habitat in Carolina. ♄

11. ANNONA *triloba.*

A. foliis ellipticis acutis glabris, floribus pendulis campanulatis, calycibus ovatis, petalis pluribus ovalibus. *Ait. Kew.* 2. *p.* 254.

A. foliis lanceolatis, fructibus trifidis. *Sp. pl.* 758. *Mill. dict. n.* 8. *et ic. t.* 35. *Du Roi. herbk.* 1. *p.* 59.

Anona fructu luteſcente levi ſcrotum arietis referente. *Cateſb. car.* 2. *p.* 85. *t.* 85. *Trew. ehr. t.* 5. *Du Ham. arb.* 1. *p.* 56. *t.* 19.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 103.

Dreylappiger Flaſchenbaum. W.

Habitat in Carolina. ♄

12. ANNONA *aſiatica.*

A. foliis lanceolatis glabris nitidis lineatis. *Hort. cliff.*

222. * *Fl. zeyl.* 225. * *Roy. lugdb.* 494. *Mill. dict. n.* 7. *Vahl. ſymb.* 3. *p.* 73.
A. (*glabra Keſchta*) foliis ellipticis, fructu glabro. *Forsk. deſcr.* 102. * *Icon. tab.* 15.
A. ſylveſtris. *Burm. zeyl.* 21.
Guanabanus fructu purpureo. *Plum. ſpec.* 43. *ic.* 143. *f.* 2.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 105.
Aſiatiſcher Flaſchenbaum. W.
Habitat in Zeylona. ♄ (v. ſ. ſ. fl.)

*13. ANNONA *Ambotay.*
A. foliis oblongo-ovatis acuminatis ſubtus tomentoſis rufeſcentibus, petalis ovatis acutis ſubaequalibus. W.
A. (*Ambotay*) foliis amplis ſubtus villoſis et rufeſcentibus. *Aubl. guj.* 1. *p.* 616. *t.* 249. *
Braunhaariger Flaſchenbaum. W.
Habitat in ſylvis Gujanae. ♄
Frutex *octopedalis.* Folia *octopollicaria coriacea oblongo-ovata acuminata venoſa, venis margine arcuato-confluentibus, ſuperne viridia glabra, inferne tomento rufeſcente induta.* Petioli *breves ut ramuli juniores flexuoſi, tomentoſi.* Pedunculi *uniflori ſolitarii axillares vix longitudine petioli.* Flos *parvus vireſcens.* Fructus *ignotus.* W.

14. ANNONA *africana.*
A. foliis lanceolatis pubeſcentibus. *Hort. cliff.* 222. *Roy. lugdb.* 494. *Mill. dict. n.* 6.
Guanabanus fructu ſubcoeruleo. *Plum. gen.* 43.
Houttuyn Lin. Pfl. Syſt. 2. *p.* 105.
Africaniſcher Flaſchenbaum. W.
Habitat in America. ♄

*15. ANNONA *pygmaea.* †
A. foliis lanceolatis acutis, petalis oblongis acutis interioribus longioribus. W.
Annona pygmaea. *Bartram itin. ed. germ. p.* 21. *t.* 1.
Zwerg-Flaſchenbaum. W.
Habitat in Florida. ♄
Frutex *pedalis, ramis depreſſis.* Folia *alterna lanceolata acuta baſi in petiolum attenuata venoſa glabra.* Pedunculi *uniflori axillares ſolitarii nutantes.* Corolla *campanulata magnitudine A. ſquamoſae, petalis oblongis acutis apice reflexis, interioribus longioribus.* Fructus *ignotus.* W.

*16. ANNONA *obovata*. W.

A. foliis obovatis obtuſiuſculis, fructu oblongo glabro, petalis obovatis obtuſis exterioribus majoribus. W.

Annona grandiflora. *Bartram itin. ed. germ. p.* 20. *tab.* 2.

Groſser Flaſchenbaum. W.

Habitat in Florida. ♄

Folia *obovata obtuſiuſcula breve petiolata glabra.* Flos *ſolitarius terminalis nutans albus duplo major quam in A. muricata,* petalis *obovatis, interioribus brevioribus.* Fructus *oblongo-obovatus glaber.* W.

*17. ANNONA *grandiflora.*

A. foliis ovato-oblongis, fructu elliptico ſcabriuſculo, petalis pubeſcentibus oblongis interioribus brevioribus. W.

A. (*grandiflora*) foliis ovato-oblongis petiolatis. *Vahl ſymb.* 3. *p.* 73.

A. (*grandiflora*) foliis ovato-lanceolatis glabris maximo flore. *Lamarck. encycl.* 2. *p.* 126. *

Groſsblütiger Flaſchenbaum. W.

Habitat in Madagaſcaria *et inſula* Borboniae. ♄

Rami *glabri punctati.* Folia *petiolata alterna palmaria parum attenuata acuta, venoſo-reticulata, utrinque glabra, ſupra nitida.* Petiolus *breviſſimus rugoſus.* Calycis *foliola tomentoſa ovata unguicularia.* Pedunculi *breves axillares vel laterales ſolitarii uniflori.* Petala *exteriora bipollicaria, pollicem lata coriacea oblonga, baſi parum anguſtiora, extus villis denſis adpreſſis cinerea, intus glabra, excepto margine cinereo.* Vahl l. c.

*18. ANNONA *amplexicaulis.*

A. foliis cordato-oblongis amplexicaulibus. *Vahl ſymb.* 3. *p.* 73.

A. (*amplexicaulis*) foliis cordato-oblongis acutis diſtichis amplexicaulibus. *Lamarck. encycl.* 2. *p.* 127. *

Herzblättriger Flaſchenbaum. ♄

Habitat in Madagaſcar *et in inſula* Mauritii. ♄

Rami *teretes glabri, cortice cinereo rimoſo.* Folia *alterna amplexicaulia cordato-oblonga ſenſim attenuata, dodrantalia et ultra bifaria venoſo-reticulata acutiuſcula, ſubtus in ſpecimine ſicco purpureo-violacea.* Pedunculi *axillares breves uniflori ſolitarii.* Flores *pollicares vel parum ultra.* Petala *exteriora lanceolata obtuſa.* Vahl l. c.

1079. XYLOPIA.

Gen. plant. edit. Schreb. n. 946.

Cal. 3-phyllus. *Petala* 6. *Capſulae* ſubmonoſpermae tetragonae bivalves. *Semina* arillata.

1. XYLOPIA *muricata.*

X. foliis lanceolatis acuminatis, ſubtus ſtrigoſis, apice barbatis, pedunculis multifloris, fructibus muricatis. W.

X. pedunculis multifloris, fructibus muricatis. *Sp. pl.* 1367.

Xylopicrum fruticoſum, foliis ovatis acuminatis productis alternis, capſulis punctatis, floribus confertis ad alas. *Brown. jam.* 250. *t.* 5. *f.* 2.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 259.

Bärtiges Bitterholz. W.

Habitat in Jamaica. ♄ (v. ſ.)

*2. XYLOPIA *fruteſcens.*

X. foliis lanceolatis acuminatis, ſubtus ſericeis, apice glabris, pedunculis ſubmultifloris, fructibus glabris. W.

X. (*fruteſcens*) foliis lanceolatis ſubtus candicantibus villoſis, capſulis rubris quadragonis. *Aubl. guj.* 1. *p.* 602. *t.* 292. * (*excluſis ſynonymis.*)

Ibira. *Piſ. braſ.* 145.

Strauchartiges Bitterholz. W.

Habitat in Gujana, Braſilia. ♄ (v. ſ.)

Simillima praecedenti, ſed diſtincta: foliis *anguſtioribus ſubtus albicantibus ſericeo-villoſis, nec ſtrigoſis, apice glabris, nec faſciculo pilorum inſtructis, fructibus glabris.* Pedunculi *ſunt uni- vel multiflori.* W.

3. XYLOPIA *glabra.*

X. foliis oblongo-ovatis glabris, pedunculis unifloris ſubgeminatis, fructibus glabris. W.

X. pedunculis ſubunifloris, fructibus glabris. *Sp. pl.* 1367.

Xylopicrum foliis amplioribus nitidis ovatis, petiolis brevibus, fructibus glabris. *Brown. jam.* 251.

Xylopicron arbor barbadenſibus lignum amarum nominata. *Pluk. alm.* 395. *t.* 238. *f.* 4.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 260.

Glattes Bitterholz. W.
Habitat in Jamaica *et* Barbados. ♄

1080. U N O N A.

Gen. pl. ed. Schreb. n. 947.

Cal. 3-phyllus. *Petala* 6. *Baccae* 2-vel 3-ſpermae articulato-moniliformes.

1. UNONA *diſcreta.*
U. foliis lanceolatis ſubtus ſericeis. *Vahl ſymb.* 2. *p.* 63.
Unona diſcreta. *Lin. ſuppl.* 270.
Weidenblättrige Unone. W.
Habitat in Surinamo. *C. G. Dalberg.* ♄

Arbor *ramis virgatis anguſtis flexilibus.* Folia *bifaria alterna petiolata liguſtrina glabra integerrima digitalia.* Petioli *breviſſimi.* Pedicelli *baccarum intra corollam umbellati.* Flos *Annonae ſed fructus diverſus purpureus ſapidus aromaticus, qui diſtinguit genere uti in Theobroma et Abroma.*

Rami *pubeſcentes.* Folia *ſesquipollicaria valde attenuata anguſta ſalicina vix petiolata.* Vahl l. c.

*2. UNONA *tomentoſa.* W. †
U. foliis lanceolatis tomentoſis. W.
Desmos (*cochinchinenſis*) foliis lanceolatis tomentoſis, floribus ſolitariis clauſis. *Loureiro cochin. p.* 431. *
Filzige Unone. W.
Habitat in Cochinchinae *dumetis.* ♄

*3. UNONA *diſcolor.*
U. foliis ovato-oblongis utrinque glabris. *Vahl ſymb.* 9. *p.* 63. *t.* 63. *
Desmos (*chinenſis*) foliis ovato-lanceolatis glabris, floribus ſolitariis patentibus. *Loureiro cochin.* 431?
Verſchiedenfarbige Unone. W.
Habitat in India *orientali.* ♄

Folia *petiolata alterna tripollicaria, pollices 2 lata, baſi rotundata, apice parum acuminata, obtuſiuſcula integerrima membranacea, utrinque glabra, ſubtus glauca, tenella ſubtus villoſo-ſericea.* Vahl l. c.

*4. UNONA *concolor.* W. †
U. foliis oblongis acuminatis utrinque glabris concoloribus, pedunculis bifloris. W.
Uvaria zeylanica. *Aubl. guj.* 1. *p.* 605. *t.* 243. * (*excluſis ſynonymis.*)

Gleichfarbige Unone. W.
Habitat in Gujanae *ſylvis.* ♄

Unonae genus ad Uvariam amandavit Clariſſ. Swartz. obſ. p. 224, *ſed fructus moniliformis eſt nota, qua facile Unona a congeneribus diſtinguitur.* W.

1081. ANEMONE.

Gen. plant. edit. Schreb. n. 948.

Cal. 0 *Petala* 6-9. *Sem.* plura.

* *Hepaticae flore ſubcalyculato.*

1. ANEMONE *Hepatica.*

A. foliis trilobis integerrimis. *Mat. med.* 140. *Scop. carn. n.* 658. *Pollich. pal. n.* 515. *Hoffm. germ.* 189. *Fl. dan. t.* 610. *Ludw. ect. t.* 140. *Roth. germ. I.* 235. *II.* 602.

Hepatica. *Hort. cliff.* 223. *Fl. ſuec.* 445, 480. *Hort. upſ.* 155. *Gron. virg.* 61. *Roy. lugdb.* 487. *Hall. helv. n.* 1156. *Scop. carn. ed.* 1. *p.* 567. *Mill. dict. n.* 1. — 5. *Blackw. t.* 207. *Knorr. del.* 1. *t. L.* 13. *Crantz. auſtr. p.* 121.

Trifolium hepaticum, flore ſimplici et pleno. *Bauh. pin.* 339.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 280.

Dreylappige Anemone. W.

Habitat in Europae *nemoribus lapidoſis.* ♃ (v. v.)

** Pulſatillae *pedunculo involucrato, ſeminibus caudatis.*

2. ANEMONE *patens.*

A. pedunculo involucrato, foliis digitatis multifidis. *Gmel. ſib.* 4. *p.* 195. *Pall. itin.* 2. *p.* 31. *Beyg. ged.* 1. *p.* 141. *n.* 2. *Mattuſch. ſil. n.* 390. *Hoffm. germ.* 189. *Roth. germ. II.* 603.

Pulſatilla patens. *Mill. dict. n.* 4.

Pulſatilla folio anemones ſecundae ſ. ſubrotundo. *Bauh. pin.* 94. *prodr.* 94.

Pulſatilla polyanthos violacea, anemones folio. *Breyn. cent.* 132. *Helw. pulſ.* 52. *t.* 2. 3.

Pulſatilla, anemones folio diſſecto lanuginoſo, flore majore dilute luteo patente. *Amm. ruth.* 104.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 282.

Ausgebreitete Anemone. W.

Habitat in Tobolſko Sibiriae, Luſatia *inferiore*, Gedani, Sileſia. ♃ (v. ſ.)

Folia *radicalia ternato-digitata: Foliolis ſeſſilibus, laciniatis, acuminatis: foliolo intermedio tripartito; lateralibus bipartitis.* Scapus *uniflorus, foliis brevior.* Involucrum *remotum, villoſum.* Corolla *alba, ſubtus villoſa.* Stamina *lutea.*

*3. ANEMONE *cernua.*

A. pedunculo involucrato, foliis pinnatis, flore cernuo. *Thunb. jap.* 238.

Hängende Anemone. W.

Habitat juxta Jedo *in* Japonia. ♃

Scapus *tomentoſus medio involucratus uniflorus apice geniculatus.* Folia *radicalia petiolata pinnata digitalia;* Pinnae *oppoſitae profunde inciſae, pinnulae alternae ſubſerratae acutae villoſae, imprimis ſubtus. Differt ab A. vernali:* flore *cernuo,* foliorum *pinnis pluribus magis inciſis.* Thunb. l. c.

4. ANEMONE *vernalis.*

A. pedunculo involucrato, foliis pinnatis, foliolis tripartitis obtuſis glabriuſculis, flore erecto. W.

A. pedunculo involucrato, foliis pinnatis, flore erecto. *Oed. dan. t.* 29. *Pollich. pal. n.* 516. *Mattuſch. ſil. n.* 391. *Reyg. ged.* 2. *p.* 95. *Hoffm. germ.* 189. *Roth. germ. I.* 235. *II.* 603.

A. tubis caudatis, involucris multifidis, foliis pinnatis, pinnis latis lobatis. *Hall. helv. n.* 1147. *t.* 21.

Anemone ſulphurea. *Allion. ped. n.* 1921.

Pulſatilla vernalis. *Mill. dict. n.* 3.

Pulſatilla foliis ſimpliciter pinnatis: foliolis lobatis, flore erecto. *Fl. ſuec.* 438, 483.

Pulſatilla involucro piloſo nitido. *Fl. lapp.* 223.

Pulſatilla, apii folio, vernalis, flore majore. *Bauh. pin.* 177. *prodr.* 94. *Helw. pulſ.* 63. *t.* 9.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 284.

Frühlings-Anemone. W.

Habitat in Sueciae, Helvetiae, Germaniae *ſylvis ſteriliſſimis.* ♃ (v. v.)

*5. ANEMONE *Halleri.*

A. pedunculo involucrato, foliis pinnatis, foliolis tripartitis acuminatis villoſis, flore erectiuſculo. W.

Anemone Halleri. *Allion. pedem. n.* 1922. *t.* 80. *f.* 2.

A. (*Halleri*) foliis pinnatis, foliolis multifidis. *Villars. delph.* 3. *p.* 725.

A. tubis caudatis, involucris multifidis, foliis hirſutis pinnatis, pinnis acute lobatis. *Hall. helv. n.* 1148.

Hallerſche Anemone. W.

Habitat in alpibus Valleſiae, Sabaudiae, Delphinatus. ♃ (v. ſ.)

Similis A. vernali, ſed diverſa: foliis *valde villoſis,* pinnis *profunde trifidis,* laciniis *lanceolatis acuminatis.* W.

6. ANEMONE *Pulſatilla.*

A. pedunculo involucrato: petalis rectis, foliis bipinnatis. *Sp. pl.* 759. *Mat. med.* 141. *Gmel. ſib.* 4. *p.* 195. *Pollich. pal. n.* 517. *Mattuſch. ſil. n.* 392. *Fl. dan. t.* 153. *Ludw. ect. t.* 119. *Kniph. cent.* 1. *t.* 4. *Knorr. del* 2. *t. A.* 7. *Hoffm. germ.* 190. *Roth. germ. I.* 235. *II.* 603.

A. tubis caudatis, involucris unifoliis multifidis, foliis linearibus hirſutis, pinnis tri-et quinquepartitis. *Hall. helv. n.* 1146.

A. caule involucrato, flore nutante, limbo erecto. *Crantz. auſtr. p.* 123. *n.* 6.

A. ſeminibus caudatis, petalis lanceolatis, rectis; foliis bipinnatis. *Scop. carn. ed.* 1. *p.* 568. *ed.* 2. *n.* 665.

Pulſatilla vulgaris. *Mill. dict. n.* 1.

Pulſatilla foliis decompoſitis pinnatis, flore nutante: limbo erecto. *Hort. cliff.* 223. *Fl. ſuec.* 446, 481. *Roy. lugdb.* 487. *Dalib. pariſ* 160. *Sauv. monſp.* 254.

Pulſatilla folio craſſiore et majore flore. *Bauh. pin.* 177. *Helw. pulſ.* 62. *t.* 8.

Pulſatilla. *Cam. epit.* 398.

β. flore minori paluſtris. *Bauh. pin.* 176. *Hall.* R.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 284.

Violette Anemone. W.

Habitat in campis ſylveſtribus exaridis collibusque apricis Europae *borealis.* ♃ (v. v.)

Datur *varietas floribus albis.* *Gmel. Hall.* R.

7. ANEMONE *pratenſis.*

A. pedunculo involucrato, petalis apice reflexis, foliis bipinnatis. *Sp. pl.* 760. *Reyg. ged.* 2. *p.* 95. *Mattuſch. ſil. n.* 393. *Fl. dan. t.* 611 *Kniph. cent.* 1. *n.* 3. *Hoffm. germ.* 190. *Roth. germ. I.* 235. *II.* 604.

A. caule involucrato, flore pendulo limbo reflexo. *Crantz. auſtr. p.* 123. *n.* 7.

Pulſatilla foliis decompoſitis pinnatis, flore pendulo: limbo reflexo. *Hort. cliff.* 223. * *Fl. ſuec.* 447, 482. *Roy. lugdb.* 487. *Sauv. monſp.* 254.
Pulſatilla pratenſis. *Mill. dict. n.* 2.
Pulſatilla flore minore nigricante. *Bauh. pin.* 177. *Helw. pulſ.* 66. *t.* 12.
Pulſatilla vulgaris, ſaturatiore flore. *Cluſ. hiſt.* 1. *p.* 246.
Pulſatilla flore clauſo obſoleto: petalis reflexis. *Helw. pulſ.* 64. *t.* 11.
Houttuyn Lin. Pſl. Syſt. 7. *p.* 286.
Wieſen-Anemone. W.
Habitat in Scaniae, Germaniae *campis pratenſibus apricis aridis.* ♃ (v. v.)
Diſtincta ſpecies a pracedenti.

* * * Anemones *caule folioſo, Seminibus caudatis.*

8. ANEMONE *alpina.*

A. foliis caulinis ternis connatis ſupradecompoſitis glabriuſculis, foliolis pinnatifidis obtuſis, ſeminibus hirſutis caudatis. W.
A. foliis caulinis ternis connatis ſupradecompoſitis multifidis, ſeminibus hirſutis caudatis. *Sp. pl.* 760. *Iacq. auſtr. t.* 85. *Mattuſch. ſil. n.* 394. *Hoffm. germ.* 190. *Roth. germ. I.* 235. *II.* 605.
A. foliis triplicato-pinnatis, acute circumſerratis, caulinis ternatis, tubis caudatis. *Hall. helv. n.* 1149.
A. burſeriana. *Scop. carn. n.* 664.
A. caule involucrato, ſeminibus piloſis, caudatis. *Crantz. auſtr. p.* 105. *n.* 8. *t.* 3. *f.* 2.
A. ſylveſtris 2. *Cluſ. hiſt.* 1. *p.* 245.
A. flore candido. *Thal. herc. p.* 15.
Pulſatilla flore albo. *Bauh. pin.* 177. *Lob. ic.* 182. *Bauh. hiſt.* 3. *p.* 419.
β. Anemone alpina alba major. *Bauh. pin.* 176. *prodr.* 94. *Burſ. IX.* 80.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 287.
Alpen-Anemone. W.
Habitat in alpibus Styriacis, Helveticis, Auſtriacis, Sileſiacis, Hercynicis. ♃ (v. v.)
Folia *caulina baſi lanata.* Involucri *folia ſupradecompoſita uti radicalia.* Flos *magnitudine flore Tulipae, albus extus ſubviolaceus pubeſcens.*

9. ANEMONE *apiifolia.*

A. foliis caulinis ternis connatis ſupradecompoſitis villoſis, foliolis pinnatifido-dentatis acuminatis, ſeminibus hirſutis caudatis. W.

Anemone apiifolia. *Wulffen in Iacq. miſc.* 2. *p.* 77. *t.* 4. * *Scop. carn. n.* 663. *Hall. helv. n.* 1149. β.

A. (*apiifolia*) foliis caulinis ternis connatis ſupradecompoſitis multifidis tenuiſſimis ſubtus piloſiſſimis. *Murr. ſyſt. veg. ed.* 14. *p.* 510.

A. (*myrrhidifolia*) foliis triangulariter decompoſitis, laciniis confluentibus, petalis acutis, tubis rectis. *Villars. delph.* 3. *p.* 727. *

A. (*ſulphurea*) pedunculo involucrato, foliis triplicato-pinnatis piloſis planis acute inciſis, ſeminibus caudatis. *Mant.* 78.

A. ſylveſtris tertia. *Cluſ. hiſt.* 1. *p.* 245.

Pulſatilla lutea apii hortenſis folio. *Bauh. hiſt.* 3. *p.* 411.

Pulſatilla lutea paſtinacae ſylveſtris folio. *Bauh. pin.* 177.

Pulſatilla tertia alpina. *Dalech. hiſt.* 851.

Pulſatilla lutea. *Camer. epit.* 393. *Geſn. ic. aen.* 18.

Selleryblättrige Anemone. W.

Habitat in alpibus Helvetiae, Delphinatus, Sabaudiae, Carnioliae. ♃ (v. ſ.)

Differt ab A. alpina ut A. Halleri ab A. vernali. W

* * * * Anemoides *caule folioſo, ſeminibus ecaudatis.*

10. ANEMONE *coronaria.*

A. foliis radicalibus ternato-decompoſitis, dentibus mucronatis, involucro folioſo, ſeminibus lanatis. W.

A. foliis radicalibus ternato-decompoſitis, involucro folioſo. *Sp. pl.* 760. *Haſſelq.* 477. *Hill. anat.* 12. *Mill. dict. n.* 5.

Pulſatilla foliis decompoſitis ternatis. *Hort. cliff.* 223. *Hort. upſ.* 155. *Roy. lugdb.* 487.

Anemone tenuifolia, ſimplici flore. *Bauh. pin.* 174. *n* 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12.

β. A. tenuifolia multiplex rubra. *Bauh. pin.* 176.

Anemone. *Cam. epit.* 386.

γ. A. anguſtifolia multiplex, mutata florum facie ...otannis nova. H. R. P. *Mill. dict. t.* 31.

Houttuyn Lin. Pfl. Syſt. 2. *p.* 288.

Gemeine Anemone. W.

Habitat in Oriente, Conſtantinopoli *alla*... ♃ (v. v.)

11. ANEMONE *hortenſis.*

A. foliis radicalibus digitatis, laciniis trifidis, caulinis ternis lanceolatis connatis ſubdiviſis, ſeminibus lanatis. W.

A. foliis digitatis, ſeminibus lanatis. *Sp. pl.* 761. *Mill. dict. n.* 6.

A. ſeminibus lanatis, foliis radicalibus trilobatis et multifidis, caulinis ovato-lanceolatis. *Hall. helv. n.* 1152.

A. foliis digitatis: lobis inciſis: petalis lanceolatis numeroſis. *Ger. prov.* 380.

Pulſatilla foliis digitatis. *Hort. cliff.* 224.

Anemone hortenſis latifolia. *Bauh. pin.* 176.

A. geranii rotundo folio, purpuraſcens. *Bauh. pin.* 173.

A. hortenſis latifolia 3. *Cluſ. hiſt.* 1. *p.* 249.

Anemone 1. *Dod. pempt.* 434.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 289.

Schöne Anemone. W.

Habitat in Italia, Helvetia. ♃ (v. v.)

Involucrum *foliolis lanceolatis.* Petala 9, *lineata, extus pubeſcentia.*

12. ANEMONE *palmata.*

A. foliis cordatis trilobis, involucris triphyllis trifidis, corollis ſubdecapetalis. *Vahl ſymb.* 3. *p.* 73. *

A. foliis rotundatis lobatis crenatis, involucro multifido, petalis exterioribus villoſis majoribus. *Desfont. fl. atl.* 1. *p.* 432. *

A. foliis cordatis ſublobatis, calyce hexaphyllo colorato. *Sp. pl.* 758.

Pulſatilla foliis palmatis. *Hort. cliff.* 224. *Roy. lugdb.* 487.

Anemone cyclaminis ſ. malvae folio, lutea. *Bauh. pin.* 173. *Moriſ hiſt.* 2. *p.* 425. *ſ.* 4. *t.* 25. *f.* 3.

A. latifolia flava. *Bauh. pin.* 176.

A. hortenſis latifolia 1. *Cluſ. hiſt.* 1. *p.* 248. *Barr. ic.* 792.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 289.

Schlitzblättrige Anemone. W.

Habitat in Luſitania *ad* Tagum, *inque* Algeria. ♃

Involucrum *triphyllum, foliola ſeſſilia baſi ſubconnata, extrorſum dilatata, glabro-nervoſa trifida, laciniis lineari-lanceolatis lateralibus interdum bifidis.* Vahl l. c.

13. ANEMONE *ſibirica.*
A. caule unifloro, involucro folioſo obtuſo. *Sp. pl.* 763. *Gmel. ſib.* 4. *p.* 199. *n.* 41.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 290.
Sibiriſche Anemone. W.
Habitat in Sibiria. *D. Gmelin.* ♃
Folia *ſuborbiculata, ternata: foliola inciſo-palmata, ciliata, petiolis hirtis.* Scapus *nudus, piloſus.* Involucrum *ſub flore tripartitum, lobatum, laciniis lanceolatis.* Flos *patens: petalis* 6, *ſubrotundis, fulvis.* Stamina *flava.* Piſtillorum *capitula ſubrotunda, muricata.*

*14. ANEMONE *baldenſis.*
A. foliis biternatis, foliolis tripartitis, laciniis trifidis, involucro folioſo, ſeminibus lanatis ſtylo perſiſtenti breviſſimo. W.
A. foliis biternatis hirſutis. *Mant.* 78. *Allion pedem. n.* 1928. *t.* 44. *f.* 9. *et t.* 67. *f.* 2. *Villars. delph.* 3. *p.* 723. *t.* 49. *
A. (*fragifera*) pedunculo nudo, ſeminibus ſubrotundis lanatis muticis, foliorum laciniis acutis lanceolatis. *Murr. ſyſt. veg. ed.* 14. *p.* 510. *Iacq. ic. rar.* 1. *t.* 103. *Miſc.* 2. *p.* 55. *
Anemone alpina. *Scop. carn. n.* 662. *t.* 26.
A. ſeminibus lanatis, foliis caulinis ternatis triplicato-trifidis, lobulis lanceolatis acute trifidis bifidisve. *Hall. helv. n.* 1151.
A. alpina alba minor. *Bauh. pin.* 176. *prod.* 94.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 283.
Baldiſche Anemone. W.
Habitat alpibus Taureri-Raſtadienſis, Leontinis, Aquilejenſibus, Penninis, Delphinatus, *inque monte* Baldo *et* Ceniſio. ♃ (v. ſ.)
Semina *ſunt lanata, ſtylo brevi glabro coronata.* Caulis *uniflorus et facies plantae fere ut in A. alpina, ſed abunde diſtincta, foliis ſeminibusque.* W.

15. ANEMONE *ſylveſtris.*
A. caule bifloro folioſo, foliis tripartitis, laciniis trifidis dentatis, ſeminibus ſubrotundis lanatis, ſtigmate perſiſtenti. W.
A. pedunculo nudo, ſeminibus ſubrotundis hirſutis muticis. *Sp. pl.* 761. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 176. *Crantz. auſtr. p.* 123. *n.* 5. *Pollich. pal. n.* 515.

Kniph. cent. 5. *n.* 12. *Hoffm. germ.* 190. *Roth. germ. I.* 236. *II.* 605.

A. ſeminibus lanatis, foliis digitatis hirſutis, floribus pentapetalis. *Hall. helv. n.* 1150.

A. ſeminibus hirſutis. *It. oel.* 112. *Fl. ſuec.* 449, 484.

A. ſeminibus ſeta plumoſa terminatis. *Hort. cliff.* 224. *Roy. lugdb.* 224.

A. ſylveſtris alba major. *Bauh. pin.* 176.

A. ſylveſtris 1. *Cluſ. hiſt.* 1. *p.* 244.

β. Anemone ſylveſtris alba minor. *Bauh. pin.* 176. *prod.* 93. *Burſ. IX.* 79. *Pollich. l. c.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 291.

Wald-Anemone. W.

Habitat in Germania, Helvetia, *etc.* ♃ (v. v.)

Variat flore viridi, purpureo-viridi et viridi pleno. Gmel. R.

16. ANEMONE *virginiana.*

A. caule multifloro folioſo, foliis tripartitis, laciniis trifidis acuminatis dentatis, petalis acutis, ſeminibus hirſutis, ſtylo perſiſtenti breviſſimo. W.

A. pedunculis alternis longiſſimis, fructibus cylindricis, ſeminibus hirſutis muticis. *Sp. pl.* 761. *Mill. dict. n.* 4.

A. caule ramoſo, petalis lanceolatis. *Roy. lugdb.* 488. *Gron. virg.* 165.

A. virginiana tertiae ſimilis, flore parvo. *Herm. lugdb.* 645. *parad.* 18. *t.* 18.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 292.

Virginiſche Anemone. W.

Habitat in Virginia. ♃ (v. v.)

Caulis *nudus, terminatus,* Foliis *ternis, ternatis, medio trilobo, acuto, ſerrato, lateralibus bipartitis, lobato-ſerratis.* Pedunculus *exit e centro petiolorum, longus, nudus.* Flos *parvus petalis ovatis, viridibus, inſtar calycis cujusdam.* Piſtillorum *capitulum elongatum, cylindricum, reſolvitur in Semina hirſuta. Juxta pedunculi baſin exit caulis ulterius, terminatus pedunculo florifero et foliis duobus oppoſitis et ſic ulterius.*

17. ANEMONE *decapetala.*

A. caule unifloro, flore decapetalo, foliis ternatis lobatis radicalibus. *Mant.* 79.

A. foliis ternatis: foliolis trilobis punctatis, pedunculo

ſimplici, flore unico decapetalo. *Ard. ſpec.* 2. *p.* 27. *t.* 12.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 294.
Braſilianiſche Anemone. W.
Habitat in Braſilia. ♃
Statura *A. virginianae, ſed minuta.* Folia *radicalia, ternata, petiolata, petiolulata: foliolis ſubovatis, obtuſis trilobis ſ. inciſis.* Caulis *ſimpliciſſimus, ſemipedalis.* Involucrum *uniflorum, tripartitum, multifidum, lineare, acutum, laeve.* Pedunculus *longitudine involucri, hirtus.* Flos *parvus, albus, 10-petalus: petalis lanceolatis, perſiſtentibus: interioribus paullo minoribus.* Stamina *corolla breviora.* Germen *capitatum, conicum, villoſum.* Stigmata *acuminata, laevia.*

*18. ANEMONE *triternata.*
A. foliis ternatis, foliolis cuneatis ſubinciſis, involucris multifidis. *Vahl ſymb.* 3. *p.* 74. *t.* 65.
Dreymaldreyblättrige Anemone. W.
Habitat in Braſilia. ♃
Folia *radicalia petiolata, ternato-ſupradecompoſita glabra;* foliola *parva vix ſemiunguicularia, bi-ſ. trifida obtuſiuſcula;* laciniis *bi-ſ. tridentatis.* Scapus *digitalis altitudine foliorum uniflorus, ſub involucro et floribus incraſſatus villoſoque incanus, inferne glabriuſculus.* Involucrum *foliis ternis baſi connatis bi-ſ. trifidis: laciniis linearibus.* Corolla *dodecapetala.* Petala *lineari-lanceolata, exteriora parum longiora.* Vahl l. c.

19. ANEMONE *penſylvanica.*
A. caule dichotomo, foliis ſeſſilibus amplexicaulibus: infimis ternis trifidis inciſis. *Mant.* 247.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 293.
Penſylvaniſche Anemone. W.
Habitat in Canada. *D. Hope.* ♃.
Simillima A. dichotomae, *ut incautum fallat, ſed differt:* Foliis *ad primam dichotomiam caulis ternis (non binis,) ſeſſilibus, magis venoſis, minus horizontalibus.* Flores *toti albi, (non ſubtus rubicundi,) majores.* Antherae *flavae, (non fulvae.)* Fructus *globoſus, ſeminum apicibus ſubulatis, (non mucrone minuto recurvo) undique exſtantibus.*

20. ANEMONE *dichotoma.*
A. caule dichotomo, foliis ſeſſilibus omnibus oppoſitis

amplexicaulibus trifidis inciſis. *Amoen. acad.* 1. *p.* 155. * *Gmel ſib.* 4. *p.* 197. *Linn. decur.* 29. *t.* 15. *Mill. dict. n.* 7.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 294.
Gabeläſtige Anemone. W.
Habitat in Canada, Sibiria. ♃ (v. ſ.)
Simillima *praecedenti, ſed* Radix *late repens.* Folia *ubique oppoſita, horizontalia.* Corollae *ſubtus purpuraſcententes.* Antherae *fulvae.* Semina *nuda, ſed mucrone minuto recurvo.*

21. ANEMONE *trifolia.*
A. caule unifloro, foliis caulinis ternis ternatis, foliolis ovatis dentatis. W.
A. foliis ternatis ovatis integris ſerratis, caule unifloro. *Hort. upſ.* 155. *Hort. cliff.* 224. *Roy. lugdb.* 488. *Dalib. pariſ.* 161. *Gmel. ſib.* 4. *p.* 199. *n.* 39. *Scop. carn. n.* 659.
A. trifolia, flore albo. *Bauh. hiſt.* 3. *p.* 412.
A. trifolia. *Dod. pempt.* 436.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 294.
Dreyblättrige Anemone. W.
Habitat in Gallia, Carniolia. ♃ (v. ſ.)

22. ANEMONE *quinquefolia.*
A. caule unifloro, foliis caulinis ternis quinatis, foliis ovatis dentatis. W.
A. foliis quinatis ovalibus ſerratis, caule unifloro. *Sp. pl.* 762.
Ranunculus nemorum, fragariae foliis, virginianus. *Pluk. alm.* 310. *t.* 106. *f.* 3.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 295.
Fünfblättrige Anemone. W.
Habitat in Virginia, Canada. *Kalm.* ♃

23. ANEMONE *nemoroſa.*
A. caule unifloro, foliis caulinis ternis ternatis, foliolis lanceolatis trifido-dentatis, corolla hexapetala. W.
A. ſeminibus acutis, foliolis inciſis, caule unifloro. *Hort. cliff.* 224. *Fl. ſuec.* 450, 483. *Mat. med.* 141. *Dalib. pariſ.* 161. *Oed. dan. t.* 549. *Gmel. ſib.* 4. *p.* 198. *Pollich. pal. n.* 519. *Hoffm. germ.* 190. *Kniph. cent* 1. *n.* 2. *Pallas it.* 2. *p.* 14. *Roth. germ. I.* 236. *II.* 606.
A. ſeminibus nudis, caule unifloro, foliis caulinis, quin-

quepartitis, lobis tripartitis inciſis. *Hall. helv. n.* 1154.

A. nemoroſa alba. *Crantz. auſtr. p.* 121.

A. ſeminibus acutis, petalis ellipticis, foliis ſerratis. *Scop. carn. ed.* 1. *p.* 567 *n.* 1. *ed.* 2. *n.* 660.

A. foliis tripartito-multifidis, floribus hexapetalis, defloratis cernuis. *Neck. gallob. p.* 235.

A. nemoroſa, flore majore. *Bauh. pin.* 176.

Ranunculus ſylvarum. *Cluſ. hiſt.* 247.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 295.

Buſch-Anemone. W.

Habitat in Europae *aſperis, duris, nemoribus.* ♃ (v. v.)

24. ANEMONE *apennina.*

A. caule unifloro, foliis caulinis ternis ternatis, foliolis lanceolatis inciſo-dentatis, corolla polypetala. W.

A. ſeminibus acutis, foliolis inciſis, petalis lanceolatis numeroſis. *Sp. pl.* 762. *Hudſ. angl.* 237. *Mill. dict. n.* 3.

Ranunculus nemoroſus, flore caeruleo, apennini montis. *Mentz. pug. t.* 8. *f.* 2.

Ranunculus nemoroſus, flore purpureo-caeruleo. *Park. theatr.* 325. *Raj. angl.* 3. *p.* 259.

A. geranii robertiani folio, caerulea. *Bauh. pin.* 174.

A. hortenſis, tenuifolia, ſimplici flore 1. *Cluſ. hiſt.* 1. *p.* 254.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 296.

Apenninen-Anemone. W.

Habitat in Apenninis, Romae, Anglia. ♃ (v. v.)

*25. ANEMONE *reflexa.*

A. caule ſubbifloro, foliis caulinis ternis ternatis, foliolis ſubtrifidis apice dentatis, petalis lanceolatis obtuſis reflexis. W.

Anemone reflexs. *Stephan. in litt.*

Zurückgebogene Anemone. W.

Habitat in Sibiria. ♃ (v. ſ.)

Simillima ſequenti, qua cum convenit, foliis florum *numero ac* colore, *differt vero:* pedunculis *duplo brevioribus vix petiolo longioribus hirtis,* corollis *reflexis,* petalis *multo anguſtioribus.* W.

26. ANEMONE *ranunculoides.*

A. caule ſubbifloro, foliis caulinis ternis ternatis, foliolis

ſubtrifidis apice dentatis, petalis oblongis obtuſis patentibus. W.

A. ſeminibus acutis, foliolis inciſis, petalis ſubrotundis, caule ſubbifloro. *Hort. cliff.* 195. *Fl. ſuec.* 451, 486. *Roy. lugdb.* 488. *Dalib. pariſ.* 161. *Gmel. ſib.* 4. *p.* 199 *Pollich. pal. n.* 520. *Roth. germ. I.* 236 *II.* 606. *Fl. dan. t.* 146. *Kniph. cent.* 1. *t.* 5. *Hoffm. germ.* 190.

A. biflora, ſeminibus nudis, foliis caulinis quinquepartitis, lobis tripartitis inciſis. *Hall. helv. n.* 1153.

A. nemoroſa lutea. *Crantz. auſtr. p.* 121.

A. ſeminibus acutis; petalis ſubrotundis, foliis crenatis. *Scop. carn.* 7. *p.* 578. *n.* 2. *ed.* 2. *n* 661.

Ranunculus nemoroſus luteus. *Bauh pin.* 178.

Ranunculus phragmites luteus nemoroſus. *Bauh. hiſt.* 3. *p.* 413.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 297.

Ranunkelartige Anemone. W.

Habitat in Europae *borealis ; ratis nemoroſis.* ♃ (v. v.)

27. ANEMONE *narciſſiflora.*

A. floribus umbellatis involucratis, foliis radicalibus palmatis inciſo-dentatis. W.

A. floribus umbellatis, ſeminibus depreſſo-ovalibus nudis. *Sp. pl.* 763. *Gmel. ſib.* 4. *p.* 200. *n.* 42. *Iacq. auſtr. t.* 159. *Crantz. auſtr. t.* 3. *f.* 1. *Mattuſch. ſil. n.* 397. *Hoffm. germ.* 191. *Villars. delph.* 3. *p.* 722. *

A. ſeminibus nudis rotundis planis, foliis ſeptifidis imbricatis, floribus umbellatis. *Hall. helv. n.* 1155.

Ranunculus montanus hirſutus albus altior. *Bauh. pin.* 183. *Hall.*

Ranunculus montanus hirſutus humilior narciſſi flore. *Bauh. pin.* 182.

Ranunculi montani 2, ſpecies altera. *Cluſ. hiſt.* 1. *p.* 235.

Ranunculus alpinus narciſſi flore. *Cluſ. pann.* 366. 367. *Hall.*

Caryophyllata alpina quinquefolia. *Bauh. pin.* 322.

β. A. (*faſciculata*) floribus umbellatis congeſtis, foliis multifidis. *Sp. pl.* 763.

Ranunculus orientalis, aconiti lycoctoni folio, flore magno albo. *Tournef. cor.* 20. *itin.* 3. *p.* 92. *t.* 92.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 298. 299.

Narciſſenblütige Anemone. W.

Habitat in alpibus Galliae, Helvetiae, Auſtriae, Sileſiae, Sibiriae, *inque* Cappadocia. ♃ (v. ſ.)

A. faſciculata eſt ſola hujus ſpeciei varietas, differens tantum pedunculis brevioribus, ut patet ex ſpecimine ſicco et icone Tournefortii. W.

•28. ANEMONE *umbellata.* W.

A. floribus umbellatis involucratis, foliis radicalibus tripartitis, laciniis trifidis integerrimis. W.

A. (*faſciculata*) floribus umbellatis congeſtis, foliis tripartitis trifidis. *Vahl. ſymb. 3. p. 74. (excluſo ſynonymo.)*

Ranunculus orientalis Napelli folio lanuginoſo flore albo. *Tournef. cor.* 20.

Doldenblütige Anemone. W.

Habitat in Cappadocia. ♃ (v. ſ.)

Radix *faſciculata.* Folia *radicalia tri- ſ. quinquepartita, laciniis trifidis acutiuſculis integerrimis, pilis longis albis, praeſertim margine inſtructa.* Scapus *ſemipedalis pilis ſparſis longis obſitus.* Involucrum *e foliis ternis tripartitis integerrimis connatis.* Pedunculi 3 *ſ.* 4 *involucro longiores.* Flores *exacte praecedentis.* W.

29. ANEMONE *thalictroides.*

A. floribus umbellatis, foliis caulinis ſimplicibus verticillatis, radicalibus biternatis. *Sp. pl.* 763. *Mill. dict. n.* 8

Ranunculus nemoroſus, aquilegiae foliis, virginianus aſphodeli radice. *Pluk. alm.* 310. *t.* 106. *f.* 4.

β. Thalictrum caule unifloro, ex eodem puncto foliis quatuor ſimplicibus inſtructo. *Gron. virg.* 62.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 299.

Thalictrumartige Anemone. W.

Habitat in Virginia, Canada ♃

Involucrum *foliolis petiolatis.* Canadenſis *plantae* β, *a D. Kalmio lectae;* Radix *tuberoſa.* Folia *radicalia petiolis longis, duplicato - ternata; foliolis ſubcordatis, angulatis.* Scapus *foliis paulo altior, terminatus foliis quatuor ſimplicibus, foliolis radicalibus ſimilibus, petiolis longis inſidentibus.* Pedunculi *terminales, conferti, ſ. faſciculati* 1, 2, 3, *ſ.* 5, *ſimplices, uniflori, aequales.* Corollae *albae: petalis ovatis, obtuſis, patentibus.* Stamina *flaveſcentia.* Piſtilli *capitulum ovatum muricatum.*

1082. ATRAGENE.

Gen. plant. ed. Schreb. n. 949.

Cal. o. *Corolla* duplex, petalis numeroſis, exterioribus majoribus. *Sem.* caudata.

1. ATRAGENE *alpina.*

A. ſcandens, foliis biternatis ſerratis acuminatis, petalis exterioribus quaternis, interioribus ſpathulatis. W.

A. foliis duplicato-ternatis ſerratis, petalis exterioribus quaternis. *Sp. pl.* 764. *Gmel. ſib.* 4. *p.* 194. *Iacq. auſtr. t.* 241. *Willd. arb.* 30. * *Pall. roſſ.* 2. *p.* 69. *t.* 76. *

A. foliis ternatis, foliolis ternatis, foliolis trifoliis acutis. *Hall. helv. n.* 1145.

Clematis foliis compoſitis et decompoſitis: foliolis ternatis ſerratis. *Mill. ic. t.* 284.

Clematis foliis cordatis crenatis indiviſis trilobis ternatisque. *Roy. lugdb.* 487.

Clematis alpina geranifolia. *Bauh. pin.* 300. *prodr.* 155. *Pluk. alm.* 109 *t.* 84. *f.* 7. *Moriſ. hiſt.* 3. *ſ.* 15. *t.* 2. *f. ult.*

Clematis cruciata alpina. *Pon. bald.* 335.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 300.

Alpen-Atragene. W.

Habitat in Baldi, Auſtriae, Sibiriae *alpibus.* ♄ (v. v.)

Folia *biternata, inciſa, radicalia.* Scapi *uniflori.* Petala 4 *exteriora lanceolata, magna; interiora lanceolata, obovata, parva.*

In Iacquini figura corollae ſunt coeruleae, et pili ariſtae ſeminis ſecundi, in Pallaſii vero: corollae majores albae et pili ariſtae diſtichi. Noſtra in hortis culta convenit in flore cum icone Pallaſii, in ſemine cum Iacquiniana figura. W.

*2. ATRAGENE *ochotenſis.* †

A. ſcandens, foliis biternatis ſerratis acutis, petalis exterioribus ſenis, interioribus linearibus. W.

A. (*ochotenſis*) foliis triternatis, petalis exterioribus ſex, interioribus linearibus. *Pall. roſſ.* 2. *p.* 69. *

Ochozkiſche Atragene. W.

Habitat in Sibiria *verſus mare* Ochotenſe. ♄

*3. ATRAGENE *japonica.*

A. erecta, foliis oppoſitis triternatis, foliolis ovatis inciſis. *Thunb. jap.* 239. *

Japaniſche Atragene. W.

Habitat in Japonia. ♄ ♃

Caulis *erectus ſtriato-angulatus ſubdichotomus villoſus bipedalis.* Folia *petiolata oppoſita triternata patentia* Folio la *ovata acuta inciſa dentata tenuiſſime villoſa, ſubtus pallida, pollicaria.* Petiolus *baſi planus amplexicaulis ſemipollicaris.* Flores *e dichotomiis pauci pedunculati.* Pedunculi *filiformes elongati uniflori.* Petala *plurima ultra viginti ſubaequalia vel interiora paullo breviora ovata obtuſiuſcula, intus purpurea, extus albo-tomentoſa.* Facies *omnino Anemones ſed petala plurima huc referre jubent.* Thunb. l. c.

4. ATRAGENE *capenſis.*

A. erecta, foliis biternatis, foliolis inciſo-dentatis, petalis exterioribus numeroſis. W.

A. foliis ternatis: foliolis inciſis dentatis, petalis exterioribus quinis. *Sp. pl.* 764. *Berg. cap.* 148.

A. foliis ternatis, foliolis inciſis dentatis rigidis, flore incarnato ſemipleno. *Andrews. bot. repoſit. t.* 9. *

Pulſatilla foliis trifidis dentatis, flore incarnato pleno. *Burm. afr.* 148.

Pulſatilla apii folio rigido, flore magno. *Herm. afric.* 18.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 301.

Capiſche Atragene. W.

Habitat ad Cap. b. ſpei. ♄ ♃

Scapus *ſimplex, ſpithamaeus.* Involucrum *in medio Scapi e* ſtipulis *ventricoſis, ovatis, villoſis, foliaceis,* Folia *ternata, cuneiformia, trifida, acuta, dentata, nuda: intermedio longius petiolato.* Petala *circiter* 20, *alba: horum* 6 *inferiora latiora, ſubtus villoſa purpuraſcentia.* Seminum *capitulum ovatum, hirſutum.*

5. ATRAGENE *tenuifolia.*

A erecta, foliis ternato-ſupradecompoſitis, foliolis lineari-filiformibus. *Thunb. jap.* 239. *in nota.*

A. foliis duplicato-pinnatis, pinnulis linearibus integris. *Lin. ſuppl.* 270.

Feinblättrige Atragene. W.

Habitat ad Cap. b. Spei.

6. ATRAGENE *zeylanica.*
A. ſcandens, foliis conjugatis indiviſis. *Thunb. jap.* 239. *in nota.*
A. cirrhis diphyllis, petalis exterioribus quaternis. *Fl. zeyl.* 226. *
Atragene. *Amoen. acad.* 1. *p.* 105.
Clematis zeylanica, floribus obſoletis parvis. *Herm. zeyl.* 55.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 302.
Zeylaniſche Atragene. W.
Habitat in Zeylona. ♄
Caule/cens et ſcandens.

1083. CLEMATIS.

Gen. plant. edit. Schreb. n. 960.

Cal. 0. *Petala* 4, rarius 5. *Sem.* caudata.

* *Scandentes.*

1. CLEMATIS *cirrhoſa.*
C. foliis ſimplicibus, caule cirrhis oppoſitis ſcandente, pedunculis unifloris lateralibus. *Syſt. veg.* 426. *Kniph. cent.* 3. *n.* 28.
C. cirrhis ſcandens, foliis ſimplicibus. *Sp. pl.* 2. *p.* 766. *Hort. cliff.* 226. *Roy. lugdb.* 487. *Mill. dict. n.* 7.
C. peregrina, foliis pyri inciſis. *Bauh. pin.* 300.
C. altera baetica. *Cluſ. hiſt.* 1. *p.* 123.
C. peregrina, foliis pyri inciſis: nunc ſingularibus, nunc ternis. *Tournef. cor.* 20.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 303.
Einfachblättrige Waldrebe. W.
Habitat in Baetica, *arbores operiens et deprimens.* ♄ (v. v.)
Cirrhi oppoſiti, a foliis diſtincti. Folia *ſaepius ſimplicia, ſerrata, petiolata, ovata, ex iisdem geniculis caulis plura.* Pedunculi *axillares, uniflori; ſub ſingulo flore calyculus remotus, hemiſphaericus, minutus.*

*2. CLEMATIS *florida.*
C. foliis decompoſitis, foliolis binatis ternatisque, petalis ovatis. *Thunb. jap.* 240. *Ait. Kew.* 2. *p.* 258.
Anemone vel Anemonoides. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 280. *t.* 55. *f.* 1.

Groſsblütige Waldrebe. W.
Habitat in Japonia. ♄ (v. v.)

Caulis *ſcandens ſtriatus purpureus omni parte glaber.* Folia *oppoſita binata vel ternata; pinnae ſeſſiles ovatae acutae integrae, rarius inciſae, villoſae, media majori, unguiculares.* Petioli *baſi perfoliati ſemiteretes horizontaliter patentes digitales.* Flores *ex axillis pedunculati ſolitarii.* Pedunculus *villoſus uniflorus foliis longior.* Corolla *magna ſpecioſa patens, petalis ovatis acuminatis luteſcentibus.* Stamina *lineari-lanceolata purpuraſcentia inaequalia corolla dimidio breviora. Differt a C. Viticella:* foliis *ſemper decompoſitis*, foliolis *acutis*, petalis *ovatis.* Thunb. l. c.

3. CLEMATIS *Viticella.*

C. foliis compoſitis decompoſitisque, foliolis ſublobatis integerrimis, petalis obovatis patentibus. W.

C. foliis compoſitis decompoſitisque: foliolis ovatis ſublobatis integerrimis. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Mill. dict. n.* 8. *Scop. carn. n.* 670. *Kniph. cent.* 11. *n.* 30. *Willd. arb.* 64. *

Clematitis caerulea vel purpurea repens. *Bauh. pin.* 300.
Clematitis altera. *Cluſ. hiſt.* 1. *p.* 122. *Cam. epit.* 696.
Clematitis caerulea, flore pleno. *Bauh. pin.* 301.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 304.
Italiäniſche Waldrebe. W.
Habitat in Italiae, Hiſpaniae, *ſepibus.* ♄ (v. v.)

Folia *floralia ovata.* Petala *deltoidea, ſ. ſuperne dilatata.*

4. CLEMATIS *Viorna.*

C. foliis compoſitis decompoſitisque, foliolis ovatis trifidisque integerrimis, corollis campanulatis, petalis acuminatis. W.

C. foliis compoſitis decompoſitisque: foliolis quibusdam trifidis. *Sp. pl.* 765. *Gron. virg.* 62. *Mill. dict. n.* 10.

Flammula ſcandens, flore violaceo clauſo. *Dill. elth.* 144. *t.* 118 *f.* 144.

Clematis purpurea repens, petalis florum coriaceis. *Raj. hiſt.* 1928.

Scandens caroliniana planta, viornae folio. *Pet. ſicc.* 27.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 305.
Glockenblütige Waldrebe. W.
Habitat in Virginia, Carolina. ♄ (v. ſ.)

5. CLEMATIS *criſpa.*

C. foliis pinnatis ternatisque, foliolis ovato-lanceolatis tri-lobisve integerrimis, corollis campanulatis acuminatis margine undulatis, ariſtis ſeminum ſubulatis nudis. W.

C. foliis ſimplicibus ternatisque; foliolis integris trilobisve. *Sp. pl.* 765. *Mill. dict. n.* 15. *Willd. arb.* 65. *

C. flore criſpo. *Dill. elth.* 86. *t.* 73. *f.* 84.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 306.

Krausblütige Waldrebe. W.

Habitat in Carolina. ♄ (v. v.)

Seminum *ariſtae craſſae ſubulatae non plumoſae, ut in reliquis, ſed pubeſcentes pilis brevibus adpreſſis.* W.

*6. CLEMATIS *calycina.*

C. involucro calycino approximato, foliis ternatis, intermedio tripartito. *Ait. Kew.* 2. *p.* 259. *Vahl ſymb.* 2. *p.* 75. *

C. *(balearia)* ſcandens, foliis compoſitis tenuiter laciniatis, floribus calyculatis lateralibus, petalis interne guttatis. *Lamarck. encycl.* 2. *p.* 44. *

Gekelchte Waldrebe. W.

Habitat in Minorca ♄ (v. v.)

Folia *ternata, foliolis profunde inciſis, intermedio tripartito.* Pedunculi *uniflori axillares nutantes.* Involucrum *calyciforme ſub corolla, monophyllum bilobum.* Petala *quatuor oblonga externe pubeſcentia, interne pallide flava, ſtriis nonnullis rubicundis notata.* W.

7. CLEMATIS *orientalis.*

C. foliis compoſitis, foliolis cuneiformibus trilobis ſubdentatis acuminatis, petalis interne villoſis. W.

C. foliis compoſitis: foliolis inciſis angulatis lobatis cuneiformibus, petalis interne villoſis. *Sp. pl.* 765. *Mill. dict. n.* 11.

C. orientalis, apii folio, flore e viridi flaveſcente poſterius reflexo. *Tournef. cor.* 20.

Flammula ſcandens, apii folio glauco. *Dill. elth.* 144. *t.* 119. *f.* 145.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 306.

Orientaliſche Waldrebe. W.

Habitat in Oriente. ♄ (v. f.)

Flores cernui: petalis recurvatis, ſupra villoſis.

*8. CLEMATIS *glauca.* W.

C. foliis compoſitis, foliolis ovatis ſublobatis obtuſis mucronatis, petalis glabris margine pubeſcentibus. W.

C. foliis compoſitis glaucis, foliolis ovatis ſublobatis, petalis margine tomentoſis. *Willd. arb.* 65. *t.* 4. *f.* 1. *

Meergrüne Waldrebe. W.

Habitat in Sibiria, Oriente. ♄ (v. v.)

Diverſiſſima a praecedente, foliolis *non trilobis acuminatis*, petalis *interne glabris ut et toto habitu.* W.

9. CLEMATIS *hexapetala.*

C. foliis compoſitis, foliolis ovatis ſerratis, pedunculis diphyllis, corolla patente hexapetala. *Lin. ſuppl.* 271.

C. foliis ternatis, foliolis cordato-ovatis crenato-inciſis, pedunculis ramoſis trichotomis diphyllis, floribus polypetalis dioicis. *Forſt. prod. n.* 230.

Sechsblättrige Waldrebe. W.

Habitat in Nova Zeelandia. ♄

Scandens, *ſimillima reliquis.* Flores *luteſcentes.*

*10. CLEMATIS *triflora.*

C. foliis ternatis, foliolis ovatis ſerratis mucronatis, pedunculis trifloris. *Vahl ſymb.* 3. *p.* 74. *

C. (*mauritiana*) foliis ternatis, foliolis ſubcordatis ſerratis, ſeminum cauda longiſſima plumoſa. *Lamarck. encycl.* 2. *p.* 43. *

Dreyblumige Waldrebe. W.

Habitat in inſula Borboniae. ♄

Corolla *tetrapetala magnitudine C. virginianae.* Petala *oblonga utrinque villoſa.* Semina *purpurea villoſa; cauda longa plumoſa.* Vahl l. c.

11. CLEMATIS *virginiana.*

C. foliis ternatis: foliolis cordatis ſublobato-angulatis ſcandentibus, floribus dioicis. *Amoen. acad.* 4. *p.* 274. * *Willd. arb.* 66. *

C. virginiana pannonicae ſimilis. *Pluk. mant.* 51. *t.* 389. *f.* 4.

C. floridenſis, flore albo odoratiſſimo. *Albin. anat.* 1. *p.* 79. *t.* 7.

C. aquatica trifoliata late ſcandens, floribus albis odoratis. *Gron. virg.* 62. *n.* 270.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 307.

Virginiſche Waldrebe. W.

Habitat in America *ſeptentrionali.* ♄ (v. v.)

Femina gaudet ſtaminibus, ſed antherae polline deſtituuntur.

*12. CLEMATIS *japonica.*

C. foliis ternatis, foliolis elliptico-ovatis ſerratis, floribus cylindricis. *Thunb. jap.* 240.

Japaniſche Waldrebe. W.

Habitat in Japonia. ♄

Caulis *filiformis ſcandens ſtriatus purpureus villoſus.* Folia *e geniculis plura petiolata ternata.* Foliola *breviter petiolata elliptico-oblonga acuminata a medio ad apicem aequaliter ſerrata nervoſa villoſa pollicaria, intermedio majori.* Petiolus *capillaris laxus palmaris.* Flores *e geniculis pedunculati ſolitarii cylindrici purpurei.* Pedunculi *uniflori longitudine foliolorum.* *Differt a C. virginica:* foliolis *utrinque attenuatis nec cordatis,* floribus *cylindricis nec ovatis.* Thunb. l. c.

*13. CLEMATIS *trifoliata.*

C. foliis oppoſitis ternatis glabris, foliolis ovatis repando-dentatis, caule ſcandente. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 337.

Scandens foliis ternatis. *Thunb. jap. p.* 357. *n.* 43.

Dreyblättrige Waldrebe. W.

Habitat in Japonia. ♄

14. CLEMATIS *dioica.*

C. foliis ternatis integerrimis, floribus dioicis. *Amoen. acad.* 6. *p.* 398. * *Mill. dict. n.* 13.

C. ſcandens, foliis quinquenerviis ovatis nitidis pinnato-ternatis. *Brown. jam.* 255.

C. prima ſ. ſylveſtris latifolia, foliis ternis. *Sloan. jam.* 84. *hiſt.* 1. *p.* 199. *t.* 128. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 308.

Weſtindiſche Waldrebe. W.

Habitat in America *calidiore.* ♄

*15. CLEMATIS *indiviſa.* W. †

C. foliis ternatis, foliolis ovatis integerrimis mucronatis, pedunculis axillaribus paniculatis diphyllis.

Clematis integrifolia. *Forſt. prod. n.* 231.

Ungetheilte Waldrebe. W.

Habitat in Nova Zeelandia. ♄

*16. CLEMATIS *paniculata.*

C. foliis quinato-pinnatis, foliolis cordatis ovatis integris. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 337.

Clematis criſpa. *Thunb. jap.* 239.
Clematis Vitalba japonica. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 309. *t.* 55. *f.* 2?
Riſpenblütige Waldrebe. W.
Habitat in Iaponia. ♄

Caulis *flexuoſus ſcandens ſtriatus glaber.* Rami *alterni.* Folia *petiolata pinnato-quinata.* Foliola *petiolata cordata ovata acuta indiviſa integerrima glabra inferiora majora, ſuperiora minora.* Petioli *flexuoſi.* Flores *axillares albi.* Pedunculi *triternato-paniculati filiformes.* Thunb. Act. l. c.

17. CLEMATIS *Vitalba.*

C. foliis pinnatis, foliolis ſcandentibus cordatis inciſo-lobatis. W.
C. foliis pinnatis: foliis cordatis ſcandentibus. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Gron. virg.* 62. *Sauv. monſp.* 249. *Gort. gelr.* 315. *Mill. dict. n.* 4. *Iacq. auſtr. t.* 308. *Scop. carn. n.* 669. *Pollich. pal. n.* 521. *Hoffm. germ.* 191. *Roth. germ. I.* 232. *II.* 595. *Willd. arb.* 67. *
C. caule ſcandente, foliis pinnatis, ovato-lanceolatis. *Hall. helv. n.* 1142.
C. ſcandens, foliis longe petiolatis pinnatis aut dentatis. *Crantz. auſtr. p.* 126. *n.* 1.
Clematis latifolia integra. *Bauh. hiſt.* 2. *p.* 125.
Clematis tertia. *Cam. epit.* 697.
β. Clematis latifolia dentata. *Bauh. hiſt.* 2. *p.* 125.
Vitalba. *Dod. pemp.* 404.
γ. Clematis ſylveſtris latifolia. *Bauh. pin.* 300.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 308.
Gemeine Waldrebe. W.
Habitat in ſepibus Europae *auſtralis.* ♄ (v. ſ.)

Petala *coriacea.* Variat *foliolis integris et inciſis.*

*18. CLEMATIS *chinenſis.*

C. foliis quinato-pinnatis, foliolis lanceolatis. *Retz. obſ.* 2. *p.* 18. *t.* 2.
Chinenſiſche Waldrebe. W.
Habitat in China. ♃ ♄ (v. ſ.)

Caules *debiles ſuſtentandi tetragoni vix ſcandentes.* Folia *pinnata, foliolis quinis, paribus longe remotis petiolatis lanceolatis baſi ovatis.* Pedunculi *oppoſiti axillares tri-vel quinqueflori, pedicellis lateralibus bracteolis binis a flore longe remotis inſtructi, termi-*

nali nudo. Flores *parvi tetrapetali, petalis lineari-lanceolatis obſcure purpureis, margine interiore linea tomentoſa pictis.* Pappus *piſtillorum antheris vix brevior.* Retz. l. c.

Valde affinis C. paniculatae, an forte varietas? W.

19. CLEMATIS *Flammula.*

C. foliis inferioribus pinnatis laciniatis: ſummis ſimplicibus integerrimis lanceolatis. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Sauv. monſp.* 249. *Mill. dict. n.* 6. *Knorr. del* 1. *t. C.* 9. *Willd. arb.* 68. *

C. caule ſcandente, foliis pinnatis trilobatis. *Hall. helv. n.* 1143.

C. foliis pinnatis: foliolis alternis; caulinis flexuoſis ſcandentibus. *Ger. prov.* 382.

Clematis ſ. Flammula ſcandens tenuifolia alba. *Bauh. hiſt.* 2. *p.* 127. *Hall.* R.

C. urens 1. 2. *Tabern.* 832. 833. R.

Clematis ſ. Flammula repens. *Bauh. pin.* 300. *Raj. hiſt.* 621.

Flammula. *Dod. pempt.* 403. *Dalech. hiſt.* 1171.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 310.

Scharfe Waldrebe. W.

Habitat in Monſpelii, *inque* Rhetiae *ſepibus.* ♃ ♄ (v. v.)

** *Erectae.*

20. CLEMATIS *maritima.*

C. foliis pinnatis linearibus, caulibus ſimplicibus hexagonis. *Sp. pl.* 767.

C. maritima repens. *Bauh. pin.* 300. *prodr.* 135. *Allion. nic.* 122. *

Houttuyn Lin. Pfl. Syſt. 7. *p.* 310.

Meerſtrands-Waldrebe. W.

Habitat ad litora maris Adriatici, Venetii, Monſpelii. ♃

Planta *perennis.* Caules *e ſarmentis procumbentibus, plures, erectiuſculi, erecti.* Folia *oppoſita, pinnata: foliolis ſaepius 5 paribus, remotis, linearibus, rigidiuſculis, ſubſeſſilibus, laevibus, indiviſis ſ. rarius bi- vel trifidis.* Flores *C. rectae.* Varietatem *C. Flammulae ſtatuunt Magnol. et Rajus; meam potius C. rectae judicarem ex ſolo mutatam?*

*21. CLEMATIS *anguſtifolia.*

D. foliis pinnatis, foliolis lanceolatis obtuſis, inferiori-

bus tripartitis, caulibus ſimplicibus ſtriatis erectis, corollis polypetalis. W.

C. (*anguſtifolia*) erecta, foliis pinnatis recurvis, foliolis lineari-lanceolatis, floribus polypetalis. *Iacq. ic rar.* 1. *t.* 104. *Collect.* 1. *p.* 137. *

Clematis hexapetala. *Pall. itin.* 3. *app. n.* 96. *t. Q. f.* 2.

C. erecta foliis anguſtis, cornu cervi diviſura. *Amman. ruth. n.* 108.

Atragene foliis pinnatis, foliolis ex lineari-lanceolatis ſimplicibus bi-et trifidis. *Gmel. ſib.* 4. *p.* 194.

Schmalblättrige Waldrebe. W.

Habitat in Sibiria *et* Auſtria *littorali.* ♃

Folia *pinnata, foliolis anguſto-lanceolatis obtuſis.* Foliolorum *par baſi proximum, ut et foliola in foliis inferioribus ſunt profunde tripartita.* Corolla *alba hexa-vel octopetala.* W.

22. CLEMATIS *erecta.*

C. foliis pinnatis: foliolis ovato-lanceolatis integerrimis, caule erecto, floribus pentapetalis tetrapetaliſque. *Hort. cliff.* 225. *Hort. upſ.* 155. *Roy. lugdb.* 486. *Sauv. monſp.* 249. *Mill. dict. n.* 1. *Iacq. auſtr. t.* 291. *Gmel. it.* 1. *p.* 125. *Scop. carn. n.* 667. *Hoffm. germ.* 191. *Roth. germ. I.* 232. *II.* 596.

C. caule erecto, foliis pinnatis, ovato-lanceolatis. *Hall. helv. n.* 1144.

C. foliis longe petiolatis, integris, caule erecto. *Crantz. auſtr. p.* 126.

Flammula recta. *Bauh. pin.* 300.

Flammula. *Cam. epit.* 698. *Cluſ. hiſt. p.* 124.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 311.

Aufrechte Waldrebe. W.

Habitat in collibus Auſtriae, Pannoniae, Tatariae, Monſpelii, Helvetiae. ♃ (v. v.)

Umbellae erectae et ſtrictae ſunt.

*23. CLEMATIS *ochroleuca.*

C. foliis ſimplicibus ovatis pubeſcentibus integerrimis, floribus erectis. *Ait. Kew.* 2. *p.* 260.

C. erecta humilis non ramoſa, foliis ſubrotundis, flore unico ochroleuco. *Pluk. mant.* 51. *t.* 379. *f.* 5.

Gelbliche Waldrebe. W.

Habitat in America *boreali.* ♃ (v. v.)

24. CLEMATIS *integrifolia.*

C. foliis ſimplicibus ſeſſilibus ovato-lanceolatis, floribus

cernuis. *Hort. cliff.* 225. *Hort. upſ.* 156. *Roy. lugdb.* 486. *Mill. dict. n.* 2. *Gmel. ſib.* 4. *p.* 194. *Jacq. auſtr. t.* 363. *Willich. in nov. Act. A. N. C. IV. p.* 107. *Kniph. cent.* 1. *n.* 15. *Hoffm. germ.* 191.

C. (*nutans*) foliis ſeſſilibus integris, caule erecto. *Crantz. auſtr. p.* 124.

C. inclinata. *Scop carn. n.* 668.

Clematis caerulea erecta. *Bauh. pin.* 300.

Clematis caerulea pannonica. *Cluſ. hiſt.* 1. *p.* 123.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 312.

Einfache Waldrebe. W.

Habitat in Hungaria, Tataria. ♃ (v. v.)

Petala *coriacea.*

1084. THALICTRUM.

Gen. pl. ed. Schreb. n. 951.

Cal. o. *Petala* 4. ſ. 5. *Sem.* ecaudata.

1. THALICTRUM *alpinum.*

T. caule ſimpliciſſimo ſubnudo, racemo ſimplici terminali. *Hort. cliff.* 227. *Fl. ſuec.* 457, 491. *Roy. lugdb.* 485. *Mill. dict. n.* 11. *Gunn. norv. n.* 41. *Fl. dan. t.* 11.

T. caule ſubnudo ſimplici, foliolis obtuſis. *Fl. lapp.* 225. *

T. montanum minimum praecox, foliis ſplendentibus. *Moriſ. hiſt.* 3. *p.* 325. *ſ.* 9. *t.* 20. *f.* 14.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 314.

Alpen-Thalictrum. W.

Habitat in alpibus Lapponiae, Arvoniae. ♃ (v. ſ.)

Petala 4, *Stamina* 12, *Piſtilla* 8, *Racemus ſimplex floribus cernuis.*

2. THALICTRUM *foetidum.*

T. caule paniculato filiformi ramoſiſſimo folioſo. *Sp. pl.* 768. *Mill. dict. n.* 8.

T. caule filiformi ramoſiſſimo in paniculam diſperſo ſubjectis foliolis. *Hort. cliff.* 227. * *Roy. lugdb.* 485. *Sauv. monſp.* 251.

Thalictrum ſaxatile. *Villars. delph.* 3. *p.* 714.

T. foliis patulis pubeſcentibus, trilobatis, floribus nutantibus, petalis ſubhirſutis. *Hall. helv. n.* 1140.

T. minimum foetidiſſimum. *Bauh. pin.* 337. *prodr.*

147. *Bauh. hiſt.* 3. *p.* 488. *Burſ.* XIX: 49. *Moriſ. hiſt.* 3. *p.* 325. *ſ.* 9. *t.* 20. *f.* 13. *Pluk. alm.* 367. *t.* 65. *f.* 4.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 314.
Stinkendes Thalictrum. W.
Habitat Monſpelii, *inque* Valleſia, Helvetia. ♃ (v. v.) Odor *hircoſus*.

3. THALICTRUM *tuberoſum*.
T. floribus pentapetalis, radice tuberoſa. *Sp. pl.* 768. *Hort. cliff.* 226. *Roy. lugdb.* 486. *Mill. dict. n.* 6. *et ic. t.* 265. *f.* 2.
T. minus, aſphodeli radice, magno flore. *Tournef. inſt.* 271.
T. minus, grumoſa radice, floribus majoribus. *Herm. flor.* 9. *Boerh. lugdb.* 1. *p.* 45.
Ranunculus thalictri folio, aſphodeli radice. *Moriſ. hiſt.* 2. *p.* 438. *ſ.* 4. *t.* 28. *f.* 13.
Oenanthe hederae foliis. *Bauh. pin.* 163.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 315.
Knolliges Thalictrum. W.
Habitat in Hiſpania, Pyrenaeis. ♃ (v. v.)

4. THALICTRUM *Cornuti*.
T. floribus dioicis, foliolis ovatis trifidis, paniculis terminalibus. *Ait. Kew.* 2. *p.* 262.
T. floribus pentapetalis, radice fibroſa. *Sp. pl.* 768. *Roy. lugdb.* 436.
T. canadenſe. *Corn. canad.* 168.
T. americanum. *Park. theatr.* 265.
T. majus, foliis aquilegiae, flore albo. *Moriſ. hiſt.* 3. *p.* 325. *ſ.* 9. *t.* 20. *f.* 15.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 316.
Cornutiſches Thalittrum. W.
Habitat in Canada. ♃ (v. ſ.)

5. THALICTRUM *dioicum*.
T. floribus dioicis, foliis ſubrotundis cordatis lobatis, lobis obtuſis, pedunculis axillaribus folio brevioribus *Ait. Kew.* 2. *p.* 262.
T. floribus dioicis. *Sp. pl.* 768. *Mill. dict. n.* 9.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 316.
Getrenntes Thalictrum. W.
Habitat in Canada. *Kalm.* ♃ (v. ſ.)
Altitudo vix pedalis. Caulis *rectus.* Flores *tetra-*

petali corollis et ſtaminibus 40 *albis.* Antherae *flaveſcentes.* Foliola *cernua ſunt, qua nota diſtinctiſſima evadit planta ex habitu.* Feminae *ſtyli ſetacei, longiores quam in congeneribus.*

*6. THALICTRUM *elatum.*

T. foliolis ovatis ſubcordatis ſubtrifidis, panicula terminali, floribus erectis, caule teretiuſculo. W.

T. (*elatum*) foliis multipartitis, foliolis cordatis vel ovatis obtuſis inciſis, caule tereti paniculato. *Murr. ſyſt. veg. ed.* 14. *p.* 513. *Iacq. hort.* 3. *t.* 95. *

Hohes Thalictrum. W.

Habitat in Hungaria. ♃ (v. v.)

Habitu accedit ad T. minus, differt vero: ſtatura *majori*, foliolis *latioribus minus profunde inciſis*, floribus *erectis non cernuis.* W.

*7. THALICTRUM *majus.*

T. foliolis ſubrotundis ſubcordatis trifidis ſubtus glaucis, panicula folioſa, floribus cernuis. W.

T. (*majus*) foliis multipartitis, floribus cernuis, ramis floriferis ex iisdem axillis binis vel ternis. *Murr. ſyſt. veg. ed.* 14. *p.* 513. *Iacq. auſtr.* 5. *t.* 420. *Hoſt. ſyn.* 306.

Groſses Thalictrum. W.

Habitat in Auſtriae, Pannoniae *ſylvaticis.* ♃ (v. v.)

Simile T. minori differt tamen: foliolis *ſubtus glaucis majoribus, ut magnitudine totius plantae, denique* panicula *folioſa nec aphylla.* W.

*8. THALICTRUM *medium.*

T. foliolis oblongis cuneiformibus acutis trifidis, ſummis indiviſis lanceolatis, floribus erectiuſculis. W.

T. (*medium*) foliis multipartitis, foliolis lanceolatis inciſis, floribus erectis. *Murr. ſyſt. veg. ed.* 14. *p.* 513. *Iacq. hort.* 3. *t.* 96.

Mittleres Thalictrum. W.

Habitat in collibus Hungariae. ♃ (v. v.)

A T. minori diverſum: foliolis *cuneiformibus ſummis indiviſis lanceolatis et* floribus *erectiuſculis nunquam revera cernuis.* W.

9. THALICTRUM *minus.*

T. foliolis ſubrotundis trifidis inciſis, floribus cernuis. W.

T. foliis ſexpartitis, floribus cernuis. *Sp. pl.* 769. *Mill. dict. n.* 7. *Scop. carn. n.* 674. *Pollich. pal. n.* 522.

Fl. dan. t. 732. *Iacq. auſtr.* 5. *p.* 9. *t.* 419. *Hoffm. germ.* 192. *Roth. germ. I.* 233. *II.* 597.
T. foliis patulis, imis obtuſe trilobis, ſupremis acutis, panicula diffuſa, floribus nutantibus. *Hall. helv. n.* 1139.
T. alpinum minus ſaxatile, rutae folio, ſtaminibus luteis. *Segu. ver.* 476. *t.* 11. *Hall.* R.
T. caule folioſo, foliolis caulinis acutis, panicula divaricata, floribus nutantibus. *Fl. ſuec.* 454, 489. *It. gotl.* 180. * *Dalib. pariſ.* 454.
T. caule glabro, panicula reclinata, floribus cernuis. *Crantz. auſtr. p.* 107. *n.* 4. R.
T. ſiliqua ſeminis ſtriata, minus. *Moriſ. umb. t.* 8. R.
T. minus. *Bauh. pin.* 335. *Dod. pempt.* 58.
Ruta pratenſis minor. *Tabern. ic.* 55.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 317.
Kleines Thalictrum. W.
Habitat in Europae *pratis.* ♃ (v. v.)

Dignoſcitur apicibus foliorum purpuraſcentibus et Caulis nebula coeruleſcente. Floret *praecocius cum T. aquilegifolio, alpino, purpuraſcente, anguſtifolio et contorto.*

*10. THALICTRUM *rugoſum.* †
T. caule ſtriato, foliis rugoſis venoſis, lobulis obtuſis. *Ait. Kew.* 2. *p.* 262.
Runzliches Thalictrum. W.
Habitat in America *boreali.* ♃

11. THALICTRUM *ſibiricum.*
T. foliis tripartitis: foliolis ſubreflexis argute incifis, floribus cernuis. *Sp. pl.* 769.
T. alpinum minus ſaxatile, rutae folio, ſtaminibus luteis. *Segu. ver.* 1. *p.* 476. *t.* 11?
T. orientale minimum fumariae folio. *Tournef. cor.* 20.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 318.
Sibiriſches Thalictrum. W.
Habitat in Sibiria, Armenia. ♃ (v. ſ.)

Habitus Th. minoris ſ. purpuraſcentis, at Foliola *parva, ſextuplo minora, glauca colore Rutae ſ. Fumariae.* Caulis *viridis, maxime paniculatus.* Panicula *ad divaricationes fuſca.* Floret *ſerius, nempe cum T. flavo et dioico.*

Planta ſpontanea eſt duplo minor quam T. minus, hortenſis vero multo major evadit et habitu ad T. medium

accedit, differt vero: foliolis *omnibus tripartis,* floribus*que cernuis.* W.

*12. THALICTRUM *ſquarroſum.*

T. foliolis trifidis indiviſisque, petiolis amplexicaulibus membranaceis alatis, floribus cernuis. W.

Thalictrum ſquarroſum. *Stephan. in litt.*

Geöhrtes Thalictrum. W.

Habitat in Sibiria. ♃ (v. ſ.)

Diverſum ab antecedentibus, quibus floribus nutantibus ſimile, petioli *ſtructura.* Folia *ſupradecompoſita ut in omnibus, foliola lateralia plerumque ovata acuta integerrima, terminalia vero et lateralia ſuperiora trifida vel etiam bifida.* Petiolus *baſi valde dilatatus, alis orbiculatis membranaceis margine dentatis. In petiolis partialibus foliorum ſunt et* auriculae *orbiculatae minores dentatae, qua nota facile dignoſcitur.* W.

13. THALICTRUM *purpuraſcens.*

T. foliis tripartitis, caule foliis duplo altiore, floribus cernuis. *Sp. pl.* 769.

T. virginianum elatius glaucum, ſtaminibus purpuraſcentibus. *Moriſ. hiſt.* 3. *p.* 324.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 319.

Röthliches Thalictrum. W.

Habitat in Canada. ♃

Habitus Thalictri minoris, at dimidio humilius evadit, ubi illud tripedale. Calyces *purpuraſcentes, ſtaminibus luteis.* Caulis *purpuraſcens, abſque nebula. Differt a T. minori foliolis minime apice purpuraſcentibus. Tempore florendi praecocius cum T. minori, nec cum T. flavo. Flos ut in praecedenti: Petalis* 4, *Staminibus* 14, *Piſtillis* 8.

14. THALICTRUM *anguſtifolium.*

T. foliolis lanceolato-linearibus integerrimis. *Hort. cliff.* 226. *Amoen. acad.* 3. *p.* 48. *Roy. lugdb.* 478. *Sauv. monſp.* 250. *Mill. dict. n.* 10. *Pollich. pal. n.* 523. *Kniph. cent.* 5. *n.* 86 *Scop. carn. n.* 676. *Hoffm. germ.* 192. *Roth. germ. I.* 233. *II.* 597.

T. foliis adſcendentibus linearibus; panicula erecta. *Hall. helv. n.* 1137.

T. (*Bauhini*) foliis anguſtiſſimis aeque latis integerrimis. *Crantz auſtr. p.* 105. *n.* 1. *ſec. Hall.* R.

T. (*anguſtiſſimum*) foliis anguſtis lanceolatis, ultimo ramo dentato. *Crantz. l. c. n.* 2. *Hall.* R.

T. pratenſe, anguſtiſſimo folio. *Bauh. pin.* 337. *prodr.* 146. *t.* 146. *Pluk. alm.* 364. *t.* 65. *f.* 6.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 319.
Schmalblättriges Thalictrum. W.
Habitat in Germania *rarius.* ♃ (v. v.)
Petala 4, *ſtamina* 16, *Piſtilla* 7. Affine *T. flavo. Eſt forte ſola T. flavi varietas, vidi enim ſaepius inter caules plantae cultae alios foliis latioribus ad T. flavum accedentibus.* W.

15. THALICTRUM *flavum.*
T. caule folioſo ſulcato, panicula multiplici erecta. *Hort. cliff.* 226. *Fl. ſuec.* 453, 488. *Roy. lugdb.* 485. *Dalib. pariſ.* 162. *Mill. dict. n.* 1. *Gmel. it.* 1. *p.* 139. *Scop. carn.* 2. *n.* 673. *Pollich. pal. n.* 5-4. *Kniph. cent.* 5. *n.* 87. *Hoffm. germ.* 192. *Roth. germ.* *I.* 233. *II.* 598.
T. pratenſe. *Fl. lapp.* 224.
T. foliis erectis, lobis acute trilobis, ſpica adſcendente. *Hall. helv. n.* 1138.
T. foliis triplicato - pinnatis: fructificatione paniculata erecta. *Neck. gallob. p.* 236.
T. nigrius, caule et ſemine ſtriato. *Bauh hiſt.* 3. *p.* 487. *Hall.* R.
T. majus, ſiliqua angulofa ſ. ſtriata. *Bauh. pin.* 336.
T. magnum. *Dod. pempt.* 58.
β. Thalictrum, *flavum, ſpecioſum.*
T. caule anguloſo, foliis linearibus bifidis, trifidisque, panicula multiplici erecta. *Mill. dict. n.* 2. *Doerr. naſſ. p.* 231.
T. majus flavum, floribus luteis ſ. glauco folio. *Bauh. pin.* 336. *ſtaminibus fulvis.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 319.
Gemeines Thalictrum. W.
Habitat in Europae *ſeptentrionalioris ſubhumidis*, β. *in* Hiſpania. ♃ (v. v.)
Petala 4, *Stamina* 24, *Piſtilla* 10 — 16.
Thalictrum *majus flavum Bauh. potius ad* Th. aquilegifolium *ſpectare. Scop. l. c.* R.

*16. THALICTRUM *nigricans.*
T. caule folioſo ſulcato, floribus corymboſis, foliolis radicalibus cuneiformibus trifidis, caulinis linearibus. W.
T. (*nigricans*) caule folioſo ſulcato, panicula multiplici

erecta, foliolis acutis trilobis. *Murr. ſyſt. veg. ed.* 14. *p.* 513. *Iacq. auſt.* 5. *t.* 421.
Schwärzliches Thalictrum. W.
Habitat in Auſtria. ♃ (v. v.)
Simile praecedenti, tamen foliis caulinis diverſum! W.

17. THALICTRUM *ſimplex.*
T. caule folioſo ſimpliciſſimo angulato. *Mant.* 78. *Fl. ſuec.* 2. *p.* 490. *Fl. dan.* 244.
Thalictrum anguſtifolium. *Villars delph.* 3. *p.* 712. * (*excluſis ſynonymis.*)
Houttuyn Lin. Pfl. Syſt. 7. *p.* 321.
Einfachſtengliges Thalictrum. W.
Habitat in Suecia, Dania, Gallia. ♃ (v. ſ.)
Differt a T. flavo: Herba *dimidio minore*, Floribus *nutantibus, nec erectis*; Petalis *viridibus, nec albis*; Filamentis 14, *purpuraſcentibus, nec* 17 *albis*; Panicula *rariore*; Foliis *anguſtioribus.* Praeter *ſtipulas inter ramificationem primam foliorum,* ſquamae *nullae, at in* T *flavo ſquamula utrinque lacera.*
T. anguſtifolium Villarſii eſt T. ſimplex, ut ex ſpeciminibus Clariſſ. auctoris mecum benigne communicatis vidi. W.

18. THALICTRUM *lucidum.*
T. caule folioſo ſulcato, foliis linearibus carnoſis. *Sp. pl.* 770. *Dalib. pariſ.* 162. † *Mill. dict. n.* 4.
T. minus lucidum, libanotidis coronariae foliis. *Pluk. alm.* 363. *t.* 65. *f.* 5.
T. minus alterum pariſienſium, foliis craſſioribus et lucidis. *Tournef. inſt.* 271.
Thalictrum 1, vel pratenſe 1. *Cluſ. hiſt.* 1. *p.* 234.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 321.
Glänzendes Thalictrum. W.
Habitat Pariſiis *et in* Hiſpania. ♃ (v. ſ.)
Planta, an ſatis diſtincta a T. flavo? videtur temporis filia.

19. THALICTRUM *aquilegifolium.*
T. fructibus pendulis triangularibus rectis, caule tereti. *Sp. pl.* 770. *Mill. dict. n.* 3. *Reyg. ged.* 2. *p.* 96. *Gort. ingr.* 87. *Iacq. auſtr. t.* 318. *Scop. carn. n.* 672.
T. ſeminibus triangularibus pendulis, ſtipulis caulinis.

Hort. cliff. 226. *Fl. ſuec.* 452, 487. *Roy. lugdb.* 285.

T. foliis triplicato-pinnatis, patulis, obtuſe trilobatis, ſtipulis ovatis, paniculis umbellatis. *Hall. helv. n.* 1141.

T. fructibus triangularibus, pedicellatis, pendulis. *Crantz. auſtr. p.* 107. *n.* 3.

T. majus, florum ſtaminibus purpuraſcentibus. *Bauh. pin.* 337.

T. majus, folliculis anguloſis, caule laevi. *Bauh. hiſt.* 3. *p.* 487.

β. T. (*atro-purpureum*) fructibus pendulis triangularibus rectis, caule tereti ſtricto pluries ramoſo nebuloſo. *Murr. ſyſt. veg. ed* 14. *p.* 513. *Iacq. hort.* 3. *t.* 61.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 321.

Akeleyblättriges Thalictrum W.

Habitat in Scania, Helvetia, Auſtria, Gedani, Ingria. ♃ (v. v.)

Petala 4, *Stamina* 60, *Piſtilla* 16.

Stamina 56 *ad* 57. Piſtilla 7, 8, 10. *Crantz. l. c.* R.

20. THALICTRUM *contortum.*

T. fructibus pendulis triangularibus contortis, caule ſubancipiti. *Sp. pl.* 770.

T. hybridum feminibus contortis. *Amoen. acad.* 3. *p.* 47.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 322.

Gedrehtes Thalictrum. W.

Habitat in Sibiria. *G. Demidoff.* ♃ (v. v.)

Petala 4, *Stamina* 60, *Piſtilla* 8, *antecedenti ſimile, ſed humilius floribus albis.*

Planta noſtra T. aquilegifolio eſt ſimillima ſed, folia *ſunt glauca,* flores*que albi, at ſemina nullo modo contorta ſed exacte cum iis T. aquilegifolii conveniunt. Numerus vero partium eſt idem ut eum Linnaeus indicavit.* Semina *vero imperfecta ſunt torta.* W.

21. THALICTRUM *petaloideum.*

T. ſcapo ſubumbellato, florum filamentis oblanceolatis coloratis anthera latioribus. *Sp. pl.* 77. *Pall. itin.* 3. *p.* 231.

β. T. (*ſtamineum*) ſtaminum filamentis petaliformibus lanceolatis. *Lin. ſuppl.* 271.

Dichtblütiges Thalictrum. W.

Habitat in Dauria. ♃ (v. ſ.)

Folia *radicalia duo, ſupradecompoſita ut in conge-*

neribus, ſed reliquis longe minora. Foliolis *ovatis, obtuſis, ſaepe trilobis.* Scapus *vix pedalis, ſtriatus, aphyllus, raro unico folio decompoſito et pedunculo laterali.* Flores *fere in umbellam terminalem congeſti.* Calyx *tetraphyllus, deciduus, rotundatus.* Stamina *numeroſa: Filamentis membranaceis, ſuperne magis dilatatis, coloratis, incarnatis Antheris latioribus.* Germina *ſeſſilia.*

T. ſtamineum ab hoc nullo modo eſt diverſum. W.

22. THALICTRUM *ſtyloideum.*

T. foliis tripartito-pinnatis, ſtylis baſi alatis. *Lin. ſuppl.* 271.

Breitgriffliges Thalictrum. W.

Habitat in Sibiria. ♃

Caulis *folio uno alterove.* Folia *radicalia et caulina tripartita, foliolis bipinnatis.* Pinnae *ſubcordatae tri- aut quinquedentatae.* Panicula *terminalis. Species diſtinctiſſima* ſtylis *ſubulatis utrinque baſi in alam ſemiorbicularem verticalem dilatatis.*

*23. THALICTRUM *japonicum.* †

T. ſeminibus laevibus, foliis tripinnatis, pinnulis inciſis ſerratis. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 337.

Didynamiſta Salviae ſimilis. *Thunb. jap.* 364. *n.* 74.

Japaniſches Thalictrum. W.

Habitat in Japonia.

1085. ADONIS.

Gen. plant. edit. Schreb. n. 952.

Cal. 4-phyllus. *Petala* quinis plura absque nectario. *Sem.* nuda.

1. ADONIS *aeſtivalis.*

A. floribus pentapetalis, fructibus ovatis. *Mill. dict. n.* 2. *Pollich. pal. n.* 525. *Schol. barb. n.* 452. *Mattuſch. ſil. n.* 401. *Doerr. naſſ. p.* 34. *Knorr. del* 2. *t. A.* 12. *Kniph. cent.* 5. *n.* 6. *Hoffm. germ.* 192. *Roth. germ. I.* 241. *II.* 623.

A. ſylveſtris, flore phoeniceo, ejusque foliis longioribus. *Bauh. pin.* 178.

A. flore pallido. *Cam. epit.* 648.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 324.

Sommer-Adonis. W.

Habitat inter ſegetes Europae *auſtralis.* ⊙ (v. v.)

Affinis nimium A autumnali. Flores rubri.

Datur *varietas floribus ochroleucis, baſi intus nigra vel violacea macula notatis, quam habet* Scholl. l. c. Petala 5 *ad* 8. R.

2. ADONIS *autumnalis.*

A. floribus octopetalis, fructibus ſubcylindricis. *Hort. upſ.* 156. *Scholl. barb. n.* 453. *Hoffm. germ.* 193. *Roth. germ. I.* 242. *II.* 623.

A. radice annua. *Hort. cliff.* 321. *Roy. lugdb.* 488. *Dalib. pariſ.* 162. *Sauv. monſp.* 253.

Adonis miniata. *Iacq. auſtr.* 4. *t.* 354.

A. radice annua, flore octopetalo. *Hall. helv. n.* 1158.

A. annua. *Mill. dict. n.* 1.

A. flore majore. *Bauh. pin.* 178.

A. hortenſis, flore minore atro - rubente. *Bauh. pin.* 178.

Flos adonis. *Cluſ. hiſt.* 1. *p.* 336.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 324.

Herbſt - Adonis. W.

Habitat inter ſegetes Europae *auſtralis.* ⊙ (v. v.)

Flores *atropurpurei.* Planta *uniflora, ramis ſterilibus.*

Cum priori conjungit Crantz. auſtr. p. 110. R.

*3. ADONIS *flammea.*

A. floribus octopetalis, petalis lanceolatis acutis, fructibus cylindricis, caule hirto. W.

A. floribus octopetalis, calycibus hirſutis, fructibus cylindricis. *Murr. ſyſt. veg. ed.* 14. *p.* 514. *Iacq. auſt.* 4. *t.* 355. *Hoſt. ſynop.* 308.

Brennendrothe Adonis. W.

Habitat inter ſegetes Auſtriae. ⊙

Radix *inſipida, quae amara in A. autumnali.* Caulis *hirtus.* Calyx *hirſutus.* Petala *lanceolata acuta, nec oblonga obtuſa, quandoque uno alterove dente inſtructa,* Fructus *cylindricus.* W.

4. ADONIS *vernalis.*

A. flore dodecapetalo: fructu ovato. *Gmel. ſib.* 4. *p.* 200. *Mill. dict. n.* 3. *et. ic. t.* 14. *Scop. carn. n.* 678. *Pall. it.* 1. *p.* 62. *et it. p.* 388. *Mattuſch. ſil. n.* 402. *Blackw. t.* 504. *Kniph. cent.* 2. *n.* 2. *Hoffm. germ.* 193. *Roth. germ. I.* 242. *II.* 624.

A. radice perenni, flore dodecapetalo. *Hall. helv. n.* 1157. *Hort. cliff.* 231. *Fl. ſuec.* 456, 492. *It. oel.* 73. *Roy. lugdb.* 488. *Sauv. monſp.* 253. *Pollich. pal. n.* 526.

A. (*Helleborus*) perennis, petalis plus octo. *Crantz. auſtr. p.* 110.

A. ſeminibus aequalibus apice craſſioribus, infra apicem ſt, latis, hirſutis. *Gouan. illuſtr.* 33.

A. apennina. *Iacq. auſtr. t.* 44. *ſec. Linneum in Syſt. Veg. p.* 427. R.

Helleborus niger tenuifolius, buphthalmi flore. *Bauh. pin.* 186.

Bupthalmum. *Dod. pempt.* 261.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 325.

Frühlings-Adonis. W.

Habitat in Oelandiae, Boruſſiae, Bohemiae, Germaniae, Helvetiae, *collibus apricis.* ♃ (v. v.)

Haec Soror A. apenninae, ut A. aeſtivalis eſt A. autumnalis.

5. ADONIS *apennina.*

A. floribus pentdecapetalis, fructu ovato. *Pall. itin.* 2 *p.* 93. *et* 388. *Kniph. cent.* 3. *n.* 2. *Gmel. it.* 1. *p.* 81.

A. ſeminibus angulatis acuminato-mucronatis glabris. *Gouan. illuſtr. p.* 33. R.

Helleborus niger ferulaceus, caule geniculato, flore magno tulipae minoris inſtar. *Mentz. pug. t.* 3. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 328.

Apenniniſche Adonis. W.

Habitat in Sibiria, Apenninis. ♃ (v. ſ.)

Planta *multiflora; ramis ſcilicet etiam floriferis.*

Difficillime diſtinguuntur plantae 5 *et* 4 *ſimillimae.* Caulis 4 *adſperſus pilis;* 5 *punctis farinoſis pellucidis.* Rami 4 *ex caulis parte inferiore:* 5 *ex alis ſuperioribus.* Pedunculus 4 *breviſſimus;* 5 *longior.* Folia *in* 4 *minus pinnata;* 5 *magis diviſa, latiora, nitidiora.* Calyx *in* 4 *griſeus ſtriatus villoſus concavus;* 5 *luteo-vireſcens, glaber, avenius, planus.* Petala *in* 4 *lanceolata, ſaepius duodecim;* 5 *ovata, imbricata, ſaepius quindecim.* Flos *in* 4 *unicus; in* 5 *plures.* Stamina *in* 5 *valde reflexa.* Planta 5 *diutius aeſtate durat,* 4 *vero vernalis eſt.*

6. ADONIS *filia.*

A. foliis bipinnatis, foliolis linearibus pinnatifidis. *Lin. ſuppl.* 271.

Mohrrübenblättrige Adonis. W.

Habitat ad Cap. b. ſpei. ♃

Filia *A. capenſis, cui adeo ſimilis, ut ne quidquam differat caule, umbella, calyce, petalis, ſtaminibus.* Folia *umbellatarum (Dauci) pinnata.* Folia *partialia* inferiora *longius petiolata.* Folia *propria pinnatifida.* Pinnae *inaequales.*

7. ADONIS *capenſis.*

A. foliis biternatis, foliolis ſubcordatis. *Lin. ſuppl.* 272.

A. floribus decapetalis, fructibus depreſſis, foliis duplicato-ternatis foliolis ſerratis cordatis. *Sp. pl.* 772. *Berg. cap.* 149.

Actaea trifoliata hirſuta, foliis ſcabris, flore ſulphureo rariore. *Burm. afr.* 147. *t.* 51.

Chriſtophoriana africana ranunculoides, foliis rigidis. *Boerh. lugdb.* 2. *p.* 62.

Ranunculus aethiopicus, foliis rigidis, floribus ex luteo vireſcentibus. *Comm. hort.* 1. *p.* 1. *t.* 1. *Raj. ſuppl.* 316.

Ranunculus perelegans Franqueville, foliis aculeatis. *Lob. illuſtr.* 135. *

Aſtrantia africana, laſerpitii foliis latioribus rigidis et ſpinoſis. *Moriſ. hiſt.* 3. *p.* 279.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 328.

Capiſche Adonis. W.

Habitat ad Cap. b. ſpei. ♃ (v. ſ.)

Folia *biternata, radicalia: foliolis ſubcordatis, ſerratis, rigidis: Petiolis hirtis.* Scapi *aphylli, teretes, piloſi, terminati* Umbella *compoſita.* Involucra *et* Involucella *hexaphylla, lanceolata.* Pedunculi *partiales tres exteriores; tres interiores et unicus centralis uniflorus; reliqui quadriflori.* Calyx 5-*phyllus, ovato-oblongus, concavus, albidus.* Petala 10, *lanceolata, alba, calyce duplo longiora.* Filamenta *multa, linearia, petalis ½ breviora.* Antherae *binae, ovatae, parvae.* Piſtilla *multa, in capitulum congeſta.* Styli *extrorſum flexi, longitudine ſtaminum.* Stigmata *obtuſa.* Foliatio *involuta.*

8. ADONIS *veſicatoria.*

A. foliis biternatis, foliolis ſerratis glabris, floribus decapetalis. *Lin. ſuppl.* 272. *Ait. Kew.* 2. *p.* 264.

Imperatoria ranunculoides africana enneaphyllos, Laſerpitii lobatis foliis rigidis margine ſpinoſis. *Pluk. alm.* 198. *t.* 95. *f.* 2.

Blaſenziehende Adonis. W.

Habitat ad Cap. b. ſpei. ♃

Folia *Afris pro Veſicatoriis adhibentur.*

1086. RANUNCULUS.

Gen. pl. ed. Schreb. n. 953.

Cal. 5-phyllus. *Petala* 5, intra ungues poro mellifero. *Sem.* nuda.

* *Foliis ſimplicibus.*

1. RANUNCULUS *Flammula.*

R. foliis ovato-lanceolatis petiolatis, caule declinato. *Hort. cliff.* 228. *Fl. ſuec.* 458, 493. *Roy. lugdb.* 489. *Dalib. pariſ.* 163. *Pollich. pal. n.* 527. *Reyg. ged.* 1. *p.* 143. *Scop. carn. n.* 682. *Flor. dan. t.* 575. *Hoffm. germ.* 193. *Roth. germ. I.* 237. *II.* 609.

R. foliis ovato-oblongis integerrimis, caule procumbente. *Fl. lapp.* 325.

R. caule declinato, foliis elliptico-lanceolatis, ſubſerratis. *Hall. helv. n.* 1182.

R. foliis ſubintegerrimis, calycibus pentaphyllis coloratis: floribus hexapetalis. *Neck. gallob. p.* 238.

R. longifolius paluſtris minor. *Bauh. pin.* 180.

Flammula Ranunulus. *Dod. pempt.* 432.

β. Ranunculus paluſtris, foliis ſerratis. *Bauh. pin.* 180. *Dodon. pempt.* 433.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 330.

Beiſsender Ranunkel. W.

Habitat in Europae *paſcuis udis.* ♃ (v. v.)

2. RANUNCULUS *reptans.*

R. foliis linearibus, caule repente. *Fl. lapp.* 236. *t.* 3. *f.* 5. *Fl. ſuec.* 459, 495. *Mattuſch. ſil.* 405. *Fl. dan. t.* 108. *Kniph. cent.* 9. *n.* 83. *Hoffm. germ.* 193. *Roth. germ. I.* 237. *II.* 610.

R. caule reptante, foliis linearibus, faſciculatis. *Hall. helv. n.* 1183.

R. flammeus minimus, ſpergulae folio. *Buxb. cent.* 5. *app.* 41.

R. repens, gramineis foliis, e ſingulis geniculis radices agens. *Amm. ruth.* 80. *t.* 13. *f.* 1.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 331.

Feinblättriger Ranunkel. W.

Habitat in Suecia, Ruſſia, Helvetia, Germania *ad ripas lacuum, antecedenti valde affinis et forte varietas.* ♃ (v. ſ.)

Non eſt varietas praecedentis, differt caule *filiformi repente*, foliis *linearibus in petiolum attenuatis integerrimis*, pedunculis *uncialibus unifloris ſolitariis erectis*, floribus *multo minoribus.* W.

3. RANUNCULUS *Lingua.*

R. foliis lanceolatis, caule erecto. *Hort. cliff.* 228. *Fl. ſuec.* 457. *Roy. lugdb.* 489. *Dalib. pariſ.* 163. *Scop. carn. n.* 683. *Pollich. pal. n.* 528. *Crantz. auſtr. p.* 120. *n.* 19. *Flor. dan. t.* 755. *Hoffm. germ.* 193. *Roth. germ. I.* 237. *II.* 610.

R. erectus, ramoſus, foliis enſiformibus, ſubſerratis. *Hall. helv. n.* 1181.

R. longifolius paluſtris major. *Bauh. pin.* 34.

R. lanceolatus major. *Tabern. ic.* 48.

β. R. flammeus, latiori plantaginis folio marginibus piloſis. *Pluk. alm.* 312.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 331.

Groſser Ranunkel. W.

Habitat in Europae *borealioris foſſis, aquis limoſis.* ♃ (v. v.)

Caulis *ramoſus, folioſus, ſaepe ſeſquipedalis.* Folia *ſubdenticulata, ſcabra pilis brevibus appreſſis.* Pedunculi *oppoſitifolii uniflori.*

4. RANUNCULUS *nodiflorus.*

R. foliis ovatis petiolatis, floribus ſeſſilibus. *Hort. cliff.* 228. *Dalib. pariſ.* 164.

R. pariſienſis pumilus, plantaginellae folio. *Pet. gaz.* 40. *t.* 25. *f.* 4. *Vaill. act.* 1719. *p.* 52. *t.* 4. *f.* 4.

β. R. ſiculus, folio rotundo vix ſerrato. *Pet. gaz.* 39. *t.* 24. *f.* 9.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 332.

Knotenblütiger Ranunkel. W.

Habitat Pariſiis *et in* Siciliae *locis paludoſis.* (v. ſ.)

Habitu Limoſellae aquaticae haud diſſimilis. W.

5. RANUNCULUS *gramineus.*

R. foliis lanceolato-linearibus indiviſis, caule erecto laeviſſimo paucifloro. *Syſt. veg.* 428. *Mill. dict. n.* 5.

R. foliis lanceolato-linearibus ſeſſilibus, caule erecto, radice bulboſa. *Sp. pl.* 2. *p.* 773. *Hort. cliff.* 228. *Roy. lugdb.* 489. *Dalib. pariſ.* 163. *Sauv. monſp.* 75.

R. foliis lanceolato-linearibus ſeſſilibus, caule erecto, radice faſciculata. *Ger. prov.* 384.

Ranunculus, gramineo folio, bulboſus. *Bauh. pin.* 181.

β. R. montanus gramineo folio. *Bauh. pin.* 180.

γ. R. montanus, folio gramineo, multiplex. *Bauh. pin.* 181.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 333.

Grasblättriger Ranunkel. W.

Habitat in Galliae *pratis udis.* ♃ (v. ſ.)

Folia *lanceolato-linearia, ſubnervoſa ſ. ſtriata.* Caulis *teres, non ſtriatus.* Flores *lutei* 1, 2, 3, *minime albi.*

Diſtinguendus floribus luteis, radice tuberoſa a R. pyrenaeo, *qui eſt floribus albis, radice bulboſa.*

6. RANUNCULUS *pyrenaeus.*

R. foliis linearibus indiviſis, caule erecto ſtriato ſubbifloro. *Mant.* 248. *Iacq. miſcell.* 1. *p.* 154. *t.* 18. *f.* 1. *Villars delph.* 3. *p.* 732.

R. foliis radicalibus lineari-lanceolatis nervoſis, ſcapo unifloro, germinibus uncinatis. *Hohenwarth et Reiner. it.* 1. *p.* 187. *

Ranunculus plantagineus. *Allion. pedem. n.* 1445. *t.* 76. *f.* 1.

R. radicibus faſciculatis: imo caule reticulato, foliis gramineis. *Hall. helv. n.* 1180.

R. foliis linearibus, caule ſubnudo, radice faſciculata comoſa. *Ger. prov.* 384.

R. alpinus pumilus, gramineo folio, flore albo. *Tournef.* 292.

R. pumilus gramineis foliis. *Bauh. hiſt.* 3. *p.* 850.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 333.

Pyrenäiſcher Ranunkel. W.

Habitat in Pyrenaeis *inque alpibus* Helvetiae, Delphinatus, Carinthiae. ♃ (v. ſ.)

Radix *ſubbulboſa, obducta reticulo tomentoſo: Radiculis faſciculatis.* Caulis *erectus, filiformis, laevis, digitalis folio uno alterove.* Folia *radicalia et caulina*

linearia, indiviſa, integerrima, laevia, erecta, pollicaria. Flos *terminalis* 1 *ſ.* 2. *pedunculatus, parvus, albus.* Fructus *ſubrotundus.* Stamina *ſubrotunda, mucronata.*

Allionii icon repraeſentat plantam duplo majorem foliisque latiſſimis. W.

7. RANUNCULUS *parnaſſifolius.*

R. foliis nervoſis, radicalibus ſubrotundo-ovatis ſubcordatis, caulinis ſeſſilibus ovato-lanceolatis. W.

R. foliis ſubovatis nervoſis lineatis integerrimis petiolatis, floribus umbellatis. *Sp. pl.* 774. *Gouan. illuſtr. p.* 34. *Villars delph.* 3. *p.* 735.

R. caule multifloro hirſuto, foliis radicalibus petiolatis ovatis integerrimis ſupra hirſutis, ſubtus nervoſis, calycibus purpureis. *Wulffen in Iacq. Collect.* 1. *p.* 191. *t.* 9. *f.* 3. * *Hohenwarth et Reiner itin.* 1. *p.* 190. *t.* 5. *

R. radicibus faſciculatis, imo caule reticulato, foliis cordiformibus. *Hall. helv. n.* 1179. *ſec. Gouan.* R.

R. montanus, graminis parnaſſi folio. *Tornef. inſt.* 286.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 334.

Parnaſſienblättriger Ranunkel. W.

Habitat in alpibus Helveticis, Delphinatus, Carinthiae *inque* Pyrenaeis. ♃ (v. ſ.)

Gemma *radicalis promit* Folia *duo, lanceolata, membranacea, ſupra terram eminentia.* Folia *ovata, obtuſa, coriacea, nervoſa, integerrima, ſupra pubeſcentia, imprimis baſi involuta, longius petiolata.* Caulis *ſimpliciſſimus, indiviſus, aphyllus.* Involucrum *terminale, diphyllum.* Pedunculi 2 — 4, *uniflori, pubeſcentes.* Calyx *purpuraſcens.* Corolla *albida vel purpuraſcens nervis purpureis.*

*8. RANUNCULUS *ophiogloſſoides.*

R. caule ſimplici erecto, foliis nervoſis, inferioribus ovatis ſubcordatis petiolatis, floralibus ſeſſilibus lanceolatis. W.

R. (*ophiogloſſifolius*) foliis integerrimis obtuſis, imis cordatis, caulinis ovato-lanceolatis, ſupremis linearibus, caule erecto. *Villars delph.* 3. *p.* 731. *t.* 49. *

Natterzungenblättriger Ranunkel. W.

Habitat in alpibus Delphinatus. ♃

Accedit ad praecedentem, differt vero, licet folii

inferioribus conveniat: foliis *caulinis petiolatis, petiolis dilatatis,* foliis *floralibus lanceolatis,* caule *erecto,* floribus *triplo minoribus luteis, denique radice fuſiformi.* W.

9. RANUNCULUS *amplexicaulis.*

R. foliis ovatis acuminatis amplexicaulibus, caule multifloro, radice faſciculata. *Hort. cliff.* 229. *Roy. lugdb.* 484. *Sauv. monſp.* 76. *Kniph. cent.* 2. *n.* 66.

R. montanus, folio plantaginis. *Bauh. pin.* 180. *Moriſ. hiſt.* 2. *p.* 444. *ſ.* 4. *t.* 30. *f.* 36.

R. dulcis, foliis latis rapiſtri perfoliatis, floribus albis. *Mentz. pug. t.* 8. *f.* 8.

R. pyrenaeus flore albo. *Cluſ. curr. app. alt.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 334.

Stengelumfaſſender Ranunkel. W.

Habitat in Helvetia, Pyrenaeis, Apenninis. ♃ (v. v.)

10. RANUNCULUS *bullatus.*

R. foliis ovatis ſerratis, ſcapo nudo unifloro. *Sp. pl.* 774. *Hort. cliff.* 229. *Desfont. fl. atl.* 1. *p.* 435. *

R. latifolius bullatus, aſphodeli radice. *Bauh. pin.* 181.

R. luſitanicus. *Dod. pempt.* 429.

R. grumoſa radice 1. *Cluſ hiſt.* 1. *p.* 238.

β. R. latifolius bullatus autumnalis, flore pleno et prolifero. *Moriſ. hiſt.* 2. *p.* 247. *ſ.* 4. *t.* 31. *f.* 49. 50. 51.

R. luſitanicus, folio ſubrotundo, parvo flore. *Tournef. inſt.* 286.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 335.

Blaſiger Ranunkel. W.

Habitat in Luſitania, Creta, *inque* Africa *boreali.* ♃

*11. RANUNCULUS *ſalſuginoſus.*

R. foliis ovatis ſubcordatisve apice tri- ſeu quinquedentatis, caule repente. W.

Ranunculus ſalſuginoſus. *Pall. itin. ed. min* 3. *p.* 173.

R. (*plantaginifolius*) foliis cordato ovatis, apice vel integris vel tridentatis, flagellis reptantibus. *Murr. comment. goett.* 1777. *p.* 39. *t.* 2. *

R. (*ruthenicus*) foliis ovatis ad apicem paucidentatis, caule repente. *Iacq. hort.* 3. *p.* 19. *t.* 31. *

R. foliis petiolatis ovatis integris aut varie inciſis, pedunculis radicatis nudis. *Gmel. ſib.* 4. *p.* 205.

R. repens flore in caule ſingulari, foliis varie ſectis. *Amman. ruth. p.* 81. *n.* 107. *t.* 13. *f.* 2.

Salz-Ranunkel. W.

Habitat locis ſalſis in Sibiria *tranſbaicalenſi.* ♃ (v. ſ.)

12. RANUNCULUS *Ficaria.*

R. foliis cordatis angulatis petiolatis, caule unifloro. *Fl. ſuec.* 460, 496 *Hort. cliff.* 228. *Mat. med.* 141. *Roy. lugdb.* 490. *Dalib. pariſ.* 167. *Oed. dan. t.* 499. *Gort. ingr. p.* 88. *Scop. carn. n.* 684. *Pollich. pal. n.* 529. *Ludw. ect. t.* 52. *Kniph. cent.* 1. *n.* 73. *Hoffm. germ.* 194.

R. calycibus triphyllis cucullatis. *Crantz. auſtr. p.* 120.

R. foliis radicalibus glabris cordato-ſubrotundo-anguloſis ſ. integerrimis: pedunculis unifloris. *Neck. gallob. p.* 237.

Ficaria. *Hall. helv. n.* 1160. *Blackw. t.* 51.

Ficaria ranunculoides. *Roth. germ. I.* 241. *II.* 622.

Chelidonia rotundifolia minor. *Bauh. pin.* 309.

Chelidonium minus. *Fuchſ. hiſt.* 866.

β. Chelidonia rotundifolia major. *Bauh. pin.* 209. *prodr.* 137.

Houttuyn, Lin. Pfl. Syſt. 7. *p.* 336.

Feigwarzen-Ranunkel. W.

Habitat in Europae *ruderatis, umbroſis, ſpongioſis.* ♃ (v. v.)

*13. RANUNCULUS *frigidus.* W.

R. foliis radicalibus cuneiformi-ovatis apice quinquedentatis, caulinis ſeſſilibus palmatis. W.

Kalter Ranunkel. W.

Habitat in Sibiriae *alpibus.* ♃ (v. ſ.)

Hunc ſub nomine R. gracilis accepi, cui floris magnitudine et forte colore ſimilis, tamen foliis abunde diſtinctus. Folia *radicalia plura petiolata vix ſemiuncialia ovato-cuneiformia integerrima, apice truncata dentibusque quinque magnis obtuſis inſtructa.* Caulis *erectus uni-vel biflorus.* Folia *caulina bina vel terna alterna remota ſeſſilia baſi cuneiformia profunde quinqueloba, lobis lanceolatis integerrimis acutiuſculis.* Calycis *foliola elliptica obtuſa hirſuta.* Petala *obcordata calyce majora videntur alba in ſpecimine ſicco.* W.

14. RANUNCULUS *Thora.*

R. foliis reniformibus ſubtrilobis crenatis: caulino ſeſſili, floralibus lanceolatis, caule ſubbifloro. *Sp. pl.* 775. *Scop. carn. n.* 685. *Hoffm. germ.* 194. *Iacq. auſtr.* 5. *t.* 442. *Villars delph.* 3. *p.* 729.

R. foliis duris, venoſis, reniformibus, rotunde crenatis, convexitate retuſa, acute ſerrata. *Hall. helv.* 1178.

R. foliis inferioribus ſubrotundo-reniformibus, crenatis; floralibus lanceolatis integerrimis, caule inferne nudo. *Iacq. vind.* 249 *obſ.* 1 *p.* 25. *t.* 15.
R. Pthora *Crantz. auſtr. p.* 119. *n.* 17.
Aconitum pardalianches 1, ſ. Thora major. *Bauh. pin.* 184.
Thora major. *Cam. epit.* 825.
Pthora valdenſium. *Lob. ic.* 604. *Cluſ. hiſt.* 1. *p.* 239.
β. Aconitum pardalianches alterum, ſ. Thora minor. *Bauh. pin.* 184.
Thora minor. *Cam. epit.* 826.
Limeum pardalianches genus, uno tantum folio. *Dalech. hiſt.* 1738.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 338.
Giftiger Ranunkel. W.
Habitat in alpibus Helveticis, Pyrenaicis, Auſtriacis *et* Delphinatus. ♃ (v. ſ.)

Calyx *coloratus, reflexus.* Plantae *α. β. conjunguntur ab Hallero, diſtinguuntur a Seguiero.*

* * *Foliis diſſectis et diviſis.*

15. RANUNCULUS *creticus.*

R. foliis radicalibus reniformibus crenatis ſublobatis, caulinis tripartitis lanceolatis integerrimis, caule multifloro. *Sp. pl.* 775. *Mill. dict. n* 3.
R. foliis radicalibus cordato-ſubrotundis, caulinis tripartitis ſeſſilibus lanceolatis integerrimis. *Roy. lugdb.* 490.
R. aſphodeli radice, creticus. *Bauh. pin.* 181.
R. creticus latifolius. *Cluſ. hiſt.* 1. *p.* 239.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 339.
Candiſcher Ranunkel. W.
Habitat in Creta. ♃ (v. ſ.)

Tota planta hirſuta. Folia *radicalia ampla quatuor uncias longa et quinque uncias lata, reniformia, leviter ſeptemloba, lobis obtuſiſſimis trifidis laciniis obſolete tridentatis.* Caulis *erectus ramoſus ſeſquipedalis et forte ultra, pennae anſerinae craſſior.* Folia *caulina baſi cuneiformiter attenuata profunde tripartita, lobis oblongis obtuſis divaricatis integerrimis vel interdum trifidis.* Corolla *flava magnitudine R. Linguae.* Petala *ſubrotunda.* W.

16. RANUNCULUS *caſſubicus.*

R. foliis radicalibus ſubrotundo-cordatis crenatis; caulinis digitatis dentatis, caule multifloro. *Sp. pl.* 775. *Reyg. ged.* 1. *p.* 143. *n.* 4.

Ranunculus rotundifolius vernus ſylvaticus major vel caſſubicus, folio thorae calthae. *Breyn. prodr.* 1. *p.* 45.

R. aconitifolius folio rotundo ad radicem praeſtolante. *Loeſel pruſſ.* 225. *t.* 72.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 340.

Caſſubiſcher Ranunkel. W.

Habitat in Caſſubia, Sibiria, Gedani. ♃

Simillimus videtur ſequenti, ſed ſecundum Loeſelii iconem Foliola *caulina ſunt lanceolata dentata, nec linearia integerrima.* W.

17. RANUNCULUS *auricomus.*

R. foliis radicalibus reniformibus crenatis inciſis; caulinis digitatis linearibus, caule multifloro. *Hort. cliff.* 229. *Fl. ſuec.* 462, 498. *Roy. lugdb.* 490. *Dalib. pariſ.* 167. *Leyſ. hal.* 487. *Gmel. ſib.* 4. *p.* 202. *n.* 47. *Crantz. auſtr. p.* 112. *Gort. ingr. p.* 89. *Scop. carn.* *n.* 687. *Pollich. pal. n.* 530. *Fl. dan. t.* 665. *Kniph. cent.* 2. *n.* 67. *Hoffm. germ.* 194. *Roth. germ. I.* 238. *II.* 611.

R. foliis radicalibus integris et ſemitrilobatis, rotunde crenatis, caulinis multipartitis linearibus integerrimis. *Hall. helv. n.* 1177.

R. rotundifolius vernus ſylvaticus. *Bauh. hiſt.* 3. *p.* 341. R.

R. nemoroſus ſ. ſylvaticus, folio ſubrotundo. *Bauh. pin.* 178.

Ranunculus 1 ſylveſtris. *Dalech. hiſt.* 1028.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 340.

Verſchiedenblättriger Ranunkel. W.

Habitat in Europae *paſcuis humidiuſculis.* ♃ (v. v.)

Flores *nobis menſe Frondeſcentiae apetali, Floreſcentiae di tripetali, Graſſificationis pentapetali.* Folium *radicale cordatum.*

18. RANUNCULUS *abortivus.*

R. foliis radicalibus cordatis crenatis, caulinis ternatis angulatis, caule ſubtrifloro. *Sp. pl.* 776.

R. foliis radicalibus crenatis, caulinis digitatis petiolatis. *Roy. lugdb.* 490.

R. foliis radicalibus reniformibus crenatis petiolatis; caulinis paucis digitatis ſeſſilibus. *Gron. virg.* 166.
R. virginianus, flore parvo, molliori folio. *Herm. lugdb.* 514.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 341.
Virginiſcher Ranunkel. W.
Habitat in Virginia, Canada.•

*19. RANUNCULUS *trilobus.*

R. caule erecto, foliis glabris, caulinis trilobis, pedunculis ſtriatis, ſeminibus compreſſis tuberculoſis. *Desfont. fl. atl.* 1. *p.* 437. *t.* 113.
Dreylappiger Ranunkel. W.
Habitat in arvis humidis prope Mayane *in* Africa *boreali.* ♃

Radices *fibroſae numeroſae.* Caulis *erectus glaber ſtriatus fere pedalis.* Folia *glabra; radicalia crenata; caulina inferiora et media petiolata, profunde triloba, lobis cuneiformibus, inaequaliter dentatis; ſuperiora laciniata, laciniis anguſto-lanceolatis.* Pedunculi *ſtriati uniflori.* Flores *minimi.* Corolla *lutea.* Semina *in capiculum rotundum aggregata parva compreſſa orbiculata cum acumine, utrinque tuberculoſa. Affinis R. parvifloro, differt:* caule *erecto,* foliis *caulinis glabris profunde trilobis.* Desfont. l. c.

20. RANUNCULUS *ſceleratus.*

R. foliis inferioribus palmatis; ſummis digitatis, fructibus oblongis. *Hort cliff.* 230. *Fl. ſuec.* 463, 499. *Roy. lugdb.* 490. *Gron. virg.* 63. *Dalib. pariſ.* 167. *Gmel. ſib.* 4. *p.* 203. *t.* 83. *f. a.* *Gunn. norv. n* 84. *Scop. carn. n* 688. *Pollich. pal. n.* 531. *Doerr. naſſ. p.* 193. *Fl. dan. t.* 371. *Hoffm. germ.* 194. *Roth. germ. I.* 238. *II.* 612.
R. foliis levibus, ſemitrilobatis, rotunde ſerratis, fructu ovato. *Hall. helv. n.* 1175.
R. fructu oblongo impatiente. *Crantz auſtr. p.* 111.
R. paluſtris. *Tabern.* 42. *Blackw. t.* 259
R. paluſtris, apii folio, laevis. *Bauh. pin.* 180.
R. ſylveſtris 1. *Dod. pempt.* 426.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 342.
Blaſenziehender Ranunkel. W.
Habitat ad Europae *foſſas et paludes.* ⊙ (v. v.)

21. RANUNCULUS *aconitifolius.*

R. foliis quinquelobis dentatis, lobis acuminatis, inter-

mediis trifidis, floralibus ſummis digitatis ſeſſilibus lanceolatis W.

R. foliis omnibus quinatis lanceolatis inciſo-ſerratis. *Hort. cliff.* 229. *Hort. upſ.* 156 *Fl. ſuec.* 2. *n.* 497. *Roy. lugdb.* 490. *Scop. carn. n.* 680. *Pall. itin.* 2. *p.* 568. *Crantz. auſtr. p.* 112 *n.* 4. *Pollich. pal. n.* 532 *Mattuſch. ſil. n.* 409. *Hoffm. germ.* 195. *Roth. germ. I.* 238. *II.* 613.

R. caule ramoſo multifloro, foliis venoſis, quinquelobis, lobis rhomboideis, acutis, acute ſerratis. *Hall. helv. n.* 1164.

R. albus, folio denſo. *Bauh hiſt.* 3. *p.* 860.

R. montanus, aconiti folio, albus, flore minore. *Bauh. pin.* 182.

R. montanus 4. *Cluſ. hiſt.* 1. *p.* 236.

β. R. folio aconiti, flore albo multiplici. *Bauh. pin.* 179. *Kniph. cent.* 2. *n.* 65.

R. pleno flore albo. *Cluſ. hiſt.* 1. *p.* 236.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 344.

Eiſenhutblättriger Ranunkel. W.

Habitat in alpibus Helvetiae, Galliae, Auſtriae, Sileſiae *inque montibus* Thüringiae. ♃ (v. v.)

Similis *R. platanifolio*, *ſed* folia *quinquepartita ad petiolum uſque*, *et* caulis *dimidio brevior. Ioh. Bauh.* *Simillimus ſequenti at diverſus:* foliorum *lobis acuminatis*, *foliorum* floralium *ſummorum laciniis lanceolatis*, corolla *duplo minori. Ceterum conveniunt*, foliis *radicalibus 5-7-vel-9-lobis, lobis tribus vel quinque intermediis leviter tripartitis*, floribus *albis*. W.

22. RANUNCULUS *platanifolius.*

R. foliis quinquelobis dentatis, lobis obtuſis, intermediis trifidis, floralibus ſummis digitatis ſeſſilibus lineari-ſubulatis. W.

R. foliis palmatis laevibus inciſis, caule erecto, bracteis linearibus. *Mant.* 79. *Oed. dan. t.* 111. *Gouan. illuſtr.* 35.

R. montanus, aconiti folio, flore majore. *Bauh. pin.* 182.

R. montanus albus. *Dalech. hiſt.* 1031.

R. albus flore ſimplici, ſ. batrachium album. *Tabern.* 43. *ex Gouan.* R.

R. albus flore ſimplici. *Lob. ic.* 668. *Bauh. hiſt.* 3. *p.* 859.

Houttuyn Lin. Pfl. Syſt. 7. p. 345.
Platanusblättriger Ranunkel. W.
Habitat in Germaniae, Italiae *alpinis, et* Pyrenaeis. ♃ v. f.)

Statura *R. aconitifolii, floribus itidem albis, ſed* Folia *non compoſita, verum palmata, minus craſſa, magis inciſa:* Flores *majores.*

*23. RANUNCULUS *ſpicatus.*

R. foliis quinquelobis dentatis, caule ſimplici, ſeminibus ſpicatis. W.

R. foliis radicalibus rotundatis lobatis inciſis, caule ſimplici villoſo paucifloro, ſeminibus longe ſpicatis. *Desfont. fl. atl. 1. p. 438. t. 115.*

Aehrenſamiger Ranunkel. W.

Habitat in paludibus Algeriae. ♃

Radices *bulboſae.* Bulbi *numeroſi faſciculati oblongi.* Folia *radicalia petiolata villoſa orbiculata, baſi emarginata inaequaliter lobata, inciſa et crenata, crenulis rotundatis.* Caulis *erectus villoſus, leviter ſtriatus ſimplex, folia bi-aut quadripartita emittens, quandoque aphyllus.* Pedunculi *pauci uniflori.* Calyx *pentaphyllus villoſus.* Corolla *lutea magnitudine R. acris.* Petala 5 *obovata apice rotundata.* Semina *plana marginata in ſpicam denſam cylindraceam ſesquiuncialem diſpoſita.* Desf. l. c.

24. RANUNCULUS *illyricus.*

R. foliis ſericeo-villoſis ternatis, foliolis trifidis inciſis integerrimis, calyce reflexo W.

R. foliis ternatis integerrimis lanceolatis. *It. oel.* 71. *Fl. ſuec.* 471, 500. *Jacq. auſtr. t.* 222. *Pall. itin.* 3. *p.* 584. *Hoffm. germ.* 195. *Roth. germ. I* 238. *II.* 613.

R. foliis tripartitis: laciniis linearibus, caule multifloro, radice tuberoſa. *Hort. cliff.* 230. *Roy. lugdb.* 491. *Sauv. monſp.* 181.

R. foliis digitato-tripartitis, laciniis lanceolatis, integerrimis, calyce reflexo. *Crantz. auſtr. p.* 113. *n.* 5.

R. lanuginoſus anguſtifolius, grumoſa radice, major et minor. *Bauh. pin.* 181.

R. grumoſa radice 4. *Cluſ. hiſt.* 1. *p.* 240.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 546.

Illyriſcher Ranunkel. W.

Habitat in Oelandia, Hungaria, Narbona, Italia, Auſtria. ♃ (v. v.)

*25. RANUNCULUS *flabellatus.*

R. foliis ſimplicibus dentatis, ternatisque, foliolis laciniatis, caule ſimplici. W.

R. caule ſimplici hirſuto, foliis radicalibus flabelliformibus inciſo-lobatis, caulinis paucis multipartitis, caule ſubunifloro. *Desfont. fl. atl.* 1. *p.* 338. *t.* 114. *

Fächerblättriger Ranunkel. W.

Habitat circa Algeriam *in collibus incultis et humidis.* ♃

Simillimus R. aſiatico, differt vero: caule *ſimplici ſubunifloro,* Foliis *primordialibus cuneiformi-ovatis indiviſis inaequaliter dentatis baſi integerrimis,* Calycinis *foliolis ovatis acutis nec lanceolatis,* Corolla *flava,* Capitulo *ſeminum elliptico nec cylindraceo.* W.

26. RANUNCULUS *aſiaticus.*

R. foliis ternatis biternatisque: foliolis trifidis inciſis caule inferne ramoſo. *Sp. pl.* 777. *Mill. dict. n.* 11. *et ic t.* 216.

R. foliis tripartitis laciniatis, caule inferne ramoſo, radice tuberoſa. *Hort. cliff.* 230. *Hort. upſ.* 156. *Roy. lugdb.* 495.

R. grumoſa radice, ramoſus. *Bauh. pin.* 181.

R. aſiaticus polyclonos ſ. grumoſa radice 2. *Cluſ. hiſt.* 1. *p.* 241.

β. R. grumoſa radice, flore flavo vario. *Bauh. pin.* 181.

γ. R. grumoſa radice, flore albo. *Bauh pin.* 181.

δ. R. grumoſa radice, flore albo leviter crenato. *Bauh. pin.* 181.

ε. R. grumoſa radice, flore niveo. *Bauh. pin.* 181.

ζ. R grumoſa radice, flore phoeniceo minimo ſimplici. *Bauh. pin.* 181.

η. R. aſphodeli radice, flore ſanguineo. *Bauh. pin.* 181.

R. (*ſanguineus*) foliis ternatis biternatisque, foliolis trifidis obtuſis, caule ſimplici. *Mill. dict. n.* 10.

ϑ. R. aſphodeli radice, flore ſubphoeniceo rubente. *Bauh. pin.* 181.

R. aſphodeli radice, prolifer miniatus. *Bauh. pin.* 181.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 347.

Garten-Ranunkel. W.

Habitat in Aſia *et* Mauritania. ♃ (v. v.)

Caulis *erectus, ramoſus, pubeſcens, teres uti Pedunculi.* Folia *infima ſimplicia, lobata, inciſa, acuta, ſubtus pubeſcentia ut Petioli, ſuperiora tripartita et biternata.* Calyx *non reflexus.* Flos *coronarius.*

*27. RANUNCULUS *japonicus.*

R. foliis inciſo-ternatis, lobis inciſis dentatis, cauleque hirſuto. *Thunb. Act. Soc. Lin. Lond.* 2. *p.* 337.

Ranunculus aſiaticus. *Thunb. jap.* 241.

Japaniſcher Ranunkel. W.

Habitat in Japonia. ♃

Folia *radicalia rotundata, ſuprema fiſſa, laciniis lanceolatis, omnia trifida, laciniis iterum inciſis, dentata acuta utrinque hirſuta.* Petioli *pollicares usque pedales hirſuti.* Caules *ſtriati hirſuti vix folioſi flexuoſi, apice ramoſi in pedunculos.* Thunb. Act. l. c.

28. RANUNCULUS *rutaefolius.*

R. foliis pinnatis ternatisque, foliolis tripartito-multifidis inciſis, caule ſubſimplici, corolla polypetala, radice tuberoſa. W.

R. foliis ſupradecompoſitis, caule ſimpliciſſimo unifolio unifloro, radice tuberoſa. *Hort. cliff.* 230. *Roy. lugdb.* 491. *Mill. dict. n.* 6. *Crantz. auſtr. p.* 113. *n.* 6. *Allion. ped. n.* 1451. *t.* 67. *f.* 1.

R. foliis pinnatim decompoſitis, caule unifloro. *Villars. delph.* 3. *p.* 740. *

R. caule ſubtrifloro, foliis bipinnatis, pinnis ſeſſilibus ſulateſcentibus lobatis ſubpalmato-multifidis, radice fuſiformi longe undique fibroſa. *Wulffen in Jacq. Collect.* 1. *p.* 186. *t.* 6. *et* 7. *

R. caule unifloro, foliis imbricatis pinnatis, pinnis lobatis, obtuſe palmatis. *Hall. helv. n.* 1165.

R. rutaceo folio, flore ſuave rubente. *Bauh. pin.* 181. *Moriſ. hiſt.* 2. *p.* 448. *ſ.* 4. *t.* 31. *f.* 54.

R. alpinus coriandrifolio. *Pona bald.* 341. *Barr. ic.* 456.

R. folio rutaceo. *Bauh. hiſt.* 3. *p.* 414.

R. praecox 1, rutae folio. *Cluſ. hiſt.* 1. *p.* 232. *mala.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 347.

Rautenblättriger Ranunkel. W.

Habitat in alpibus Auſtriae, Delphinatus, Helvetiae. ♃ (v. ſ.)

Flos *albus.*

Caulis *plerumque uniflorus et unifolius, tamen ſaepius caule trifloro occurrit teſte Clariſſ. Wulffen; et ego ſpecimen e Sibiria habeo caule ramoſo.* Corolla *eſt ſemper decem vel-octopetala.* W.

29. RANUNCULUS *glacialis.*

R. foliis ternatis; foliolis tripartito-multifidis, caulinis ſeſſilibus, caule ſubtrifloro, calycibus hirſutis. W.

R. calycibus hirſutis, caule bifloro, foliis multifidis. *Fl. ſuec.* 464, 501. *Fl. dan. t.* 19.

R. caule bifloro, calyce hirſuto. *Fl. lapp.* 233. *t.* 3. *f.* 1.

R. caule ſubtrifloro, foliis repetito-trilobatis pulpoſis, calyce hirſuto. *Wulffen in Iacq. Collect.* 1. *p.* 189. *t.* 8. *et t.* 9. *f.* 1. 2.* *Hohenwarth et Reiner it.* 1. *p.* 193. *Hoſt. ſynop.* 311.

R. calycibus hirſute villoſis, caule ſubbifloro, foliis tripartitis, foliolis trifidis, laciniis ſubtrilobatis, lobis ſubacutis. *Schrank. ſalisb.* n. 519. *

R. foliis ternatim decompoſitis, foliolis carnoſis, intermediis ſimplicibus, caule multifloro. *Villars. delph.* 3. *p.* 738. *

R. foliis pulpoſis repetito-trilobatis, calycibus villoſis. *Hall. helv. n.* 1166.

R. montanus purpureus calyce villoſo. *Bauh. hiſt.* 3. *p.* 861. *Scheuch. alp.* 139. *t.* 20. *f.* 1. *Tournef. inſt.* 289.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 348.

Gletſcher-Ranunkel. W.

Habitat in ſummis alpibus Lapponiae, Helvetiae, Delphinatus, Auſtriae, Salisburgi. ♃ (v. ſ.)

Figura foliorum in Flora lapponica et Flora danica non eſt optima, ut primo intuitu diverſam eſſe ſpeciem crediderim, ſed ſpecimina ſicca lapponica quae poſſideo, eandem foliorum formam ac alia ex alpibus Europae auſtralioris habent. W.

*30. RANUNCULUS *Seguieri.*

R. foliis tripartitis, lobis multifido-laciniatis acutis, omnibus petiolatis, caule multifloro, calycibus glabris. W.

R. (*Seguieri*) foliis palmatis orbiculatim multifidis, caule proſtrato multifloro. *Villars delph.* 3. *p.* 737. *t.* 49. *

Ranunculus Columnae. *Allion. pedem. n.* 1453. *t.* 67. *f.* 3. 4.

R. (*Columnae*) foliis palmato-quinquepartitis, laciniis cuneiformibus pinnatifido-trifidis quinquefidisve, caule tereti villoſo ramoſo, pedunculis unifloris, radice aſphodeli fibroſa. *Wulffen in Iacq. Collect.* 4. *p.* 345. *

R. alpinus apii folio, flore albo magno. *Ponted. comp.* 117. *Segu. veron.* 1. *p.* 490. *t.* 12. *f.* 2. 3.
R. minor rutaefolio, flore ſimplici, grumoſa radice italicus. *Barr. icon.* 1153.
R. alter ſaxatilis aſphodeli radice. *Colum. ephr.* 312. *t.* 313?
Italiäniſcher Ranunkel. W.
Habitat in alpibus Delphinatus, Italiae, Carnioliae. ♃ (v. ſ.)
Calyx *coloratus glaber, apice tantum pilis aliquot inſtructus.* W.

31. RANUNCULUS *nivalis.*
R. foliis quinquelobis integerrimis, caulino ſeſſili digitato, caule unifloro. W.
R. calyce hirſuto, caule unifloro, foliis radicalibus palmatis: caulinis multipartitis ſeſſilibus. *Fl. lapp.* 232. *t.* 3. *f.* 2. *Fl. ſuec.* 465, 502. *Gunn. norv.* 627. *Retz. obſ.* 2. *p.* 19. *Swartz. Act. holm.* 1789. *p.* 47.
Ranunculus lapponicus. *Fl. dan. t.* 144.
β. Ranunculus idem pygmaeus. *Fl. lapp.* 232. *t.* 3. *f.* 3. *Houttuyn Lin. Pfl. Syſt.* 7. *p.* 349.
Schnee-Ranunkel. W.
Habitat in alpibus Lapponiae, Norvegiae. ♃
Planta tota glaberrima nequit eſſe R. monſpeliacus.

*32. RANUNCULUS *montanus.* W.
R. foliis quinquelobis dentatis, caulino ſeſſili digitato, laciniis lineari-lanceolatis integerrimis, caule unifloro. W.
Ranunculus nivalis. *Iacq. auſtr. t.* 325. 326. *Hoſt. ſyn.* 311.
R. (*nivalis*) foliis radicalibus quinquepartitis orbiculatim multifidis, caulinis palmatis integris ſeſſilibus, caule unifloro. *Villars delph.* 3. *p.* 742.
R. uniflorus, foliis glabris reniformibus ſemiquinquelobis. *Hall. helv. n.* 1168.
R. minimus alpinus luteus. *Bauh. hiſt.* 3. *p.* 861. *f. interior.*
Berg-Ranunkel. W.
Habitat in alpibus Helvetiae, Delphinatus, Salisburgi, Auſtriae. ♃ (v. ſ.)
Abunde diſtinctus a praecedente, quocum antea mixtus erat. W.

*33. RANUNCULUS *Gouani.* W.

R. foliis quinquelobis dentatis, caulino ſeſſili palmato, laciniis lanceolatis dentatis, caule unifloro. W.

R. (*pyrenaeus*) foliis radicalibus reniformibus trilobis quinquelobisve, caulino palmato ſeſſili. *Gouan. illuſt.* 33. *t.* 17. *f.* 1. 2. *

Gouaniſcher Ranunkel. W.

Habitat in Pyrenaeis *inque alpibus* Marmaroſienſibus Hungariae. ♃ (v. ſ.)

Accedit ad R. montanum ſed eſt omnibus partibus duplo major, et folio caulino dentato diverſus. W.

34. RANUNCULUS *alpeſtris.*

R. foliis radicalibus ſubcordatis obtuſis tripartitis: lobis trilobatis: caulino lanceolato integerrimo, caule ſubunifloro. *Sp. pl.* 778. *Iacq. auſtr. t.* 110. *Pall. it.* 2. *p.* 568.

R. uniflorus, foliis laevibus, ſemitrilobis, rotunde ſerratis, caulinis lingulatis. *Hall. helv. n.* 1167.

R. foliis ſubrotundis, ſplendentibus, ſemitrifidis, petalis cordatis albis. *Crantz. auſtr. p.* 113. *n.* 7.

R. foliis trilobis: lobis acutis; caule unifloro. *Scop. carn. n.* 1. *p.* 563. *n.* 4. *ed.* 2. *n.* 679.

R. alpinus humilis rotundifolius, flore majore et minore. *Bauh. pin.* 181.

R. alpinus humilis albus, folio ſubrotundo. *Segu. ver.* 1. *p.* 489. *t.* 12. *f.* 1.

R. montani 1 ſpecies 1 et 2. *Cluſ. hiſt.* 1. *p.* 234.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 349.

Alpen-Ranunkel. W.

Habitat in alpibus Auſtriacis, Helveticis. ♃ (v. ſ.)

Folia *ſimilia R. aquatilis natantibus.* Calyx *reflexus, glaber.* Petala *alba, biloba.*

35. RANUNCULUS *lapponicus.*

R. foliis tripartitis lobatis obtuſis, caule ſubnudo unifloro. *Sp. pl.* 778. *Fl. ſuec.* 461. 505. *

R. caule unifolio unifloro, foliis tripartitis. *Fl. lapp.* 251. *t.* 3. *f.* 4. *

Houttuyn Lin. Pfl. Syſt. 7. *p.* 350.

Lappländiſcher Ranunkel. W.

Habitat in alpibus Lapponicis. ♃ (v. ſ.)

*36. RANUNCULUS *hyperboreus.*

R. foliis profunde trilobis, lobis oblongis divaricatis caule

filiformi repente. *Rottb. Act. Haf.* 10. *p.* 458. *t.* 4. *f.* 16. *Retz. prod. fl. ſcand. ed. n.* 691. *Fl. dan. t.* 331.
Ranunculus Ammanni. *Gunn. norv. n.* 8°6. (*excluſo ſynonymo Ammanni.*)
R. foliis ſubrotundis trilobis integerrimis, caule repente. *Gmel. ſib.* 4. *p.* 204. *t.* 836. * (*excluſis ſynonymis Linnaei et Royeni.*)
Nordiſcher Ranunkel. W.
Habitat in Groenlandia, Islandia, Norvegia, *et* Sibiria, *prope* Ircutiam. ♃ (v. ſ.)

A R. lapponico caule *et* foliis *diverſiſſimus, nec ullo modo ſimilis R. ſalſuginoſo, quocum olim conjuxerat Gunnerus.* W.

37. RANUNCULUS *monſpeliacus.* †

R. foliis tripartitis crenatis, caule ſimplici villoſo ſubnudo unifloro. *Sp. pl.* 778. *Sauv. monſp.* 181. †
R. ſaxatilis, magno flore. *Bauh. pin.* 182. *prodr.* 96.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 351.
Franzöſiſcher Ranunkel. W.
Habitat Monſpelii.
Videtur differre a R. Gouani. W.

38. RANUNCULUS *penſylvanicus.*

R. calycibus reflexis, caule erecto, foliis ternatis trifidis, inciſis ſubtus piloſis. *Lin. ſuppl.* 272.
Ranunculus canadenſis. *Iacq. ic. rar.* 1. *t.* 105. *Miſc.* 2. *p.* 343. *
Penſylvaniſcher Ranunkel. W.
Habitat in Canada, Penſylvania. ☉ ♂ (v. v.)

Caulis *erectus ramoſus teres pilis patulis.* Folia *ternata ſublineata, ſubtus uti petioli piloſa.* Foliola *trifida acuta inciſa ſerrataque.* Pedunculi *teretes laeves vagi.* Calyx *reflexus, foliolis ovatis concavis.* Petala *quinque lutea vix calycis magnitudine.* Stamina *parva longitudine corollae.* Germinum *capitulum ovatum corolla longius.* Semina *ovata compreſſa mucronata.*

*39. RANUNCULUS *ternatus.*

R. calycibus reflexis, foliis omnibus ternatis, foliolis trifidis, caule multifloro. *Thunb. jap.* 241.
Dreyblättriger Ranunkel. W.
Habitat in Japonia.

40. RANUNCULUS *bulbòſus.*

R. calycibus retroflexis, pedunculis ſulcatis, caule erecto multifloro, foliis compoſitis. *Fl. ſuec.* 469, 504. *Dalib. pariſ.* 165. *Pollich. pal n.* 533. *Scop. carn. n.* 692. *Hoffm germ.* 195. *Kniph.cent.* 7. *n.* 74. *Fl. dan. t.* 551. *Roth. germ I.* 239. *II.* 614.

R. radice ſimplici globoſa. *Hort. cliff.* 230. *Roy. lugdb.* 491.

R. radice ſubrotunda, foliis hirſutis, ſeptilobis, lobis petiolatis. *Hall. helv. n.* 1174.

R. calycibus retroflexis, radice ſimplici globoſa. *Crantz. auſtr. p.* 114. *n.* 8.

R. bulboſus. *Lob. ic.* 666. *Fl. lapp.* 429.

R. pratenſis, radice verticilli modo rotunda. *Bauh. pin.* 179.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 351.

Knolliger Ranunkel.

Habitat in Europae *pratis, paſcuis.* ♃ (v. v.)

*41. RANUNCULUS *Philonotis.*

S. calycibus demum reflexis ſetoſis, pedunculis ſulcatis, foliis tripartitis inciſo-lobatis hirſutis, fructu ſubgloboſo. *Retz. prod fl. ſcand. ed.* 2. *n.* 693. *Retz. obſ.* 6. *p* 31.

R. (*Philonotis*) radice faſciculata, herba piloſa, caule erecto ramoſo, foliis ternatis, foliolis trilobis, lobis crenato-inciſis, apicibus puncto albo terminatis, pedunculis ſulcatis, calyce reflexo, fructu globoſo, ſeminibus compreſſis acuminatis. *Ehrh. Beitr.* 2. *p.* 145. *Hoffm germ.* 194.

R. (*hirſutus*) radice fibroſa annua, caule hirſuto, calycibus papilloſo-hiſpidis acuminatis, demum reflexis. *Ait. Kew.* 2. *p.* 268. *Curt. Lond.*

R. rectus, foliis pallidioribus hirſutis. *Bauh. hiſt.* 3. *p.* 417. *f.* 3. *Petiv. t.* 38.

R. paluſtris apii folio lanuginoſus. *Bauh. pin.* 180.

Rauher Ranunkel. W.

Habitat in Europae *udis.* ⊙ (v. v.)

*42. RANUNCULUS *polyrhizos.*

R. foliis radicalibus palmatis, caulinis ſeſſilibus digitatis, caule multifloro, radice faſciculata. W.

Ranunculus polyrhizos. *Stephan in litt.*

Vielwurzlicher Ranunkel. W.

Habitat in Sibiria. ♃ (v. ſ.)

Radix *faſciculata*, Folia *radicalia Ranunculi ſcelerati, triloba, lobis lateralibus bilobis, laciniis obtuſe tridentatis ut folia ipſa cuneiformibus.* Caules *digitales bini vel tres ex una radice erecti vel adſcendentes digitales ſimplices.* Folia *caulina ſeſſilia digitata, baſi cuneiformia, laciniis linearibus obtuſiuſculis integerrimis.* Pedunculi 2, 3 *vel* 4 *longi uniflori teretes non ſulcati tenuiſſime ut caulis pubeſcentes.* Calycis *foliola colorata glabra obtuſa patentia.* Petala *lutea magnitudine R. Flammulae.* Capitulum *ſeminum ſubrotundum.* Semina *ovata compreſſa laevia, ſtigmate obtuſo perſiſtenti coronata.* W

43. RANUNCULUS *repens.*

R. calycibus patulis, pedunculis ſulcatis, ſarmentis repentibus, foliis compoſitis. *Fl. ſuec.* 468, 505. *Gmel. ſib.* 4. *p.* 206. *Crantz auſtr. p.* 114. *n.* 9. *Scop. carn. n.* 689. *Pollich pal. n.* 531. *Blackw. t.* 31. *Hoffm. germ.* 196. *Roth. germ. I.* 239. *II.* 6 5.

R. caule reptante, radicato, foliis ſemitrilobatis, lobis petiolatis. *Hall. helv. n.* 1173.

R. foliis ternatis, foliolis petiolatis trifidis: medio productiore, caule multifloro. *Hort. cliff.* 230. *Gron. virg.* 166. *Roy. lugdb.* 491.

R. pratenſis repens hirſutus. *Bauh. pin.* 179. *Fl. lapp.* 230.

R. hortenſis 1. *Dod. pempt.* 425.

β. R. dulcis batrachium ſalutiferum. *Tabern.* 51. *Hall.*

Houttuyn Lin. Pfl. Syſt. 7. *p.* 352.

Kriechender Ranunkel. W.

Habitat in Europae *cultis.* ♃ (v. v.)

44. RANUNCULUS *polyanthemos.*

R. calycibus patulis, pedunculis ſulcatis, caule erecto, foliis multipartitis. *Fl. ſuec.* 467, 506. *Scholl. barb. n.* 447. *Pollich. pal. n.* 535. *Mattuſch. ſil. n.* 412. *Doerr. naſſ. p.* 194. *Hoffm. germ.* 196. *Roth. germ. I.* 239. *II.* 616.

R. ramoſus multiflorus, foliis ſemiſeptilobis, lobis tridentatis acutis. *Hall. helv. n.* 1171.

R. foliis inferioribus peltatis, tri-quadripartito-multifidis: caulinis linearibus ſubdenticulatis ſeſſilibus. *Neck. gallob. p.* 240.

R. polyanthemos ſimplex. *Lob. ic.* 666.
R. ſylveſtris. *Tabern. ic.* 42. *hiſt.* 117.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 353.
Vielblumiger Ranunkel. W.
Habitat in Europae *borealis gramineſis.* ♃ (v. v.)

Calyce *hirſuto et* Pedunculis *ſulcatis abunde a ſequenti diſtinctus.* W.

45. RANUNCULUS *acris.*

R. calycibus patulis, pedunculis teretibus, foliis tripartito multifidis: ſummis linearibus. *Fl. ſuec.* 465, 507. *Dalib. pariſ.* 164. *Gmel. ſib.* 4. *p.* 206. *n.* 53. *Scop. carn. p.* 690. *Pollich. pal n.* 536. *Hoffm. germ.* 196. *Knorr. del* 1. *t. H.* 1. *Roth. germ. I.* 239. *II.* 616.

R. foliis peltatis quinquangularibus multipartitis: laciniis linearibus, caule multifloro. *Hort. cliff.* 231. *Roy. lugdb.* 492.

R. foliis hirſutis, ſemitrilobatis, lobis lateralibus bipartitis, foliis caulinis ſemitrilobis. *Hall. helv. n.* 1160.

R. napellifolius. *Crantz. auſtr. p.* 114. *n.* 10. *t.* 4. *f.* 2. *cum praecedente conjungit.* R.

R. foliis inferioribus tri-quinque et partito-multifidis, ſuperioribus lineari-lanceolatis ſeſſilibus integerrimis. *Neck. gallob. p.* 239.

R. pratenſis erectus acris. *Bauh. pin.* 178. *Fl. lapp.* 228.

R. hortenſis. *Dod. pempt.* 426.

β. R. hortenſis erectus, flore pleno. *Bauh. pin.* 179. *Knorr. del.* 1. *t. H.* 2.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 354.
Scharfer Ranunkel. W.
Habitat in Europae *pratis, paſcuis.* ♃ (v. v.)

*46. RANUNCULUS *cappadocicus.* W.

R. calycibus patulis, pedunculo tereti, caule ſubbifido foliis cordatis trilobis dentatis. W.

R. orientalis dulcis Doronici radice. *Tournef. cor* 20.
Cappadociſcher Ranunkel. W.
Habitat in Cappadocia. ♃ (v. ſ.)

Radix *dura craſſitie pennae anſerinae facie Doronici.* Folia *radicalia ſubrotundo-cordata breviſſime triloba, lobis acuminatis dentibus paucis groſſis inſtructis,*

utrinque pili ſparſi adpreſſi. Folium *caulinum unicum ut radicalia petiolatum, et in pedunculi medio folium parvum ſeſſile lanceolatum integerrimum.* Caulis *ſpithamaeus pilis adpreſſis tectus ſimplex vel ſupra folium bifidus, pedunculis unifloris.* Corolla *flava magnitudine R. polyanthemi.* Capitulum *ſeminum ſubrotundum.* Semina *compreſſa uncinata.* W.

47. RANUNCULUS *lanuginoſus.*

R. calycibus patulis, pedunculis teretibus, caule petiolisque hirſutis, foliis trifidis lobatis crenatis holoſericeis. *Oed. dan. t.* 397. *Reyg. ged.* 2. *p.* 97. *Leers herb. n.* 429. *Mattuſch. ſil. n.* 414. *Doerr. naſſ. p.* 195. *Hoffm. germ.* 196. *Roth. germ. I.* 240. *II.* 617.

R. foliofus ramoſus erectus, foliis ſemitrilobatis, hirſutis, vaginis ſericeis. *Hall. helv. n.* 1172.

R. foliis tripartitis, acute ſinuatis lanuginoſis. *Crantz. auſtr. p.* 118.

R. foliis trilobis, hiſpidis: calycibus patulis, caule multifloro tereti. *Scop. carn.* 1. *p.* 564. *n.* 5. *ed.* 2. *n.* 691.

R. tomentoſus, calycibus retroflexis, foliis radicalibus tripartitis lobis inciſis; caulinis multipartitis ſeſſilibus. *Ger. prov.* 387.

R. foliis radicalibus pinnatis. *Sauv. monſp.* 227.

R. montanus lanuginoſus, foliis ranunculi pratenſis repentis. *Bauh. pin.* 182. *prodr.* 96.

R. magnus valde hirſutus, flore luteo. *Bauh. hiſt.* 3. *p.* 417.

β. Ranunculus montanus ſubhirſutus, geranii folio. *Bauh. pin.* 182.

R. montanus ſubhirſutus latifolius. *Bauh. prodr.* 96. *t.* 96.

R. nemoroſus hirſutus, foliis caryophyllatae. *Loeſ. pruſſ.* 220. *t.* 71.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 355.

Wolliger Ranunkel. W.

Habitat Monſpelii, *in* Helvetia, Germania, Auſtria, Carniolia, Luſatia ♃ (v. v.)

Caulis *altus, hirtus, erectus.* Folia *Aconiti ſ. Ran. acris, ſed magis inciſa.* Semina *acuta, compreſſa.* Pedunculi *longi, teretes.* Macula *pallida ad ſinus foliorum, quae tactu craſſiora ſentiuntur.*

48. RANUNCULUS *chaerophyllus.*

R. calycibus retroflexis, pedunculis ſulcatis, caule erecto

unifloro, foliis compoſitis lineari-multifidis. *Sp. pl.* 780. *Guett. ſtamp.* 275. *Dalib. pariſ.* 166. *Villars delph.* 3. *p.* 752.

R. foliis compoſitis, caule ſimplici villoſo ſubnudo unifloro, radice tuberoſa. *Linn. goett.* 129.

R. grumoſa radice, folio Ranunculi bulboſi. *Bauh. pin.* 181. *prodr.* 96.

R. chaerophyllos, aſphodeli radice. *Bauh. pin.* 181. *Barr. ic.* 581.

R. montanus leptophyllos, aſphodeli radice. *Col. ecphr.* 1. *p.* 312. *t.* 311.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 356.

Kerbelblättriger Ranunkel W.

Habitat in Gallia, Italia. ♃ (v. ſ.)

*49. RANUNCULUS *millefoliatus.*

R. foliis ſupradecompoſitis linearibus, calycibus piloſis, caule ramoſo ſericeo villoſo. *Vahl ſymb.* 2. *p.* 63. *

R. foliis multifariam decompoſitis lineari-ſubulatis, caule ſubaphyllo villoſo paucifloro, calycibus erectis hirſutis. *Desfont. fl. atl.* 1. *p.* 441. *t.* 116. *

Feinblättriger Ranunkel. W.

Habitat in regno Tunetano. ♃

50. RANUNCULUS *parvulus.*

R. hirtus, foliis trilobis inciſis, caule erecto ſubunifloro. *Mant.* 79. *Gmel. It.* 2. *p.* 196.

R. foliis radicalibus et caulinis inferioribus trilobis crenatis, ſummis trifidis dentatis, caule bifloro erecto. *Gouan. monſ.* 270. *

R. arvenſis parvus, folio trifido. *Bauh. pin.* 79.

R. ſaxatilis minimus hirſutus. *Bauh. pin.* 182. *prodr.* 96.

R. minimus apulus. *Col. ecphr. t.* 316.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 357.

Kleiner Ranunkel. W.

Habitat Monſpelii, *in* Italia, Ruſſia.

Caulis *ſolitarius, hirtus, filiformis, longitudine digiti ſ. ſemipedalis.* Folia *pauca, petiolata, ſimplicia, trifida, dentata, hirta.* Flos *fere foliis major, luteus.* Calyx *membranaceus, villoſus.*

*51. RANUNCULUS *oxyſpermus.*

R. foliis radicalibus oblongis obtuſis ſinuato-dentatis, caulinis ſeſſilibus digitatis inciſis, ſeminibus ariſtatis. W.

Ranunculus oxyſpermus Marſchallii. *Stephan. in litt.*
Spitzſamiger Ranunkel. W.
Habitat in Sibiria *ad fluvium* Tereck. ☉ (v. ſ.)

Planta *videtur annua.* Folia *radicalia petiolata ovata obtuſa inaequaliter ſinuato-dentata utrinque uti petioli piloſa.* Caulis *erectus pedalis et ultra ramoſus piloſus.* Folia *caulina ſeſſilia digitata, laciniis inferiorum inaequaliter pinnatifidis, ſuperiorum linearibus integerrimis.* Calyx *reflexus.* Corolla *flava magnitudine R. bulboſi.* Capitula *ſeminum elliptica.* Semina *compreſſiuſcula acuta ſubulato-ariſtata, ariſtis erectis.* W.

52. RANUNCULUS *arvenſis.*

R. ſeminibus aculeatis, foliis ſuperioribus decompoſitis linearibus. *Hort. cliff.* 229. *Flor. ſuec.* 470, 508. *It. oel.* 105. * *Roy. lugdb.* 491. *Dalib. pariſ.* 166. *Oed. dan.* 219. *Gmel. ſib.* 4. *p.* 205. *Scop. carn. n.* 693. *Pollich. pal. n.* 537. *Kniph. cent.* 12. *n.* 82. *Hoffm. germ.* 196. *Roth. germ. I.* 240. *II.* 618.
R. ſeminibus aculeatis, foliis tripartitis, lobis longe petiolatis, bipartitis et tripartitis, acute inciſis. *Hall. helv. n.* 1176.
R. echinatus. *Crantz. auſtr. p.* 118.
R. arvenſis echinatus. *Bauh. pin.* 179.
R. ſylveſtris 3. *Dod. pempt.* 427.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 558.
Acker-Ranunkel. W.
Habitat in Europae *auſtralioris agris.* ☉ (v. v.)

53. RANUNCULUS *muricatus.*

R. ſeminibus aculeatis, foliis ſimplicibus lobatis obtuſis glabris, caule diffuſo. *Hort. cliff.* 229. *Hort. upſ.* 157. *Roy. lugdb.* 491. *Sauv. monſp.* 204. *Hoffm. germ.* 196.
R. paluſtris echinatus. *Bauh. pin.* 180. *prodr.* 95. *hiſt.* 3. *p.* 858.
R. creticus echinatus latifolius. *Alp. exot.* 263. *t.* 262.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 559.
Stachelſamiger Ranunkel. W.
Habitat in Europae *auſtralis foſſis humentibus.* ☉ (v. v.)

54. RANUNCULUS *parviflorus.*

R. ſeminibus muricatis, foliis ſimplicibus laciniatis acu-

tis hirſutis, caule diffuſo. *Sp. pl.* 780. *Hudſ. angl.* 242.

R. hirſutus annuus, flore minimo. *Pluk. alm.* 311. *t.* 55 *f.* 1. *Raj. angl* 3. *p.* 248. *t.* 12. *f.* 1. *Moriſ. hiſt.* 2. *p.* 440. *ſ* 4. *t.* 28. *f.* 21.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 359.

Kleinblütiger Ranunkel. W.

Habitat in Europa *auſtrali* ⊙ (v. v.)

Differt a R. muricato *flore parvo; Caule magis proſtrato; Foliis profunde lacinulatis; Seminibus latere tantum tuberculis, nec aculeis muricatis.*

55. RANUNCULUS *orientalis.*

R. ſeminibus ſpinoſo-ſubulatis recurvis, calycibus reflexis, foliis multifidis. *Sp. pl.* 781.

R. lesbius pulſatillae folio flore magno. *Tournef. cor.* 20.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 360.

Orientaliſcher Ranunkel. W.

Habitat in Oriente. ⊙ (v. ſ.)

Caulis *digitalis ramis tribus vel quatuor unifloris.* Folia *petiolata ſupradecompoſita vel potius ex tribus foliolis parvis irregulariter pinnatis compoſita, pinnulis lineari-oblongis obtuſis.* Calyx *reflexus,* Corolla *ratione plantae magna.* Capitulum *ſeminum ſubrotundum.* Semina *oblonga compreſſa ſubulato-ariſtata, ariſtis uncinatis divaricatis.* W.

56. RANUNCULUS *grandiflorus.*

R. caule erecto bifolio, foliis multifidis; caulinis alternis ſeſſilibus. *Sp. pl.* 781. *Roy. lugdb.* 492.

R. orientalis, aconiti folio, flore maximo. *Tournef. cor.* 25.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 360.

Groſsblütiger Ranunkel. W.

Habitat in Oriente. ♃ (v. ſ.)

57. RANUNCULUS *falcatus.*

R. foliis cuneiformibus tripartitis, laciniis multifidis filiformibus, ſeminibus falcatis, ſcapo nudo unifloro. W.

R. foliis filiformi-ramoſis, ſeminibus falcatis, ſcapo nudo unifloro. *Hort. upſ.* 157. *Sauv. monſp.* 205. *Gron. orient.* 69. *Iacq. vind.* 250. *auſtr. t.* 48. *Hoffm. germ.* 197. *Roth germ. II* 618.

R. teſticulatus. *Crantz. auſtr. p.* 119.

Myoſurus foliis ramoſis. *Hort. cliff.* 117. *Roy. lugdb.* 492.
Melampyrum luteum minimum. *Bauh. pin.* 234.
Ranunculus ceratocephalus, ſeminibus falcatis in ſpicam adactis. *Moriſ. hiſt.* 2. *p.* 440. *ſ.* 4. *t.* 28. *f.* 22.
R. alopecuroides, ajugae foliis. *Boc. ſic.* 28. *t.* 14. *f.* 3.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 360.
Sichelſamiger Ranunkel. W.
Habitat inter ſegetes Europae *auſtralis et* Orientis. ⊙ (v. v.)

*58. RANUNCULUS *polyphyllus.*

R. foliis ſubmerſis oblongis petiolis capillaceis, natantibus cuneiformibus trilobis, emerſis ellipticis, caule erecto. W.
Ranunculus polyphyllus. *Waldſtein et Kitaibel pl. rar. hung.*
Vielblättriger Ranunkel. W.
Habitat in aquis Hungariae. ⊙ (v. ſ.)

Caulis *dodrantalis erectus ſuperne ramoſus*, Folia *ſubmerſa numeroſiſſima totum caulem tegentia oblonga unguicularia petiolo capillari unciali ſuffulta*, Folia *natantia parva unguicularia cuneiformia triloba integerrima petiolo craſſiori ſemiunciali.* Rami *unciales erecti ex aqua prominent, geruntque folia breve petiolata quadri- vel ſexlinearia elliptica utrinque attenuata obtuſa.* Flores *admodum parvi lutei.* W.

59. RANUNCULUS *hederaceus.*

R. foliis ſubrotundis trilobis integerrimis, caule repente. *Hort. cliff.* 231. *Roy. lugdb.* 492. *Dalib. pariſ.* 167. *Gort. gelr.* 325. *Pal. itin.* 2. *p.* 511. *Leers. herborn. n.* 431. *Pollich. pal. n.* 538. *Reich. flor. n.* 1010. *Doerr. naſſ. p.* 195. *Fl. dan. t.* 321. *Weber. ſpicil. fl. goett. p.* 16. *Hoffm. germ.* 197. *Roth. germ. I.* 240. *II.* 619.
R. aquaticus hederaceus luteus. *Bauh. pin.* 180. *Moriſ. hiſt.* 2. *p.* 441. *ſ.* 4. *t.* 29. *f.* 29.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 362.
Epheublättriger Ranunkel. W.
Habitat in aquis vadoſis Angliae, Belgii, Germaniae, Sibiriae. ♃ (v. ſ.)

Corolla *parva, petalis albis acutis.* W.

60. RANUNCULUS *aquatilis.*

R. foliis ſubmerſis capillaceis, emerſis peltatis. *Fl. ſuec.* 2. *n.* 509. *Gmel ſib.* 4. *p.* 207. *Crantz. auſtr. p.* 118. *Scop. carn. n.* 681. *Pollich. pal. n.* 539.

R. (*heterophyllus*) foliis ſubmerſis capillaceis, emerſis reniformi-palmatis. *Hoffm. germ.* 197. *Roth. germ. I.* 240. *II.* 619.

R. fluitans, petiolis unifloris, foliis imis capillaribus, laciniis divergentibus. *Hall. helv. n.* 1163.

R. foliis natantibus peltato-multipartitis: immerſis capillaceis; nectariis tubulatis. *Neck. gallob. p.* 242.

R. foliis inferioribus capillaceis, ſuperioribus peltatis. *Fl. lapp.* 234. *Roy. lugdb.* 492.

R. aquaticus, folio rotundo et capillaceo. *Bauh. pin.* 180.

R. aquatilis. *Dod. pempt.* 387.

β. R. foliis ſubmerſis capillaceis. *Fl. ſuec.* 1. *n.* 472. *Dalib. pariſ.* 168.

Ranunculus aquatilis β. *Roth. germ. II.* 620.

R. foliis omnibus capillaceis circumſcriptione rotundis. *Hort. cliff.* 232. *Roy. lugdb.* 492.

R. aquaticus albus, circinatis tenuiſſime diviſis foliis. *Pluk. alm.* 311. *t.* 55. *f.* 2.

Millefolium aquaticum cornutum majus. *Bauh. pin.* 141.

γ. R. fluitans, petiolis unifloris, foliis capillaribus, laciniis divergentibus. *Hall. helv. n.* 1162. *Pollich. l. c.* β.

R. (*aquatilis*) foliis omnibus capillaceis, laciniis divergentibus. *Hoffm. germ.* 197. *Roth. germ I.* 241. *II.* 620.

R. aquaticus capillaceus. *Bauh. pin.* 180.

Millefolium aquaticum, foliis abrotani, ranunculi flore et capitulo. *Bauh. pin.* 141.

Ranunculus trichophyllos aquaticus medio luteus. *Col. ecphr.* 1. *p.* 315. *t.* 316.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 363.

Waſſer-Ranunkel. W.

Habitat in Europae *aquis undoſis, foſſis, rivulis.* ♃ (v. v.)

Variat foliorum forma, tamen a ſequenti diverſiſſimus. W.

*61. RANUNCULUS *fluviatilis.*

R. foliis omnibus dichotomo-capillaceis, caule natante. W.

R. (*fluviatilis*) foliis omnibus capillaceis, laciniis parallelis longiſſimis. *Hoffm. germ.* 197. *Roth. germ. I.* 241. *II.* 621.

Ranunculus peucedanifolius. *Desfont. fl. atl.* 1. *p.* 444.

Ranunculus aquatilis δ. *Sp. pl.* 782.

R. fluitans, petiolis unifloris, foliis longiſſimis, laciniis parallelis. *Hall. helv. n.* 1161. *Fl. dan. t.* 376.

R. foliis omnibus capillaceis circumſcriptione oblongis. *Roy. lugdb.* 492.

R. albus fluitans peucedani folio. *Herm. lugdb.* 517.

Millefolium aquaticum foliis foeniculi, ranunculi flore et capitulo. *Bauh. pin.* 141.

Fluſs-Ranunkel. W.

Habitat in fluviis Europae *et* Africae *borealis.* ♃ (v. v.)

In fluviis et lacubus ſemper foliis longiſſimis dichotomis occurrit, nec ullo modo mutatur. W.

1087. TROLLIUS.

Gen. plant. edit. Schreb. n. 954.

Cal. 0. *Petala* circiter 14. *Capſ.* plurimae, ovatae, polyſpermae.

1. TROLLIUS *europaeus.*

T. corollis conniventibus, nectariis longitudine ſtaminum. *Sp. pl.* 782. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 190. *Hall. helv. n.* 1189. *Pall. it.* 3. *p.* 253. *Scop. carn. n.* 694. *Fl. dan. t.* 133. *Kniph. cent.* 4. *n.* 90. *Knorr. del.* 2. *t. T.* 4. *Hoffm. germ.* 198. *Roth. germ. I.* 236. *II.* 608.

T. (*altiſſimus*) flore pedunculato clauſo globoſo. *Crantz. auſtr. p.* 134.

Helleborus flore clauſo erecto petiolato, caule ſimpliciſſimo. *Fl. ſuec.* 474, 510.

Helleborus foliis angulatis multifidis, flore globoſo. *Hort. cliff.* 227. *Roy. lugdb.* 484.

Helleborus caule ſimplici, flore pedunculato. *Fl. lapp.* 226.

Ranunculus montanus, aconiti folio, flore globoſo *Bauh. pin.* 182.

Ranunculus flore globoſo. *Dod. pempt.* 430.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 364.
Europäiſche Trollblume. W.
Habitat in alpibus et ſubalpinis Sueciae, Germaniae, Angliae. ♃ (v. v.)
Corollae *clauſae flavae.* Nectaria *flava, ipſis ſtaminibus non longiora.* Germina *ſ.* Piſtilla *erecta.*

2. TROLLIUS *aſiaticus.*
T. corolla patula, nectariis ſtamine longioribus. *Sp. pl.* 782. *Mill. dict. n.* 2. *Pall. it.* 2. *p.* 528.
Helleborus aconiti folio, flore globoſo croceo. *Amman. ruth.* 101.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 366.
Aſiatiſche Trollblume. W.
Habitat in Sibiria. ♃ (v. v.)
Parum a praecedente diſtinctus nectariis fulvis ſtamine longioribus; Corollis *ſaturate luteis hiantibus.* Germina *ſ.* Piſtilla *extrorſum flexa quaſi procumbentia.*

1088. ISOPYRUM.

Gen. pl. ed. Schreb. n. 955.

Cal. 0. *Petala* 5. *Nectaria* 3-fida, tubulata. *Capſulae* recurvae, polyſpermae.

1. ISOPYRUM *fumarioides.*
I. ſtipulis ſubulatis, petalis acutis. *Sp. pl.* 783. *Hort. upſ.* 157. *Mill. dict. n.* 1. *Gmel. ſib.* 4. *p.* 191. *Willich. obſ. n.* 44. *Kniph. cent.* 12. *n.* 65.
Helleborus fumariae foliis. *Amm. ruth.* 74. *t.* 12.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 367.
Erdrauchartiges Iſopyrum. W.
Habitat in nemoribus Sibiriae. ⊙ (v. v.)
Flores lutei.
Conf. Aquilegiae corniculis, Moſchatellinae foliis, planta puſilla ex inſula cheuſan. *Pluk. amalth.* 2. *p.* 360. *f.* 3.

2. ISOPYRUM *thalictroides.*
I. ſtipulis ovatis, petalis obtuſis. *Sp. pl.* 783. *Scop. carn. n.* 695. *Mill. dict. n.* 2. *Iacq. auſtr. t.* 105. *Mattuſch. ſil. n.* 419. *Crantz. auſtr.* 135. *Hoffm. germ.* 198.
Thalictrum batrachioides, flore albo, italicum. *Bocc. muſ.* 84. *t.* 79. *f.* 1. *

Ranunculus nemoroſus, thalictri folio. *Bauh. pin.* 178. *Raj. hiſt.* 584. *Moriſ. hiſt.* 2. *p.* 437. *f.* 4. *t.* 28. *f.* 12.
Ranunculus praecox II, thalictri folio. *Cluſ. hiſt.* 1. *p.* 233.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 368.
Thalictrumartiges Iſopyrum. W.
Habitat in alpibus Italicis, Gratianopoli, *inque agro* Viennenſi, Carnioliae, *locis umbroſis; vernalis.* ♃ (v. ſ.)
Petala *alba.* Germina 1, 2, 3. *Scop. Crantz.* R.

3. ISOPYRUM *aquilegioides.* †
I. ſtipulis obſoletis. *Sp. pl.* 783. *Mill. dict. n.* 3. *Hall. helv. n.* 1190.
Aquilegia montana, flore parvo, thalictri folio. *Bauh. pin.* 144. *prodr.* 75. *Bauh. hiſt.* 3. *p.* 484. *Raj. hiſt.* 707. *Moriſ. hiſt.* 3. *p.* 458. *ſ.* 12. *t.* 1. *f.* 5.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 369.
Akeleyartiges Iſopyrum. W.
Habitat in alpibus Helveticis, Tridentinis, Apenninis.
Synonymon *Bauhinorum ad* Aquilegiam viſcoſam *amandat de la Chenal in Act. Helv. tom.* 8. *p.* 146. R.
Planta obſcura. W.

1089. HELLEBORUS.

Gen. plant. ed. Schreb. n. 956.

Cal. 0. *Petala* 5 ſ. plura. *Nectaria* bilabiata, tubulata. *Capſ.* polyſpermae, erectiuſculae.

1. HELLEBORUS *hyemalis.*
H. flore folio inſidente, foliis radicalibus peltatis multifidis, laciniis integerrimis. W.
H. flore folio inſidente. *Hort. cliff.* 227. *Hort. upſ.* 158. *Roy. lugdb.* 484. *Hall. helv. n.* 1191. *Mill. dict. n.* 5. *Crantz. auſtr. p.* 133. *Iacq. auſtr. t.* 202. *Blackw. t.* 576. *Kniph. cent.* 1. *n.* 40. *Hoffm. germ.* 198.
H. ranunculoides praecox tuberoſus, flore luteo. *Moriſ. hiſt.* 3. *p.* 459. *ſ.* 12. *t.* 2. *f.* 4.
Aconitum unifolium bulboſum. *Bauh. pin.* 183. *Hill. anat. t.* 11.
Aconitum hyemale. *Camer. epit.* 728. R.
Aconitum luteum minus. *Dodon. pempt.* 352. R.

Houttuyn Lin. Pfl. Syſt. 7. *p.* 370.
Winter-Chriſtwurz. W.
Habitat in Lombardia, Helvetia, Italia, Apenninis, Auſtria. ♃ (v. v.)

Scapus *ſimpliciſſimus, vix pollicaris, monophyllus,* Folio *horizontali, terminali.* Flos *unicus, terminalis, ſeſſilis, erectus, luteus.*

*2. HELLEBORUS *ranunculinus.*

H. caule unifloro folioſo ſimplici, foliis palmatis lobis inciſo-dentatis. W.
H. (*ranunculinus*) floribus ſolitariis pedunculatis, foliis digitato-multifidis inciſis. *Smith. ic. ined.* 1. *p.* 37. *t.* 37.
H. niger orientalis Ranunculi folio, flore nequaquam globoſo. *Tournef. cor.* 20.
Trollius humilis flore patulo. *Buxb. cent.* 1. *tab.* 22.
Ranunkelartige Chriſtwurz. W.
Habitat in Cappadocia. ♃ (v. ſ.)

3. HELLEBORUS *niger.*

H. ſcapo ſubbifloro ſubnudo, foliis pedatis. *Hort. upſ.* 157. *Roy. lugdb.* 484. *Mat. med.* 142. *Mill. dict. n.* 3. *Crantz. auſtr. p.* 133. *n.* 3. *Iacq. auſtr. t.* 201. *Mattuſch. ſil. n.* 420. *Blackw. t.* 506. 507. *Ludw. ect. n.* 139. *Kniph. cent.* 1. *n.* 41. *Knorr. del.* 1. *t. N.* 6. *Hoffm. germ* 199.
H. ſcapo florifero ſubnudo, petiolo communi bipartito. *Hort. cliff.* 227.
H. nectariis obſubulatis, filamentis ſtaminum duplo brevioribus. *Scop. carn. ed.* 1. *p.* 556. *n.* 2. *ed.* 2. *n.* 696.
H. niger, flore roſeo. *Bauh. pin.* 186. *Hill. anat. t.* 1.
Elleborus niger, legitimus. *Cluſ. hiſt.* 1. *p.* 275.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 371.
Schwarze Chriſtwurz. W.
Habitat in Auſtriae, Hetruriae, Apennini *aſperis.* ♃ (v. v.)

Folia conſiſtentia, hyeme perennantia.

Folia *omnia radicalia, coriacea.* Scapus *rudimento, unius folii, aphyllus, bifidus; ramis aphyllis, bifloris bracteatis.* Flores *verticillares, albi.*

4. HELLEBORUS *viridis.*

H. caule bifido, ramis folioſis bifloris, foliis digitatis,

Mill. dict. n. 2. Grim. iſen. in Nov. Act. A. N. C. tom. 3. app. p. 329. Jacq. auſtr. t. 106. Mattuſch. ſil. n. 421. Doerr. naſſ. p. 123. Blackw. t. 509. 510. Kniph. cent. 1. n. 42. Knorr. del. 1. t. N. 5. Hoffm. germ. 199. Roth. germ. I. 234. II. 600.

H. caule aequali folioſo, foliis radicalibus caulem tandem ſuperantibus. *Hort. cliff. 227. Hort. upſ. 158. Roy. lugdb. 484.*

H. foliis multipartitis, ſerratis, caule paucifloro. *Hall. helv. n. 1192.*

H. foliis digitatis, flore viridi. *Crantz. auſtr. p. 134.*

H. nectariis obconicis, filamentis ſtaminum triplo brevioribus. *Scop. carn. p. 556. n. 1. ed. 2. n. 697.*

H. niger hortenſis, flore viridi. *Bauh. pin. 185.*

Elleborum nigrum alterum. *Cam. epit. 941.*

Houttuyn Lin. Pfl. Syſt. 7. p. 373.

Grüne Chriſtwurz. W.

Habitat in montibus Vignenſibus, Euganeis, rel. ♃ (v. v.)

Folia *omnia radicalia.* Scapus *aphyllus bifidus: Ramis folioſis, bifloris* Fores *nutantes, virides.* Piſtilla 3, *raro* 4, *rariſſime* 5. *Crantz. l. c.* R.

*5. HELLEBORUS *orientalis.*

H. caule multifloro, foliis pedatis ſubtus hirtis. W.

H. caule ſuperne diviſo folioſo et multifloro, foliis duplo altiore, foliis amplis pedato-digitatis ſubtus pubeſcentibus. *Lamarck. encycl. 3. p. 92.* *

H. niger orientalis ampliſſimo folio, caule praealto, flore purpuraſcente. *Tournef. cor. 20.*

Wahre Chriſtwurz. W.

Habitat in Oriente. ♃ (v. ſ.)

Eſt verus Helleborus Hippocratis. Similis quodammodo H. viridi, differt vero: caule *ramoſo multifloro,* corolla *viridi-purpuraſcente,* foliolis *foliorum radicalium ellipticis craſſioribus ſubtus ad venas prominentes pilis ſparſis obſitis. Ab H. foetido diverſus* foliis *radicalibus ſubtus hirtis multo majoribus et foliis floralibus trilobis apice dentatis.* W.

6. HELLEBORUS *foetidus.*

H. caule multifloro folioſo, foliis pedatis. *Sp. pl 784. Hoffm. germ. 199. Roth. germ. I. 234. II. 600.*

H. caule inferne anguſtato-multifolio multifloro; foliis

caule brevioribus. *Hort. cliff.* 227. *Roy. lugdb.* 484. *Dalib. pariſ.* 169. *Sauv. monſp.* 180. *Mill. dict. n.* 1. *Pollich. pal. n.* 540. *Doerr. naſſ. p.* 123. *Blackw. t.* 57. *Kniph. cent.* 12. *n.* 52. 53. *Regn. bot.*
H. ramoſus, multiflorus, foliis multipartitis ſerratis, ſtipulis ovato-lanceolatis, coloratis. *Hall. helv. n.* 1193.
H. niger foetidus. *Bauh. pin.* 185.
Helleboraſter maximus. *Lob. ic.* 679.
H. foemina. *Sterb. fung.* 372. *t.* 36. *f. C.*
Houttuyn Lin. Pfl. Syſt. 7. *p.* 374.
Stinkende Chriſtwurz.
Habitat in Germania, Helvetia, Gallia. ♃ (v. v.)
Folia *omnia in caule; verſus radicem nulla.*
Folia *floralia ovata integerrima.* W.

*7. HELLEBORUS *lividus.*
H. caule multifloro folioſo, foliis ternatis. *Ait. Kew* 2. *p.* 272. *Curt. mag.* 72.
Helleborus foetidus β. *Sp. pl.* 784.
H. niger trifoliatus. *Aldini hort. farneſ.* 93. *t.* 92. *Moriſ. hiſt.* 3. *p.* 460. *ſ.* 12. *t.* 4. *f.* 7.
Schmutzigweiſse Chriſtwurz. W.
Habitat - - - - ♃ (v. ſ.)

8. HELLEBORUS *trifolius.*
H. ſcapo unifloro, foliis ternatis. *Amoen. acad.* 2. *p.* 356. *t.* 4. *f.* 18. *Kalm. it.* 3. *p.* 379. *Lepech. it.* 1. *p.* 190. *Pall. itin.* 3. *p.* 34. *Flor. dan. t.* 566.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 376.
Kleinſte Chriſtwurz. W.
Habitat in Canadae, Sibiriae *ſylvis nemoroſis cum Oxalide, Circaea, etiam in* Islandia. ♃ (v. ſ.)
Scapus *aphyllus.* Flos *minutus, Parnaſſiae ſimilis, albus.*

1090. CALTHA.

Gen. plant. edit. Schreb. n. 957.

Cal. 0. *Petala* 5. *Nectaria* 0. *Capſ.* plures, polyſpermae.

1. CALTHA *paluſtris.*
C. caule erecto. W.

Caltha. *Fl. lapp.* 227. *Fl. ſuec.* 473, 511. *Hort. cliff.* 228. *Roy. lugdb.* 484. *Clayt. virg.* 166. *Dalib. pariſ.* 169. *Gmel. ſib.* 4. *p.* 191. *Hall. helv. n.* 1188. *Crantz. auſtr. p.* 136. *Pollich. pal. n.* 541. *Fl. dan. t.* 668. *Kniph. cent.* 7. *n.* 7. *Knorr. del.* 1. *t. H.* 2. *Hoffm. germ.* 199. *Roth. germ. I.* 234. *II.* 601.
C. paluſtris, flore ſimplici. *Bauh. pin.* 276.
C. (*major*) foliis orbiculatis crenatis, flore majore. *Mill. dict. n.* 1.
Populago paluſtris. *Scop. carn. ed.* 2. *n.* 698.
Populago major. *Tabern. ic.* 750.
β. Populago minor. *Tabern. ic.* 750.
C. (*minor*) foliis orbiculato-cordatis crenatis, flore minore. *Mill. dict. n.* 2.
γ. Caltha paluſtris, flore pleno. *Bauh. pin.* 276. *Tabern.* 751.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 377.
Gemeine Kuhblume. W.
Habitat in humidiuſculis Europae, Aſiae *et* Americae *borealis.* ♃ (v. v.)

*2. CALTHA *natans.* †
C. caule procumbente natante. W.
Caltha natans. *Pall. itin. ed. min.* 3. *p.* 248.
C. caulibus procumbentibus, foliis cordato-reniformibus aequaliter crenatis, floribus puſillis. *Gmel. ſib.* 4. *p.* 192. *t.* 82. *
Populago flore minore rubello. *Gmel. index Ircut.* 605.
Schwimmende Kuhblume. W.
Habitat in aquis ſtagnantibus Sibiriae *circa urbem* Ircutiam. ♃
Habitus praecedentis ſed omnibus partibus minor, differt: caule *procumbente natante,* petalis *ſubrotundis albis margine rubicundis, externe vireſcentibus,* capſulis *triquetris.* W.

1091. HYDRASTIS.

Gen. pl. ed. Schreb. n. 958.

Cal. 0. *Petala* 3. *Nectaria* 0. *Bacca* compoſita acinis monoſpermis.

1. HYDRASTIS *canadenſis.*
H. canadenſis. *Sp. pl.* 784.

Warnera canadenſis. *Mill. ic.* 190. *t.* 285.
Hydrophyllum verum canadenſium. *Sp. pl.* 1. *p.* 146.
Houttuyn Lin. Pfl. Syſt. 7. *p.* 379.
Canadiſche Hydraſtis. W.
Habitat in Canadae *aquis.* ♃ (v. ſ.)

Simillis Hydrophyllo. Folia *bina, petiolata, baſi emarginata, palmata, ſerrata: lobis utrinque lobulo laterali.*
